

ISSN 2146-1147

*TWMS Journal of  
Applied  
and Engineering  
Mathematics*

Volume 13, No.1, 2023

**Turkic  
World  
Mathematical  
Society**



İŞIK UNIVERSITY  
<http://jcem.iskun.edu.tr>

# World Mathematical Society Journal of Applied and Engineering Mathematics



[Home](#)

[Focus](#)

[Author Guidelines](#)

[Current](#)

[Archive](#)


[Indexing](#)

[Publication Ethics](#)

[Subscribe](#)

[Search](#)

[Announcements](#)





**Turkic  
World  
Mathematical  
Society**



Siddik B. Yarman (Istanbul, TURKEY)

## Digital Repository Universitas Jember

Editor in Chief:

Elman Hasanoglu (Istanbul, TURKEY)

Managing Editor:

Selcuk Akbulut (Istanbul, TURKEY)

Associate Editors:

Banu Uzun (Istanbul, TURKEY)

Georgi Georgiev (V.Tarnovo, BULGARIA)

Serkan Sütü (Istanbul, TURKEY)

Contributing Editors:

Mert Savcin (Istanbul, TURKEY)

Publisher:

FEYZİYE SCHOOLS FOUNDATION

IŞIK UNIVERSITY

Faculty of Engineering and Natural Sciences

Department of Mathematics

Library of Congress USA catalogue QA1.T885

Indexed and abstracted by:

1. Emerging Sources Citation Index (ESCI) by Web of Science;



2. SCOPUS;

# Digital Repository Universitas Jember

3. EBSCO;

4. VINITI (RUSSIA);

5. COSMOSIMPACT FACTOR and by others.



This Journal is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

## ABOUT THE JOURNAL

### People

- [Editorial Board](#)


### Policies

- [Focus and Scope](#)
- [Open Access Policy](#)

### Submissions

- [Author Guidelines](#)
- [Copyright Notice](#)
- [Contact](#)

### Editorial Board



Top

Ugur	Abdulla	(Florida, USA)	<a href="#"><u>www</u></a>
Ravi P.	Agarwal	(Madras, India)	<a href="#"><u>www</u></a>
Hijaz	Ahmad	(Rome, Italy)	<a href="#"><u>www</u></a>
Farajzadeh	Ali	Kermanshah, Iran	<a href="#"><u>www</u></a>
Fikret	Aliev	(Baku, Azerbaijan)	<a href="#"><u>www</u></a>

Allaberen	Ashyralyev	(Ashgabat, Turkmenistan)	<a href="#">www</a>
Agamirza	Bashirov	(Gazimagusa, Cyprus)	<a href="#">www</a>
Ismihan	Bayramoglu	(Izmir, Turkey)	<a href="#">www</a>
İsmail Naci	Cangül	(Bursa, Turkey)	<a href="#">www</a>
Ahmet Sinan	Çevik	(Konya, Turkey)	<a href="#">www</a>
Can F.	Delale	(Istanbul, Turkey)	<a href="#">www</a>
Sadik	Dost	(Victoria, Canada)	<a href="#">www</a>
Marcelo	Epstein	(Calgary, Canada)	<a href="#">www</a>
Manish	Jain	(Rewari, India)	<a href="#">www</a>
Varga	Kalantarov	(Istanbul, Turkey)	<a href="#">www</a>
Bulent	Karasozen	(Ankara, Turkey)	<a href="#">www</a>
Kazuya	Kobayashi	(Tokyo, Japan)	<a href="#">www</a>
Nazim	Mahmudov	(Gazimagusa, Cyprus)	<a href="#">www</a>
Agarwal	Praveen	(Jaipur, India)	<a href="#">www</a>
Sunil	Purohit	(Kota, India)	<a href="#">www</a>
Soheil	Salahshour	(Istanbul, Turkey)	<a href="#">www</a>
Hari Mohan	Srivastava	(Victoria, Canada)	<a href="#">www</a>
Eldar	Veliev	(Kharkov, Ukraine)	<a href="#">www</a>
Gu	Xian-Ming	(Sichuan, China)	<a href="#">www</a>
Tolga	Yarman	(Istanbul, Turkey)	<a href="#">www</a>
Kewen	Zhao	(Sanya, China)	<a href="#">www</a>

## Focus and Scope

[Top](#) ↑

TWMS Journal of Applied and Engineering Mathematics (TWMS J. of Apl. & Eng. Math.) is an international open-access, peer-reviewed, four times a year (January, April, July, October) journal which is published on-line. The Journal publishes full-length original articles, reviews on specialized topics, technical notes, book reviews in the field of applied and interdisciplinary mathematics and is an ideal academic platform for researchers to share their latest research findings and directions in applied and engineering mathematics. The scope of the Journal covers, but not limited to the following fields:

- applied probability and statistics,
- approximation theory
- applied functional analysis, linear algebra, discrete mathematics, and matrix computations,

- applied electromagnetic theory and wave motion,
- combinatorics,
- control theory,
- financial mathematics,
- fuzzy sets and logic,
- game theory,
- graph theory,
- information theory,
- inverse problems,
- mathematical economy,
- nonlinear processes in physics,
- numerical analysis and numerical solutions of ordinary and partial differential equations,
- operation research.

## Open Access Policy



TWMS Journal of Applied and Engineering Mathematics is an **Open Access** journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author. This is in accordance with the BOAI definition of Open Access

## Author Guidelines



Before submitting a manuscript it's strongly recommended that the authors read **Publication Ethics**.

Manuscripts submitted to the "TWMS Journal of Applied and Engineering Mathematics" should be in any common version of TeX including LaTeX 2.09, MiK TeX and be written strictly according to the journal's standards. For the journal template, please, click [TeX file Template](#).

The text of a manuscript should contain/include:

- a short abstract (no more than 120 words and no less than 100 words),
- keywords,
- AMS Subject Classification, 2010,
- introduction and conclusion,
- all necessary figures and tables with titles, description, and footnotes (permissions must be obtained for use of copyrighted material),
- a reference list.

Manuscripts, which do not meet these conditions or is written in other format than the template of the journal will not

be accepted for further evaluation.

## Digital Repository Universitas Jember

Submitted papers should not have been published previously or be under consideration for publication in any other journal or book available through a library or by purchase.

Authors of articles are not restricted to being members of any particular institute, society, or association. Articles based on theses for higher degrees may be submitted, as may articles presented at conferences, provided these articles do not appear in substantially the same form in published conference proceedings.

All authors should provide their ORCID ID. (For more information about ORCID and registration for a *free* ORCID identifier please visit **ORCID.org** ).

The total page of a manuscript should not exceed 15 pages.

Publication in the journal is free (open or hidden) from any kind of charge.

We encourage you to send your paper in the electronic form so it can be refereed without postal delays, and be published more quickly.

The average time for evaluation is about two months. However, authors should note that each submission is assessed on its individual merit and in certain circumstances estimating time may differ.

Please, send both PDF and LaTeX version of your paper to [jaem@isikun.edu.tr](mailto:jaem@isikun.edu.tr) with the subject "Electronic submission for TWMS J. App. & Eng. Math." in the subject line.

---

### Copyright © Notice

Copyright © 2011 by Isik University.

Top ↑

---

### Contact

Elman Hasanoglu,  
Isik University, Department of Mathematics,  
34980, Sile/Istanbul/TURKEY  
e-mail: [jaem@isikun.edu.tr](mailto:jaem@isikun.edu.tr)

Top ↑



# World Mathematical Society Journal of Applied and Engineering Mathematics



- [Home](#)
- [Focus](#)
- [Author Guidelines](#)
- [Current](#)
- [Archive](#)
- [Indexing](#)
- [Publication Ethics](#)
- [Subscribe](#)
- [Search](#)
- [Announcements](#)


**Jaem Vol 13, No 1, 2023**

**TWMS J. App. & Eng. Math.**

	COVER	[JPG]
<u>LOCAL DISTANCE IRREGULAR LABELING OF GRAPHS</u> <i>A. I. Kristiana, R. Alfarisi, Dafik</i>		1 [PDF]
<u>K-PRODUCT CORDIAL LABELING OF FAN GRAPHS</u> <i>K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi, M. Z. Youssef</i>		11 [PDF]
<u>ON SOLUTION OF BG-VOLTERRA INTEGRAL EQUATIONS</u>		21

<u>SOME FIXED POINT RESULTS FOR <math>\beta</math>-ADMISSIBLE MULTI-VALUED F-CONTRACTIONS</u> <i>E. Nazari</i>	37 [PDF]
<u>CERTAIN SUBCLASSES OF ANALYTIC FUNCTION BY SALAGEAN <math>q</math>-DIFFERENTIAL OPERATOR</u> <i>Dileep L., Divya Rashmi S. V.</i>	46 [PDF]
<u>PATHOS DEGREE PRIME GRAPH OF A TREE</u> <i>H. M. Nagesh</i>	53 [PDF]
<u>BAYESIAN ESTIMATION USING LINDLEY'S APPROXIMATION OF INVERTED KUMARASWAMY DISTRIBUTION BASED ON LOWER RECORD VALUES</u> <i>Sana, M. Faizan, A. A. Khan</i>	65 [PDF]
<u>AN ITERATIVE METHOD FOR SOLVING TIME-FRACTIONAL PARTIAL DIFFERENTIAL EQUATIONS WITH PROPORTIONAL DELAYS</u> <i>B. Mallick, P. K. Sahu, M. Routaray</i>	74 [PDF]
<u>NUMERICAL OPTIMIZATION ALGORITHM BASED ON GENETIC ALGORITHM FOR A DATA COMPLETION PROBLEM</u> <i>B. Joulik, J. Daoudi, C. Tajani, J. Abouchabaka</i>	86 [PDF]
<u>SPHERICAL FUZZY MATRICES</u> <i>I. Silambarasan</i>	98 [PDF]
<u>DETERMINATION OF A TIME-DEPENDENT COEFFICIENT IN THE TIME-FRACTIONAL WAVE EQUATION WITH A NON-CLASSICAL BOUNDARY CONDITION</u> <i>I. Tekin, H. Yang</i>	110 [PDF]
<u>A PIECEWISE ANALYTICAL ITERATIVE METHOD FOR A WIRE-MASS MODEL</u> <i>S. Mungaski</i>	124 [PDF]
<u>CONTRACTION AND DOMINATION IN FUZZY GRAPHS</u> <i>S. Ramya, S. Lavanyag</i>	133 [PDF]
<u>REGULARIZED TRACE ON SEPARABLE BANACH SPACES</u> <i>E. Gül, T. L. Gill</i>	143 [PDF]
<u>APPLYING MULTIQUADRIC QUASI-INTERPOLATION TO SOLVE FOKKER-PLANCK EQUATION</u> <i>M. Rahimi, H. Adibi, M. Amirfakhrian</i>	152 [PDF]
<u>A GENERALIZATION OF RELATION-THEORETIC CONTRACTION PRINCIPLE</u> <i>D. Khantwal, S. Aneja, G. Prasad, U. C. Gairola</i>	166 [PDF]
<u>SEVERAL TYPES OF SINGLE-VALUED NEUTROSOPHIC IDEALS AND FILTERS ON A LATTICE</u> <i>A. Bennoui, L. Zedam, S. Milles</i>	175 [PDF]
<u>RIESZ MRA OF DYADIC DILATIONS AND THE CORRESPONDING RIESZ WAVELET ON LCA GROUPS</u> <i>R. Kumar, Satyapriya, M. Singh</i>	189 [PDF]
<u>SPECTRAL INCLUSION BETWEEN A REGULARIZED QUASI-SEMIGROUPS AND THEIR GENERATORS</u> <i>A. Tajmouati, Y. Zahouan</i>	202 [PDF]
<u>AN INTRODUCTION TO PYTHAGOREAN FUZZY HYPERIDEALS IN HYPERSEMIGROUPS</u> <i>V. S. Subha, S. Sharmila</i>	214 [PDF]
<u>FUNCTIONAL VARIABLE METHOD TO THE CHIRAL NONLINEAR SCHRODINGER EQUATION</u> <i>A. Neiramehi</i>	225 [PDF]
<u>ON THE VERTEX DEGREE POLYNOMIAL OF GRAPHS</u> <i>H. Ahmed, A. Alwardi, R. Salestina M.</i>	232 [PDF]
<u>SOME NEW EXISTENCE RESULTS FOR BOUNDARY VALUE PROBLEMS INVOLVING <math>\beta</math>-CAPUTO FRACTIONAL DERIVATIVE</u> <i>H. Afshari, M. S. Abdo, M. N. Sahlan</i>	246 [PDF]
<u>THE HUB NUMBER OF A FUZZY GRAPH</u> <i>Haifa A. A., M. M. Q. Shubatah</i>	256 [PDF]
<u>OPTIMAL CONTROL ANALYSIS OF THE MALARIA MODEL WITH SATURATED</u>	265

<u>TREATMENT</u> <i>A. K. Srivastav, M. Ghosh</i>	[PDF]
<u>A NEW CLASS OF NON NORMAL OPERATORS ON HILBERT SPACES</u> <i>A. N. Bakir</i>	276 [PDF]
<u>ENCRYPTION THROUGH SQUARE GRID AUTOMATA</u> <i>F. R. P. Mary, L. Shobana, R. Sujatha</i>	285 [PDF]
<u>SOLUTION OF THE VOLTERRA-FREDHOLM INTEGRAL EQUATIONS VIA THE BERNSTEIN POLYNOMIALS AND LEAST SQUARES APPROACH</u> <i>N. Negarchi, K. Nouri</i>	291 [PDF]
<u>FORMALLY SELF-DUAL CODES OVER <math>\mathbb{Z}_2</math></u> <i>Z. Ö. Özger, B. Yildiz</i>	300 [PDF]
<u>DEGREE BASED TOPOLOGICAL INDICES OF LINE GRAPH OF A CAYLEY TREE <math>\Gamma_n^k</math></u> <i>U. Poojari, B. S. Durgi, R. M. Banakar</i>	309 [PDF]
<u>NOVEL POSSIBILITY PYTHAGOREAN INTERVAL VALUED FUZZY SOFT SET METHOD FOR A DECISION MAKING</u> <i>M. Palanikumar, K. Arulmozhi</i>	327 [PDF]
<u>A TECHNIQUE FOR SOLVING SYSTEM OF GENERALIZED EMDEN-FOWLER EQUATION USING LEGENDRE WAVELET</u> <i>A. K. Barnwal, N. Sriwastav</i>	341 [PDF]
<u>APPLYING VIM TO CONFORMABLE PARTIAL DIFFERENTIAL EQUATIONS</u> <i>A. Harir, S. Melliani, L. S. Chadli</i>	362 [PDF]
<u>PREDICTING THE OCEAN CURRENTS USING DEEP LEARNING</u> <i>C. Bayındır</i>	373 [PDF]
<u>SD-PRIME CORDIAL LABELING OF SUBDIVISION <math>K_4</math>-SNAKE AND RELATED GRAPHS</u> <i>U. M. Prajapati, A. V. Vantiya</i>	386 [PDF]



## LOCAL DISTANCE IRREGULAR LABELING OF GRAPHS

A. I. KRISTIANA<sup>1</sup>, R. ALFARISI<sup>2\*</sup>, DAFIK<sup>1</sup>, §

**ABSTRACT.** We introduce the notion of distance irregular labeling, called the local distance irregular labeling. We define  $\lambda : V(G) \rightarrow \{1, 2, \dots, k\}$  such that the weight calculated at the vertices induces a vertex coloring if  $w(u) \neq w(v)$  for any edge  $uv$ . The weight of a vertex  $u \in V(G)$  is defined as the sum of the labels of all vertices adjacent to  $u$  (distance 1 from  $u$ ), that is  $w(u) = \sum_{y \in N(u)} \lambda(y)$ . The minimum cardinality of the largest label over all such irregular assignment is called the local distance irregularity strength, denoted by  $dis_l(G)$ . In this paper, we found the lower bound of the local distance irregularity strength of graphs  $G$  and also exact values of some classes of graphs namely path, cycle, star graph, complete graph,  $(n, m)$ -tadpole graph, unicycle with two pendant, binary tree graph, complete bipartite graphs, sun graph.

**Keywords:** Distance irregularity labeling, local distance irregularity strength, some families graph.

**AMS Subject Classification:** 05C78.

### 1. INTRODUCTION

We consider that all graphs in this paper are finite, simple and connected graph, for detail definition of graph see [6, 2]. There are many types of graph labeling in Gallian [10]. One of graph labeling is magic labeling that was introduced by Sedlacek [11]. Suppose  $G$  is magic if the edges of  $G$  can be labeled by  $1, 2, \dots, |E(G)|$  so that the sum of the labels of all the edges incident with any vertex is the same. Hartsfield and Ringel [7] introduced the concept of antimagic labeling of a graph. There are a lot of results regarding to antimagic labeling, some of them can be found in Dafik *et. al* [5, 3, 4]. They determined super edge-antimagic total labelings of  $mK_{n,n}$  and super edge-antimagicness for a class of disconnected graphs, respectively. Furthermore, Bača *et. al* in [1, 9] found the antimagicness of disjoint union of isomorphism graphs.

The distance  $d(u, v)$  is the minimum of the length of the  $u - v$  path of  $G$ . For a connected graph  $G$  of diameter  $k$ , a distance- $d$  graph  $G_d$  for  $d = 1, 2, \dots, k$  is a graph with the same vertex set  $V(G)$  and the edge set consists of the pairs of vertices that lie at distance  $d$  apart. Slamir [8] was introduced the distance irregular labeling of graphs. In this labeling, the weight of a vertices in  $G$ , is the sum of the labels of all vertices adjacent to  $u$  (distance 1 from  $u$ ), that is  $w(u) = \sum_{y \in N(u)} \lambda(y)$ . The distance irregular labeling of  $G$ , denoted by  $dis(G)$ , is the minimum cardinality of the largest label  $k$  over all such irregular assignment.

<sup>1</sup> Department of Mathematics Education, University of Jember, Indonesia.  
e-mail: arika.fkip@unej.ac.id; ORCID: <http://orcid.org/0000-0002-6556-6377>.  
e-mail: d.dafik@unej.ac.id; ORCID: <https://orcid.org/0000-0003-0575-3039>.

<sup>2</sup> Department of Primary School, University of Jember, Indonesia.  
e-mail: alfarisi.fkip@unej.ac.id; ORCID: <http://orcid.org/0000-0003-3011-4917>.

\* Corresponding author.

§ Manuscript received: April 30, 2020; accepted: October 22, 2020.

TWMS Journal of Applied and Engineering Mathematics, Vol.13, No.1 © Işık University, Department of Mathematics, 2023; all rights reserved.

The coloring problem started the four-color problem. We define a coloring of  $G$  to be an assignment of colors to the vertices of  $G$  such that no two adjacent vertices have the same color. On the other hand, the chromatic number has been related with several parameters of graphs, and as a consequence, there exists different types of vertex coloring such as list coloring, total coloring, acyclic coloring, distance- $k$  coloring, etc.

**Observation 1.1.** [8] *Let  $u$  and  $w$  be any two adjacent vertices in a connected graph  $G$ . If  $N(u) - \{w\} = N(w) - \{u\}$ , then the labels of  $u$  and  $w$  must be distinct, that is  $\lambda(u) \neq \lambda(w)$ .*

## 2. RESULTS

In this paper, we combine the concept of distance irregular labeling and coloring problem. We introduce the notion of distance irregular labeling, called the local distance irregular labeling.

**Definition 2.1.** *Suppose  $\lambda : V(G) \rightarrow \{1, 2, \dots, k\}$  such that the weight calculated at the vertices induces a vertex coloring if  $w(u) \neq w(v)$  for any edge  $uv$ . The weight of a vertex  $u \in V(G)$  is defined as the sum of the labels of all vertices adjacent to  $u$  (distance 1 from  $u$ ), that is  $w(u) = \sum_{y \in N(u)} \lambda(y)$ .*

**Definition 2.2.** *The minimum cardinality of the largest label over all such irregular assignment is called the local distance irregularity strength, denoted by  $dis_l(G)$ .*

In this paper, we find the new concept of distance irregular labeling namely local distance irregular labeling. We have found the lower bound of local distance irregular labeling and also determine the exact values of local distance irregular labeling of some classes graphs path, cycle, star graph, complete graph,  $(n, m)$ -tadpole graph, unicycle with two pendant, binary tree graph, complete bipartite graphs, sun graph in the following theorems.

**Lemma 2.1.** *Let  $G$  be a connected graph on  $n \geq 4$  vertices with the chromatic number  $\chi$ , the minimum degree  $\delta$  and the maximum degree  $\Delta$  and there is no vertex having identical neighbours, then we have the lower bound of the local distance irregular labeling of  $G$  is*

$$dis_l(G) \geq \left\lceil \frac{\chi + \delta - 1}{\Delta} \right\rceil$$

*Proof.* The smallest weight possible of vertices in  $G$  is  $\delta$ . Since, the weight of vertices in  $G$  induces vertex coloring and there are  $n$  vertices, then the largest weight of vertices in  $G$  is at least  $\chi + \delta - 1$ . This weight obtained from the sum of at most the maximum degree  $\Delta$ . Thus, the largest label is at least  $\left\lceil \frac{\chi + \delta - 1}{\Delta} \right\rceil$ .  $\square$

**Theorem 2.1.** *Let  $P_n$  be a path graph with order  $n \geq 3$ , then we have the local distance irregular labeling of  $P_n$  is*

$$dis_l(P_n) = \begin{cases} 1, & \text{if } n = 3 \\ 2, & \text{if } n \geq 4 \end{cases}$$

*Proof.* The path  $P_n$  is a tree graph with  $n$  vertices. The vertex set  $V(P_n) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(P_n) = \{v_i v_{i+1}; 1 \leq i \leq n - 1\}$ . We have  $\Delta(P_n) = 2$ ,  $\delta(P_n) = 1$  and  $\chi(P_n) = 2$ . The cardinality of vertex set and edge set, respectively are  $|V(P_n)| = n$  and  $|E(P_n)| = n - 1$ . This proof divided into two cases are as follows.

**Case 1:** For  $n = 3$ , we give the label of the vertices  $\lambda(v_1) = \lambda(v_2) = \lambda(v_3) = 1$ . Then, we have the weight of vertices in  $P_3$  as follows

$$w(v_1) = w(v_3) = 1 \qquad w(v_2) = 2$$

It is clear that the labeling provide different weight of two adjacent vertices such that we obtain  $dis_l(P_3) \leq 1$ . It concludes that  $dis_l(P_3) = 1$ .

**Case 2:** For  $n \geq 4$ , Based on Lemma 2.1 that the lower bound of local distance irregular labeling of path graph  $P_n$  is  $dis_l(P_n) \geq \left\lceil \frac{\chi + \delta - 1}{\Delta} \right\rceil = \left\lceil \frac{2 + 1 - 1}{2} \right\rceil = 1$ . However, we can not attain the sharpest lower bound. If all vertices assigned label 1, then there is at least two adjacent vertices which have the same weight namely  $w(v_3) = \lambda(v_2) + \lambda(v_4) = 1 + 1 = 2$  and  $w(v_4) = \lambda(v_3) + \lambda(v_5) = 1 + 1 = 2$

so that  $w(v_3) = w(v_4)$  for edge  $v_3v_4 \in E(P_n)$ . Hence, the lower bound of the distance irregular coloring of  $P_n$  is  $dis_l(P_n) \geq 2$ . Furthermore, we prove that the upper bound of the local distance irregular labeling of  $P_n$  is  $dis_l(P_n) \leq 2$ . Suppose the label vertices for  $n$  is even using the formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = v_1 \text{ and } v = v_n \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2) \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2) \end{cases}$$

The label vertices for  $n \equiv 3(\text{mod}4)$  using formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_{n-2} \text{ and } v = v_i, i \equiv 1, 2, 3(\text{mod}4), 1 \leq i \leq n-3 \\ 2, & \text{if } v = v_{n-1}, v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}4), 1 \leq i \leq n-3 \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = v_1 \\ 2, & \text{if } v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}2), 2 \leq i \leq n-3 \\ 3, & \text{if } v = v_{n-1} \text{ and } v = v_i, i \equiv 1(\text{mod}2), 2 \leq i \leq n-3 \\ 4, & \text{if } v = v_{n-2} \end{cases}$$

The label vertices for  $n \equiv 1(\text{mod}4)$  and  $n \neq 5$  using formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4), i \neq n-3 \\ 2, & \text{if } v = v_{n-3}, v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = v_1 \\ 2, & \text{if } v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}2), 2 \leq i \leq n-3 \\ 3, & \text{if } v = v_{n-1} \text{ and } v = v_i, i \equiv 1(\text{mod}2), 2 \leq i \leq n-5 \\ 4, & \text{if } v = v_{n-2} \text{ and } v = v_{n-4} \end{cases}$$

It is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $P_n$  is  $dis_l(P_n) \leq 2$ . It concludes that  $dis_l(P_n) = 2$ .  $\square$

**Theorem 2.2.** *Let  $C_n$  be a cycle graph with order  $n \geq 4$ , then we have the local distance irregular labeling of  $C_n$  is*

$$dis_l(C_n) = \begin{cases} 2, & \text{if } n \text{ is even} \\ 3, & \text{if } n \text{ is odd} \end{cases}$$

*Proof.* The cycle  $C_n$  is a connected graph with  $n$  vertices. The vertex set  $V(C_n) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(C_n) = \{v_1v_n, v_iv_{i+1}; 1 \leq i \leq n-1\}$ . We have  $\Delta(C_n) = 2$ ,  $\delta(C_n) = 2$  and  $\chi(C_n) = 2, n \text{ even}; \chi(C_n) = 3, n \text{ odd}$ . The cardinality of vertex set and edge set, respectively are  $|V(C_n)| = n$  and  $|E(C_n)| = n$ . This proof divided into two cases are as follows.

**Case 1:** For  $n$  is even

Based on Lemma 2.1 that the lower bound of local distance irregular labeling of cycle graph  $C_n$  is  $dis_l(C_n) \geq \lceil \frac{\chi + \delta - 1}{\Delta} \rceil = \lceil \frac{2+2-1}{2} \rceil = 2$ . However, we attain the sharpest lower bound. Furthermore, we prove that the upper bound of the local distance irregular labeling of  $C_n$  is  $dis_l(C_n) \leq 2$ . Suppose the label vertices for  $n \equiv 0(\text{mod}4)$  is even using the formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2) \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2) \end{cases}$$

The label vertices for  $n \equiv 2(\text{mod}4)$  using formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4), i \neq n \\ 2, & \text{if } v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2) \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2), i \neq n - 1 \\ 4, & \text{if } v = v_{n-1} \end{cases}$$

It is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $C_n$  is  $dis_l(C_n) \leq 2$ . It concludes that  $dis_l(C_n) = 2$ .

**Case 2:**For  $n$  is odd.

Based on Lemma 2.1 that the lower bound of local distance irregular labeling of cycle graph  $C_n$  is  $dis_l(C_n) \geq \lceil \frac{\chi + \delta - 1}{\Delta} \rceil = \lceil \frac{3+2-1}{2} \rceil = 2$ . However, we can not attain the sharpest lower bound. Suppose to contrary that  $C_n$  has 2 label vertices then there exists some possible value as follows

- For  $n \equiv 1(\text{mod}4)$ , We may be construct the label vertices with a periodical irregular assignment  $1, 1, 1, 2, 1, 1, 1, 2, \dots, 1, 1, 1, 2$  of path  $P_{n-1}$  except the last vertices assigned with 1 or 2. If the last vertices assigned with 1, then there is at least two adjacent vertices which have same weight namely  $w(v_1) = w(v_2) = 2$ . If the last vertices assigned with 2, then there is at least two adjacent vertices which have same weight namely  $w(v_n) = w(v_{n-1}) = 3$ . it is a contradict.
- For  $n \equiv 3(\text{mod}4)$ , We may be construct the label vertices with a periodical irregular assignment  $1, 1, 1, 2, 1, 1, 1, 2, \dots, 1, 1, 1, 2, 1, 1$  of path  $P_{n-1}$  except the last vertices assigned with 1 or 2. If the last vertices assigned with 1, then there is at least two adjacent vertices which have same weight namely  $w(v_n) = w(v_{n-1}) = 2$ . If the last vertices assigned with 2, then there is at least two adjacent vertices which have same weight namely  $w(v_{n-2}) = w(v_{n-1}) = 3$ . it is a contradict.

Hence, the lower bound of the local distance irregular labeling of  $C_n$  is  $dis_l(C_n) \geq 3$ . Furthermore, we prove that the upper bound of the local distance irregular labeling of  $C_n$  is  $dis_l(C_n) \leq 3$ . Suppose the label vertices for  $n \equiv 1(\text{mod}4)$  is even using the formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4), i \neq n \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}4) \\ 3, & \text{if } v = v_n \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2), 2 \leq i \leq n - 2 \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2), 2 \leq i \leq n \\ 4, & \text{if } v = v_1 \text{ and } v = v_{n-1} \end{cases}$$

The label vertices for  $n \equiv 3(\text{mod}4)$  using formula

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4), i \neq n \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}4) \\ 3, & \text{if } v = v_n \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2, & \text{if } v = v_n \text{ and } v = v_i, i \equiv 0(\text{mod}2), 2 \leq i \leq n - 2 \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2), 2 \leq i \leq n - 1 \\ 4, & \text{if } v = v_1 \text{ and } v = v_{n-1} \end{cases}$$

It is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $C_n$  is  $dis_l(C_n) \leq 3$ . It concludes that  $dis_l(C_n) = 3$ . □

**Theorem 2.3.** Let  $K_n$  be a complete graph with order  $n \geq 3$ , then we have the local distance irregular labeling of  $K_n$  is  $dis_l(K_n) = n$ .

*Proof.* Let  $K_n$  be a connected graph with order  $n \geq 3$ . The vertex set  $V(K_n) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(K_n) = \{v_i v_{i+k}; 1 \leq i \leq n, 1 \leq k \leq n-i\}$ . The cardinality of vertex set and edge set, respectively are  $|V(K_n)| = n$  and  $|E(K_n)| = \frac{n(n-1)}{2}$ . Every vertices of  $K_n$  is adjacent to all other vertices. Suppose  $u, v \in V(K_n)$  such that  $N(u) - \{v\} = N(v) - \{u\}$ , based on Observation 1.1 that  $\lambda(u) \neq \lambda(v)$  then the labels of all vertices in  $K_n$  are distinct, consequently that  $dis_l(K_n) \geq n$ . We define the label vertices using formula  $\lambda(v_i) = i$  for  $1 \leq i \leq n$ . This labeling provides vertex-weight as follows

$$w(v_i) = \frac{n(n+1)}{2} - i; 1 \leq i \leq n$$

It is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $K_n$  is  $dis_l(K_n) \leq n$ . It concludes that  $dis_l(K_n) = n$ .  $\square$

**Theorem 2.4.** *Let  $S_n$  be a star graph with order  $n \geq 4$ , then we have the local distance irregular labeling of  $S_n$  is  $dis_l(S_n) = 1$ .*

*Proof.* Let  $S_n$  be a connected graph with order  $n \geq 4$ . The vertex set  $V(S_n) = \{v, v_1, v_2, \dots, v_n\}$  and edge set  $E(S_n) = \{v v_i; 1 \leq i \leq n\}$ . The cardinality of vertex set and edge set, respectively are  $|V(S_n)| = n + 1$  and  $|E(S_n)| = n$ . We have  $\Delta(S_n) = n$ ,  $\delta(S_n) = 1$  and  $\chi(P_n) = 2$ . Based on Lemma 2.1 that the lower bound of local distance irregular labeling of star graph  $S_n$  is  $dis_l(S_n) \geq \lceil \frac{\chi + \delta - 1}{\Delta} \rceil = \lceil \frac{2 + 1 - 1}{n} \rceil = 1$ . However, we attain the sharpest lower bound. Furthermore, we prove the upper bound of the local distance irregular labeling of star graph is  $dis_l(S_n) \leq 1$ . We define the label vertices using formula  $\lambda(v) = \lambda(v_i) = 1$  for  $1 \leq i \leq n$ . This labeling provides vertex-weight such as  $w(v) = n$  and  $w(v_i) = 1$ . Each leaf adjacent to central vertex and  $w(v) \neq w(v_i)$ . Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $S_n$  is  $dis_l(S_n) \leq 1$ . It concludes that  $dis_l(S_n) = 1$ .  $\square$

**Theorem 2.5.** *Let  $T_{n,m}$  be a tadpole graph with order  $n, m \geq 3$ , then we have the local distance irregular labeling of  $T_{n,m}$  is  $dis_l(T_{n,m}) = 2$ .*

*Proof.* The  $(m, n)$ -tadpole graph, also called a dragon graph, is the graph obtained by joining a cycle graph  $C_n$  to a path graph  $P_m$  with a bridge. The vertex set  $V(T_{n,m}) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_m\}$  and edge set  $E(T_{n,m}) = \{v_1 v_n, v_i v_{i+1}; 1 \leq i \leq n-1\} \cup \{v_1 u_1, u_j u_{j+1}; 1 \leq j \leq m-1\}$ . The cardinality of vertex set and edge set, respectively are  $|V(T_{n,m})| = n + m$  and  $|E(T_{n,m})| = n + m$ . We have  $\Delta(T_{n,m}) = n$ ,  $\delta(T_{n,m}) = 1$  and  $\chi(T_{n,m}) = 3$ . Based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $(m, n)$ -tadpole graph  $T_{n,m}$  is  $dis_l(T_{n,m}) \geq \lceil \frac{\chi + \delta - 1}{\Delta} \rceil = \lceil \frac{3 + 1 - 1}{n} \rceil = 2$ . However, we attain the sharpest lower bound. Furthermore, we prove the upper bound of the local distance irregular labeling of  $(n, m)$ -tadpole graph  $T_{n,m}$  is  $dis_l(T_{n,m}) \leq 2$ . We define the label vertices for  $n \equiv 0(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 1, 2, 3(\text{mod}4), j \equiv 0, 1, 2(\text{mod}4) \\ 2, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 0(\text{mod}4), j \equiv 3(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_m \text{ for } m \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i, v = u_j \text{ and } v = u_m, i \equiv 0(\text{mod}2), j \equiv 1(\text{mod}2) \text{ and } m \equiv 0(\text{mod}4) \\ 3, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 1(\text{mod}2), j \equiv 0(\text{mod}2) \\ 4, & \text{if } v = v_1 \end{cases}$$

The label vertices for  $n \equiv 3(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i \text{ and } v = u_j, i, j \equiv 0, 1, 2(\text{mod}4) \\ 2, & \text{if } v = v_i \text{ and } v = u_j, i, j \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_m \text{ for } m \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i, v = u_j \text{ and } v = u_m, i, j \equiv 1(\text{mod}2), i \neq 1 \text{ and } m \equiv 0(\text{mod}4) \\ 3, & \text{if } v = v_i \text{ and } v = u_j, i, j \equiv 0(\text{mod}2) \\ 4, & \text{if } v = v_1 \end{cases}$$



The label vertices for  $n \equiv 1, 2(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 1, 2, 3(\text{mod}4), j \equiv 0, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 0(\text{mod}4), j \equiv 1(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_m \text{ for } m \equiv 1, 3, 0(\text{mod}4) \\ 2, & \text{if } v = v_i, v = u_j \text{ and } v = u_m, i \equiv 0(\text{mod}2), j \equiv 1(\text{mod}2) \text{ and } m \equiv 2(\text{mod}4) \\ 3, & \text{if } v = v_i \text{ and } v = u_j, i \equiv 1(\text{mod}2), j \equiv 0(\text{mod}2) \\ 5, & \text{if } v = v_1 \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $(n, m)$ -tadpole graph  $T_{n,m}$  is  $dis_l(T_{n,m}) \leq 2$ . It concludes that  $dis_l(T_{n,m}) = 2$ .  $\square$

**Theorem 2.6.** *Let  $C_n^2$  be a unicycle with two pendants of order  $n \geq 3$ , then we have the local distance irregular labeling of  $C_n^2$  is  $dis_l(C_n^2) = 2$ .*

*Proof.* Let  $C_n^2$  be a unicycle with two pendants of order  $n \geq 3$ , the vertex set  $V(C_n^2) = \{v_1, v_2, \dots, v_n, u_1, u_2\}$  and edge set  $E(C_n^2) = \{v_1v_n, v_iv_{i+1}, v_1u_1, v_2u_2; 1 \leq i \leq n-1\}$ . The cardinality of vertex set and edge set, respectively are  $|V(C_n^2)| = n+2$  and  $|E(C_n^2)| = n+2$ . We have  $\Delta(C_n^2) = 3$ ,  $\delta(C_n^2) = 1$  and  $\chi(C_n^2) = 3$ . Based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $C_n^2$  is  $dis_l(C_n^2) \geq \lceil \frac{\chi+\delta-1}{\Delta} \rceil = \lceil \frac{3+1-1}{3} \rceil = 1$ . However, we can not attain the sharpest lower bound. If all vertices assigned label 1, then there is at least two adjacent vertices which have the same weight namely  $w(v_3) = \lambda(v_2) + \lambda(v_4) = 1 + 1 = 2$  and  $w(v_4) = \lambda(v_3) + \lambda(v_5) = 1 + 1 = 2$  so that  $w(v_3) = w(v_4)$  for edge  $v_3v_4 \in E(C_n^2)$ . Hence, the lower bound of the local distance irregular labeling of  $C_n^2$  is  $dis_l(C_n^2) \geq 2$ . Furthermore, we prove the upper bound of the local distance irregular labeling of  $C_n^2$  is  $dis_l(C_n^2) \leq 2$ . We define the label vertices for  $n \equiv 0(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, i \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = u_1, v = u_2 \text{ and } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_1 \text{ and } v = u_2 \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2), i \neq 2 \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2), i \neq 1 \\ 4, & \text{if } v = v_2 \\ 5, & \text{if } v = v_1 \end{cases}$$

The label vertices for  $n \equiv 3(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i, v = v_1 \text{ and } v = u_1, i \equiv 0, 2, 3(\text{mod}4) \\ 2, & \text{if } v = v_i \text{ and } v = u_2, i \equiv 1(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_1 \text{ and } v = u_2 \\ 2, & \text{if } v = v_i, i \equiv 1(\text{mod}2), i \neq 1 \\ 3, & \text{if } v = v_i \text{ and } v = v_1, i \equiv 0(\text{mod}2), i \neq 2 \\ 4, & \text{if } v = v_2 \end{cases}$$

The label vertices for  $n \equiv 2(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = v_i \text{ and } v = u_1, i \equiv 1, 2, 3(\text{mod}4) \\ 2, & \text{if } v = u_2 \text{ and } v = v_i, i \equiv 0(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_1 \text{ and } v = u_2 \\ 2, & \text{if } v = v_i, i \equiv 0(\text{mod}2), i \neq 2 \\ 3, & \text{if } v = v_i, i \equiv 1(\text{mod}2) \\ 4, & \text{if } v = v_2 \end{cases}$$

The label vertices for  $n \equiv 1(\text{mod}4)$  using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = u_1, v = u_2 \text{ and } v = v_i, i \equiv 1, 2, 0(\text{mod}4) \\ 2, & \text{if } v = u_2 \text{ and } v = v_i, i \equiv 3(\text{mod}4) \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_1 \text{ and } v = u_2 \\ 2, & \text{if } v = v_i, i \equiv 1(\text{mod}2), i \neq 1 \\ 3, & \text{if } v = v_1 \text{ and } v = v_i, i \equiv 0(\text{mod}2) \\ 4, & \text{if } v = v_2 \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $C_n^2$  is  $dis_l(C_n^2) \leq 2$ . It concludes that  $dis_l(C_n^2) = 2$ .  $\square$

**Theorem 2.7.** *Let  $M_n$  be a sun graph with order  $n \geq 3$ , then we have the local distance irregular labeling of  $M_n$  is  $dis_l(M_n) = 2$ .*

*Proof.* Let  $M_n$  be a unicycle with  $n$  pendants of order  $n \geq 3$ , the vertex set  $V(M_n) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n\}$  and edge set  $E(M_n) = \{v_1v_n, v_iv_{i+1}; 1 \leq i \leq n-1\} \cup \{v_iu_i; 1 \leq i \leq n\}$ . The cardinality of vertex set and edge set, respectively are  $|V(M_n)| = 2n$  and  $|E(M_n)| = 2n$ . We have  $\Delta(M_n) = 3$ ,  $\delta(M_n) = 1$  and  $\chi(M_n) = 3$ . Based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $M_n$  is  $dis_l(M_n) \geq \lceil \frac{\chi+\delta-1}{\Delta} \rceil = \lceil \frac{3+1-1}{3} \rceil = 1$ . However, we can not attain the sharpest lower bound. If all vertices assigned label 1, then there is at least two adjacent vertices which have the same weight namely  $w(v_3) = \lambda(v_2) + \lambda(v_4) = 1 + 1 = 2$  and  $w(v_4) = \lambda(v_3) + \lambda(v_5) = 1 + 1 = 2$  so that  $w(v_3) = w(v_4)$  for edge  $v_3v_4 \in E(C_n^2)$ . Hence, the lower bound of the local distance irregular labeling of  $M_n$  is  $dis_l(M_n) \geq 2$ . Furthermore, we prove the upper bound of the local distance irregular labeling of  $M_n$  is  $dis_l(M_n) \leq 2$ . We define the label vertices for  $n$  is even using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = u_i; i \text{ is odd and } v = v_i; 1 \leq i \leq n \\ 2, & \text{if } v = u_i; i \text{ is even} \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_i, 1 \leq i \leq n \\ 3, & \text{if } v = v_i, i \text{ is odd} \\ 4, & \text{if } v = v_i, i \text{ is even} \end{cases}$$

The label vertices for  $n$  is odd using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = u_i; i \text{ is odd, } i \neq 1 \text{ and } v = v_i; 1 \leq i \leq n-1 \\ 2, & \text{if } v = v_n, v = u_1 \text{ and } v = u_i; i \text{ is even} \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_i, 1 \leq i \leq n-1 \\ 2, & \text{if } v = u_n \\ 3, & \text{if } v = v_i, i \text{ is odd, } i \neq 1 \\ 4, & \text{if } v = v_i, i \text{ is even} \\ 5, & \text{if } v = v_1 \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $M_n$  is  $dis_l(M_n) \leq 2$ . It concludes that  $dis_l(M_n) = 2$ .  $\square$

**Theorem 2.8.** *Let  $T_{2,h}$  be a complete binary tree graph of height  $h \geq 3$ , then we have the local distance irregular labeling of  $T_{2,h}$  is  $dis_l(T_{2,h}) = 2$ .*

*Proof.* Let  $T_{2,h}$  be a tree graph with height  $h \geq 3$ , the vertex set  $V(T_{2,h}) = \{u, u_k^i; 1 \leq i \leq h, 1 \leq k \leq 2^h\}$  and edge set  $E(T_{2,h}) = \{uu_1^1, uu_2^1\} \cup \{u_k^i u_{2k-1}^{i+1}, u_k^i u_{2k-1}^{i+1}; 1 \leq i \leq h-1, 1 \leq k \leq 2^i\}$ . The cardinality of vertex set and edge set, respectively are  $|V(T_{2,h})| = 2^{h+1} - 1$  and  $|E(T_{2,h})| = 2^{h+1} - 2$ . We have  $\Delta(T_{2,h}) = 3$ ,  $\delta(T_{2,h}) = 1$  and  $\chi(T_{2,h}) = 2$ . Based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $T_{2,h}$  is  $dis_l(T_{2,h}) \geq \lceil \frac{\chi+\delta-1}{\Delta} \rceil = \lceil \frac{2+1-1}{3} \rceil = 1$ . However, we can not attain the sharpest lower bound. If all vertices assigned label 1, then there is at least two adjacent vertices which have the same weight namely  $w(v_1^1) = w(v_2^1)$  for edge  $v_1^1 v_2^1 \in E(T_{2,h})$ . Hence, the lower bound of the local distance irregular labeling of  $T_{2,h}$  is  $dis_l(T_{2,h}) \geq 2$ . Furthermore, we prove the upper bound of the local distance irregular labeling of  $T_{2,h}$  is  $dis_l(T_{2,h}) \leq 2$ . We define the label vertices for  $h$  odd using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = u \text{ and } v = u_k^i; i \text{ is even, } 1 \leq i \leq h, 1 \leq k \leq 2^i \\ 2, & \text{if } v = u_k^i; i \text{ is odd, } 1 \leq i \leq h, 1 \leq k \leq 2^i \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 1, & \text{if } v = u_k^h, 1 \leq k \leq 2^h \\ 3, & \text{if } v = u_k^i, i \text{ is odd, } 1 \leq i \leq h-1 \\ 4, & \text{if } v = u \\ 6, & \text{if } v = u_k^i, i \text{ is even, } 1 \leq i \leq h \end{cases}$$

The label vertices for  $h$  even using formula.

$$\lambda(v) = \begin{cases} 1, & \text{if } v = u \text{ and } v = u_k^i; i \text{ is even, } 1 \leq i \leq h, 1 \leq k \leq 2^i \\ 2, & \text{if } v = u_k^i; i \text{ is odd, } 1 \leq i \leq h, 1 \leq k \leq 2^i \end{cases}$$

This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2, & \text{if } v = u_k^h, 1 \leq k \leq 2^h \\ 3, & \text{if } v = u_k^i, i \text{ is odd, } 1 \leq i \leq h \\ 4, & \text{if } v = u \\ 6, & \text{if } v = u_k^i, i \text{ is even, } 1 \leq i \leq h-1 \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $T_{2,h}$  is  $dis_l(T_{2,h}) \leq 2$ . It concludes that  $dis_l(T_{2,h}) = 2$ .  $\square$

**Observation 2.1.** Let  $K_{n,m}$  be  $k$ -partite graph with  $n, m \geq 2$ . If the vertices  $u, v \in V(K_{n,m})$  in every partite class are not adjacent, then  $N(u) = N(v)$  with  $u, v$  belong to the same partite class.

**Theorem 2.9.** Let  $K_{m,n}$  be a complete bipartite graph of  $n, m \geq 3$ , then we have the local distance irregular labeling of  $K_{m,n}$  is

$$dis_l(K_{m,n}) = \begin{cases} 1, & \text{if } n \neq m \\ 2, & \text{if } n = m \end{cases}$$

*Proof.* Let  $K_{m,n}$  be a complete bipartite graph of  $n, m \geq 3$ . The vertex set  $V(K_{m,n}) = \{v_i, u_j; 1 \leq i \leq n, 1 \leq j \leq m\}$  and edge set  $E(K_{m,n}) = \{v_i u_j; 1 \leq i \leq n, 1 \leq j \leq m\}$ . The cardinality of vertex set and edge set, respectively are  $|V(K_{m,n})| = n + m$  and  $|E(K_{m,n})| = nm$ . We have  $\Delta(K_{m,n}) = m$ ,  $\delta(K_{m,n}) = n$  and  $\chi(K_{m,n}) = 2$ . The proof divided into two cases as follows.

**Case 1:** For  $n \neq m$  ( $n < m$  or  $n > m$ ), based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \geq \lceil \frac{\chi+\delta-1}{\Delta} \rceil = \lceil \frac{2+n-1}{m} \rceil = 1$ . However, we attain the sharpest lower bound. Furthermore, we prove the upper bound of the local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \leq 1$ . We define the label vertices using formula includes  $\lambda(v_i) = \lambda(u_j) = 1$ . This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} m, & \text{if } v = v_i, 1 \leq i \leq n \\ n, & \text{if } v = u_j, 1 \leq j \leq m \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \leq 1$ . It concludes that  $dis_l(K_{m,n}) = 1$ .

**Case 2:** For  $n = m$ , based on Lemma 2.1 that the lower bound of local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \geq \lceil \frac{x+\delta-1}{\Delta} \rceil = \lceil \frac{2+m-1}{m} \rceil = 2$ . However, we attain the sharpest lower bound. Furthermore, we prove the upper bound of the local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \leq 2$ . We define the label vertices using formula includes  $\lambda(v_i) = 1$  and  $\lambda(u_j) = 2$ . This labeling provides vertex-weight as follows.

$$w(v) = \begin{cases} 2m, & \text{if } v = v_i, 1 \leq i \leq n \\ n, & \text{if } v = u_j, 1 \leq j \leq m \end{cases}$$

Therefore, it is easy to see that  $w$  is vertex coloring. Hence, we obtain the upper bound of the local distance irregular labeling of  $K_{m,n}$  is  $dis_l(K_{m,n}) \leq 2$ . It concludes that  $dis_l(K_{m,n}) = 2$ .  $\square$

### 3. CONCLUSIONS

In this section, we conclude that local distance irregular labeling is a new invariant. We find the exact value of local distance irregular labeling of some families graph, namely path graph, cycle graph, star graph, complete graph, tadpole graph, unicycle with two pendant graph, binary tree graph, complete bipartite graph, and sun graph. There is some open problems for research in this area. We state a few interesting problems as follows.

**Open Problem 3.1.** *Determine the local distance irregular labeling for graph operation namely corona product, Cartesian product, joint, comb product and others.*

**Open Problem 3.2.** *Determine the local distance irregular labeling for some family graph namely unicycle, related wheel, tree graph and cayley graph.*

### ACKNOWLEDGEMENT

The authors of this article gratefully acknowledge the support from the University of Jember in 2020.

### REFERENCES

- [1] Bača, M., Muntaner-Batle, F. A., Semaničová-Feňovčíková, A., (2014). On super  $(a, 2)$ -edge-antimagic total labeling of Disconnected Graphs, *Ars Combin.*, 113, pp. 129–137.
- [2] Chartrand, G. and Lesniak, L., (2000). *Graphs and digraphs*, 3rd ed (London: Chapman and Hall).
- [3] Dafik, Miller, Ryan, J., and Baca, M., (2011). Super edge-antimagic total labelings of  $mK_{n,n}$ , *Ars Combinatoria*, 101, pp. 97–107.
- [4] Dafik, Miller, Ryan, J., and Baca, M., (2008). On antimagic labelings of disjoint union of complete  $s$ -partite graphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, 65, pp. 41-49.
- [5] Dafik, Purnapraja, A.K. and Hidayat, R., (2015). Cycle-Super Antimagicness of Connected and Disconnected Tensor Product of Graphs, *Procedia Computer Science*, 74, pp. 93–99.
- [6] Gross, J. L., Yellen, J. and Zhang, P., (2014). *Handbook of graph Theory*, Second Edition CRC Press Taylor and Francis Group.
- [7] Hartsfield, N. and Ringel, G., (1994). *Pearls in Graph Theory*, Academic Press, United Kingdom.
- [8] Slamini, (2017). On distance irregular labelings of graphs, *Far East Journal Mathematical Sciences*, 102(5), pp. 919-932.
- [9] Baca, M., Dafik, Miller, M., and Ryan, J., (2009). Antimagic labeling of disjoint union of  $s$ -crowns, *Utilitas Mathematica*, 79, pp. 193–205.
- [10] Gallian J. A., (2013). *Graph Labeling*, *Electronic Journal of Combinatorics*, 5, (Dynamic Survey DS6)
- [11] Sedlacek, J., (1963). Problem 27. *Theory of graphs and its applications*, Smolenice, pp. 163-164.



**Dr. Arika Indah Kristiana** completed her Ph.D. in mathematics (2019) from the Department of Mathematics and Sciences, Airlangga University, Indonesia. She is currently working in graph theory such as graph labeling and coloring.



**Ridho Alfarisi** completed his M.Sc. in mathematics (2017) from the Department of Mathematics, Institut Teknologi Sepuluh Nopember, Indonesia. Currently he is a Ph.D. student at Airlangga University, Indonesia. He is working in graph theory such as metric dimension and partition dimension.



**Prof. Dr. Dafik** completed his Ph.D. in mathematics (2007) at Ballarat University, Australia. In 2013, he became a professor of graph theory and combinatorics. He is currently working in graph theory such as graph labeling, domination number, and coloring.

---

