



Hindawi

Advances in Pharmacological and Pharmaceutical Sciences

Volume 2022

ISSN: 2633-4682 (Print)

ISSN: 2633-4690 (Online)

Chief Editor

Kim Wei Chan , Institute of Bioscience, Universiti Putra Malaysia, Malaysia

Editorial Board

Ergin Murat ALTUNER , Kastamonu University, Turkey

Rizky Abdulah , Universitas Padjadjaran, Indonesia

Antreas Afantitis, NovaMechanics Ltd, Cyprus

Esther Agbaje, University of Lagos, Nigeria

Mahmoud S. Ahmed, University of Texas Medical Branch, USA

Nirav H. Amin, Loma Linda University, Loma Linda, CA, USA

Athar Ata, The University of Winnipeg, Canada

Abdelhakim Bouyahya , Laboratory of Human Pathologies Biology, Department of Biology, Faculty of Sciences, and Genomic Center of Human Pathologies, Mohammed V University, Rabat,, Morocco

Maria T. D. S. Correia , Departamento de Bioquímica, Centro de Biociências, Universidade Federal de Pernambuco. Av. Prof. Moraes Rego, 1235 - Cidade Universitária - CEP 50.670-901 - Recife - PE - Brazil, Brazil

Priyankar Dey , University of North Bengal, India

Ahmed Elkamhawy, Dongguk University, Republic of Korea

Norhaizan Mohd Esa , Universiti Putra Malaysia, Malaysia

Jia You Fang, Chang Gung University, Taiwan

Antonio Ferrer-Montiel , Universidad Miguel Hernandez, Spain

Ghulam Hussain , Department of Physiology, Faculty of Life Sciences, Government College University, Faisalabad, 38000 Pakistan, Pakistan

Gunawan Indrayanto , Surabaya University, Indonesia

Abdeslam Jaafari, Faculty of Science and Technology, Sultan Moulay Slimane University. Beni Mellal, Morocco

Heng Yen Khong , Universiti Teknologi MARA, Malaysia

Ismail Laher , University of British Columbia, Canada

Chun-Guang Li , Western Sydney University, Australia

Koteshwara Mudigonda, Suven Life Sciences Limited, India

Srinivas Mutalik, Manipal University, India

Benedetto Natalini , Università degli Studi di Perugia, Italy

Berend Olivier , Utrecht University, The Netherlands

Bernard Pirotte, University of Liège, Belgium

Suriana Sabri, Department of Microbiology, Faculty of Biotechnology and Molecular Biosciences,
Universiti Putra Malaysia, Malaysia

Chita Ranjan Sahoo , Siksha O Anusandhan Deemed to be University, India

Kuldeep Singh , Maharishi Markandeshwar (Deemed to be University), Mullana, Haryana - 133207,
India

Jirapornchai Suksaeree , Rangsit University, Thailand

Mounir Tilaoui , Laboratory of Biological Engineering. Natural Substances, Cellular and Molecular
Immuno-pharmacology, Immunobiology of Cancer Cells Cluster. Faculty of Sciences and
Technology. Béni Mellal, Morocco

Nizar Tlili, Laboratoire de Biochimie Campus Universitaire , Tunisia

Jaleh Varshosaz , Isfahan University of Medical Sciences, Iran

Hansen Wang , University of Toronto, Canada

Qi-Dong You , China Pharmaceutical University, China

Tahira Younis, GCUF, Pakistan

Tamer E. Youssef , Department of Chemistry, College of Science, Imam Abdulrahman Bin Faisal
University, Saudi Arabia

Apostolos Zarros, University of Glasgow, United Kingdom

Contents

- Molecular Docking and Dynamics Simulation of Several Flavonoids Predict Cyanidin as an Effective Drug Candidate against SARS-CoV-2 Spike Protein. Asmita Shrestha ,Rishab Marahatha, Saroj Basnet, Bishnu P. Regmi, Saurav Katuwal, Salik Ram Dahal, Khaga Raj Sharma, Achyut Adhikari, Ram Chandra Basnyat, and Niranjan Parajuli. Volume 2022. <https://doi.org/10.1155/2022/3742318> - 09 Nov 2022
- Free Radical Scavenging, Anti-Infectious, and Toxicity Activities from *Stenochlaena palustris* (Burm.f.) Bedd. Extracts. Rudi Hendra, Rohimatul Khodijah, Muhammad Almurdati, Yuli Haryani, Ari Satia Nugraha, Neni Frimayanti, Hilwan Yuda Teruna, and Rizky Abdulah. Volume 2022 <https://doi.org/10.1155/2022/5729217> - 01 Nov 2022
- Antiproliferative Effects of Ferulic, Coumaric, and Caffeic Acids in HepG2 Cells by hTERT Downregulation. Afsoon Afshari, Mahmoodreza Moein, Afshin Afsari, and Zahra Sabahi. Volume 2022 <https://doi.org/10.1155/2022/1850732> - 28 Oct 2022
- Current Trends on Solid Dispersions: Past, Present, and Future. Ruba Malkawi, Walla I. Malkawi, Yahia Al-Mahmoud, and Jawad Tawalbeh. Volume 2022 <https://doi.org/10.1155/2022/5916013> - 22 Oct 2022
- In Vitro Mechanism of Action of *Acanthospermum hispidum* in *Trypanosoma brucei*. Aboagye Kwarteng Dofuor, Temitayo Samson Ademolue, Karen Nana Akua Kuampah, Frederick Ayertey, and Theresa Manful Gwira. Volume 2022 <https://doi.org/10.1155/2022/1645653> - 18 Oct 2022
- Phytochemical Screening, Antioxidant, and Antimicrobial Activities of Seven Underinvestigated Medicinal Plants against Microbial Pathogens. Borel Ndezo Bisso, Roland Njikang Epie Nkwelle, Roland Tchuenguem Tchuenteu, and Jean Paul Dzoyem. Volume 2022 <https://doi.org/10.1155/2022/1998808> - 10 Oct 2022
- Biological Activity and Chemical Composition of *Detarium microcarpum* Guill. and Perr—A Systematic Review. Abdulrahman Mahmoud Dogara. Volume 2022 <https://doi.org/10.1155/2022/7219401> - 08 Oct 2022
- In Silico, In Vitro, and In Vivo Wound Healing Activity of *Astragalus microcephalus* Willd. Fatemeh Akbari, Mohammad Azadbakht, Ali Bagheri, and Lale Vahedi. Volume 2022 <https://doi.org/10.1155/2022/2156629> - 04 Oct 2022
- Neuroprotective Effects of the Ethanolic Leaf Extract of *Crassocephalum crepidioides* (Asteraceae) on Diazepam-Induced Amnesia in Mice. Beppe Galba Jean, Folefack Alice Irène, Atsamo Albert Donatien, Ngatanko Aïbassou Hervé-Hervé, Barga Mpoo Bertrand, Allah-Doum Nanou Gael, Nguedia Ymele Merline, Zemo Gamo Franklik, Dongmo Alain Bertrand, and Dimo Théophile. Volume 2022 <https://doi.org/10.1155/2022/1919469> - 28 Sep 2022
- In Vitro and In Vivo Antiproliferative Actions of *Solanum gilo* Raddi (Solanaceae) Fruit Extract on Breast Tissues. Edwige Nana Tchoupang, Vincent Yadji, Jean Baptiste Wangbara, Marius Trésor Kemegne Sipping, Stéphane Zingue, and Dieudonné Njamen. Volume 2022 <https://doi.org/10.1155/2022/6834626> - 26 Sep 2022
- Antibiotic Use among Patients Visiting Primary Hospitals in Northwest Ethiopia: A Multicenter Cross-Sectional Survey. Adaladlew Kassie Netere and Ashenafi Kibret Sendekie. Volume 2022 <https://doi.org/10.1155/2022/2306637> - 10 Sep 2022
- Anti-Arthritic Effect of the Hydroethanolic Root Extract of *Psydrax subcordata* in Rats. Phyllis Annan, Newman Osafo, Paul Poku Sampene Ossei, and Wonder Kofi Mensah Abotsi. Volume 2022 <https://doi.org/10.1155/2022/9748382> - 24 Aug 2022
- N-Acetylcysteine and Other Sulfur-Donors as a Preventative and Adjunct Therapy for COVID-19. Heidi N du Preez, Colleen Aldous, Hendrik G Kruger, and Lin Johnson. Volume 2022 <https://doi.org/10.1155/2022/4555490> - 10 Aug 2022
- Antiplasmodial, Antitrypanosomal, and Cytotoxic Effects of *Anthonotha macrophylla*, *Annickia polycarpa*, *Tieghemella heckelii*, and *Antrocaryon micraster* Extracts. Aboagye

Kwarteng Dofuor, Emmanuel Kofi Kumatia, Jersley Didewurah Chirawurah, and Frederick Ayertey. Volume 2022 <https://doi.org/10.1155/2022/9195753> - 23 Jul 2022

- Evaluation and Clinical Validation of Guanidine-Based Inactivation Transport Medium for Preservation of SARS-CoV-2. Hesti L. Wiraswati, Shabarni Gaffar, Savira Ekawardhani, Nisa Fauziah, Fedri R. Rinawan, Leonardus Widyatmoko, Amila Laelalugina, Annissa R. Arimdayu, Tri Kusniati, Clarisa D. Andari, and Lia Faridah. Volume 2022 <https://doi.org/10.1155/2022/1677621> - 21 Jul 2022
- Phytochemical Analysis and Antioxidant and Antidiabetic Activities of Extracts from *Bergenia ciliata*, *Mimosa pudica*, and *Phyllanthus emblica*. Basanta Kumar Sapkota, Karan Khadayat, Kabita Sharma, Bimal Kumar Raut, Dipa Aryal, Bijaya Bahadur Thapa, and Niranjana Parajuli. Volume 2022 <https://doi.org/10.1155/2022/4929824> - 07 Jul 2022
- Phytochemical Screening and In Vitro Antifungal Activity of Selected Medicinal Plants against *Candida albicans* and *Aspergillus niger* in West Shewa Zone, Ethiopia. Askale Gizaw, Lencho Megersa Marami, Ibsa Teshome, Edilu Jorga Sarba, Petros Admasu, Dagmawit Atalel Babele, Getachew Mulatu Dilba, Wakuma Mitiku Bune, Morka Dandecha Bayu, Miressa Tadesse, and Kebede Abdisa. Volume 2022 <https://doi.org/10.1155/2022/3299146> - 28 Jun 2022
- Antioxidant, Antidiabetic, and Antibacterial Potentials and Chemical Composition of *Salvia officinalis* and *Mentha suaveolens* Grown Wild in Morocco. Samiah Hamad Al-Mijalli, Hamza Assaggaf, Ahmed Qasem, Adel G. El-Shemi, Emad M. Abdallah, Hanae Naceiri Mrabti, and Abdelhakim Bouyahya. Volume 2022 <https://doi.org/10.1155/2022/2844880> - 15 Jun 2022
- Antimicrobial Potential of *Streptomyces* spp. Isolated from the Rift Valley Regions of Ethiopia. Firew Elias, Sudhamani Muddada, Diriba Muleta, and Belachew Tefera. Volume 2022 <https://doi.org/10.1155/2022/1724906> - 13 Jun 2022
- Levofloxacin Use in Patients with Suspected Tuberculosis in a Community Hospital, Thailand: A Pilot Study. Thanawat Khongyot, Sawitree Laopaiboonkun, Throngpon Kawpradid, Kannuwat Jitkamrop, Tawee Chanphakphoom, and Suriyon Uitrakul. Volume 2022 <https://doi.org/10.1155/2022/5647071> - 03 Jun 2022
- *Passiflora mollissima* Seed Extract Induced Antiproliferative and Cytotoxic Effects on CAL 27 Spheroids. Angela Fonseca-Benitez, Sandra Johanna Morantes Medina, Diego Ballesteros-Vivas, Fabian Parada-Alfonso, and Sandra J Perdomo. Volume 2022 <https://doi.org/10.1155/2022/4602413> - 31 May 2022
- Ethnopharmacology, Biological Evaluation, and Chemical Composition of *Ziziphus spinachristi* (L.) Desf.: A Review. Mahmoud Dogara Abdulrahman, Ali Muhammad Zakariya, Harmand A. Hama, Saber W. Hamad, Sawsan S. Al-Rawi, Sarwan W. Bradosty, and Ahmad H. Ibrahim. Volume 2022 <https://doi.org/10.1155/2022/4495688> 29 May 2022
- Antidiabetic Properties and Toxicological Assessment of *Antidesma celebicum* Miq: Ethanolic Leaves Extract in Sprague–Dawley Rats. Raysa Y. Pratiwi, Berna Elya, Heri Setiawan, Roshamur C. Forestrania, and Rizna T. Dewi. Volume 2022 <https://doi.org/10.1155/2022/2584698> - 24 May 2022
- Evaluation of Dermal Toxicity and Wound Healing Activity of *Cnestis ferruginea* Vahl ex DC. Akosua Dufie Ankomah, Yaw Duah Boakye, Theresah Appiah Agana, Vivian Etsiapa Boamah, Paul Poku Sampene Ossei, Francis Adu, and Christian Agyare. Volume 2022 <https://doi.org/10.1155/2022/5268613> - 23 May 2022
- Phylogeny, Phytomedicines, Phytochemistry, Pharmacological Properties, and Toxicity of *Croton gratissimus* Burch (Euphorbiaceae). Kirkland Dingili Magwilu, Joseph Mwanzia Nguta, Isaac Mapenay, and Dorine Matara. Volume 2022 <https://doi.org/10.1155/2022/1238270> - 17 May 2022
- In Vitro Antioxidant, Anti-Inflammatory, and Digestive Enzymes Inhibition Activities of Hydro-Ethanolic Leaf and Bark Extracts of *Psychotria densinervia* (K. Krause) Verdc. Jean Romuald Mba, Djamila Zouheira, Hadidjatou Dairou, Fanta S. A. Yadang, Nfor Njini Gael,

Lawrence Ayong, Jules-Roger Kuate, and Gabriel A. Agbor. Volume 2022
<https://doi.org/10.1155/2022/8459943> - 06 May 2022







- Biological Evaluation of *Garcinia kola* Heckel. Abdulrahman Mahmoud Dogara, Saber W. Hamad, Harmand A. Hama, Sarwan W. Bradosty, Soran Kayfi, Sawsan S. Al-Rawi, and Abubakar Abdullahi Lema. Volume 2022 <https://doi.org/10.1155/2022/3837965> - 28 Apr 2022
- Development and In Vitro Evaluation of Oral Capsules from *Antiaris*: A Convenient Substitute for Peripheral Neuropathy. Mary-Ann Archer, Doris Kumadoh, Samuel Nii-Bortier Gaizer, 2Adelaide Mensah, 2,5Jonathan Jato, 2,6Micheal Odoi Kyene, Susana Oteng Mintah, 7Genevieve Naana Yeboah, 3Paul kwesi Sodji, 8and Ofosua Adi-Dako. Volume 2022 <https://doi.org/10.1155/2022/5340953> - 27 Apr 2022
- Advancement of Nanofibrous Mats and Common Useful Drug Delivery Applications. Hamza Abu Owida, Jamal I. Al-Nabulsi, Feras Alnaimat, Ashraf Al Sharah, 2Muhammad Al-Ayyad, 1Nidal M. Turab, and Mustafa Abdullah. Volume 2022 <https://doi.org/10.1155/2022/9073837> - 19 Apr 2022
- Phytochemical Analysis and Antifungal Potentiating Activity of Extracts from Loquat (*Eriobotrya japonica*) against *Cryptococcus neoformans* Clinical Isolates. Borel Ndezo Bisso, Prudence Ngulula Kayoka-Kabongo, Roland Tchuenteu Tchuenguem, and Jean Paul Dzoyem. Volume 2022 <https://doi.org/10.1155/2022/6626834> - 14 Apr 2022
- Prevalence and Associated Factors of Anemia among Breast Cancer Patients Undergoing Chemotherapy: A Prospective Study. Fares M.S Muthanna, Mahmathi Karuppanan, Egbal Abdulrahman, Suriyon Uitrakul, Bassam Abdul Hassan Rasool and Ali Haider Mohammed. Volume 2022 <https://doi.org/10.1155/2022/7611733> 14 Apr 2022
- Effects of the Antiobesity Drugs Aplex and Venera on Certain Biochemical and Physiological Indices in Obese Adult Male Albino Rats. Suzan S. A. Elpasty, Eman G. E. Helal, Ashraf M. M. Algendy and Hany N. Yousef. Volume 2022 <https://doi.org/10.1155/2022/3776676> 14 Apr 2022
- Cytotoxicity against Human Hepatocellular Carcinoma (HepG2) Cells and Anti-Oxidant Activity of Selected Endemic or Medicinal Plants in Sri Lanka. Jeyaraj Thusyanthan, Nimesha Sulochani Wickramaratne, Kanishka Sithira Senathilake, Umapriyatharshini Rajagopalan, Kamani Hemamala Tennekoon, Ira Thabrew, and Sameera Ranganath Samarakoon. Volume 2022 <https://doi.org/10.1155/2022/6407688> 30 Mar 2022
- In Vitro and In Vivo Wound Healing Activity of *Astragalus floccosus* Boiss. (Fabaceae). Fatemeh Akbari, Mohammad Azadbakht, Ali Bagheri, and Lale Vahedi. Volume 2022 <https://doi.org/10.1155/2022/7865015> 29 Mar 2022
- Formulation Design and Cell Cytotoxicity of Curcumin-Loaded Liposomal Solid Gels for Anti-Hepatitis C Virus. Helmy Yusuf, Eryn K. D. D. Novitasari, Ni L. W. Purnami, Adhe W. Mahbub, Retno Sari, and Dwi Setyawan. Volume 2022 <https://doi.org/10.1155/2022/3336837> 07 Mar 2022
- Approaches for the Elimination of Microbial Contaminants from *Lippia multiflora* Mold. Leaves Intended for Tea Bagging and Evaluation of Formulation. Doris Kumadoh, Mary-Ann Archer, Michael O. Kyene, Genevieve N. Yeboah, 1Ofosua Adi-Dako, Christina Osei-Asare, Emmanuel Adase, Susana Oteng Mintah, 5Hilda Amekyeh, and Alfred A. Appiah. Volume 2022 <https://doi.org/10.1155/2022/7235489> 27 Feb 2022
- Evaluation of Hepatoprotective Potential of Polyherbal Preparations in CCl₄-Induced Hepatotoxicity in Mice. Rayhana Begum, Sonia Akther Papia, Mst Marium Begum, Hongbin Wang, Rubaba Karim, Rebeka Sultana, Priyanka Rani Das, Taslima Begum, Md. Ragibul Islam, Nargis Manwar, and Md. Sohanur Rahman. Volume 2022 <https://doi.org/10.1155/2022/3169500> - 27 Feb 2022
- *Stephania japonica* Ameliorates Scopolamine-Induced Memory Impairment in Mice through Inhibition of Acetylcholinesterase and Oxidative Stress. Md. Yusuf Al-Amin, Amitav Lahiry,

Rafia Ferdous, Md. Kamrul Hasan, Md. Abdul Kader, AHM Khurshid Alam, Zahangir Alam Saud, and Md. Golam Sadik. Volume 2022 <https://doi.org/10.1155/2022/8305271> - 21 Feb 2022

- In Vitro and In Vivo Effect of Xylopic Acid on Cytochrome P450 Enzymes. Mary A. Agbenyeku, Regina Appiah-Opong, Ernest Obese, Robert P. Biney, Emmanuel A. Adakudugu, Arnold D. Forkuo, Silas A. Osei, Mustapha K. Abeka, and Elvis O. Ameyaw. Volume 2022 <https://doi.org/10.1155/2022/4524877> - 21 Jan 2022
- Ethnomedicinal Studies, Chemical Composition, and Antibacterial Activity of the *Mammea americana* L. Bark in the Municipality of Cértegui, Chocó, Colombia. Lina Mosquera-Chaverra, Manuel Salas-Moreno, and José Marrugo-Negrete. Volume 2022 <https://doi.org/10.1155/2022/9950625> - 19 Jan 2022
- Conventional Medicinal Uses, Phytoconstituents, and Biological Activities of *Euphorbia officinarum* L.: A Systematic Review. Imane Chamkhi, Mohamed Hnini, and Jamal Aurag. Volume 2022 <https://doi.org/10.1155/2022/9971085> - 06 Jan 2022
- Knowledge, Attitude, and Practices of Moroccan Retail Pharmacists towards Veterinary Medicines. Zeinab Snoussi and Samir Ahid. Volume 2022 <https://doi.org/10.1155/2022/9973945> - 04 Jan 2022

Research Article

Free Radical Scavenging, Anti-Infectious, and Toxicity Activities from *Stenochlaena palustris* (Burm.f.) Bedd. Extracts

Rudi Hendra ^{1,2}, Rohimatul Khodijah,¹ Muhammad Almurdati,^{1,3} Yuli Haryani ¹,
Ari Satia Nugraha ⁴, Neni Frimayanti ³, Hilwan Yuda Teruna ¹,
and Rizky Abdulah ^{2,5}

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Riau, Pekanbaru, Indonesia

²Center of Excellence in Pharmaceutical Care Innovation, Universitas Padjadjaran, Bandung, Indonesia

³Sekolah Tinggi Ilmu Farmasi Riau, Pekanbaru, Indonesia

⁴Drug Utilisation and Discovery Research Group, Faculty of Pharmacy, University of Jember, Jember, Indonesia

⁵Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia

Correspondence should be addressed to Rudi Hendra; rudi.hendra@lecturer.unri.ac.id

Received 6 July 2022; Revised 10 October 2022; Accepted 22 October 2022; Published 1 November 2022

Academic Editor: Abdeslam Jaafari

Copyright © 2022 Rudi Hendra et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Stenochlaena palustris (Burm.f.) Bedd. (*Blechnaceae*) or Kelakai is a traditional medicinal plant found in the Indonesian islands of Borneo and Sumatra. It has been used to treat wound healing, infection, and diabetes. This study looked into the free radical scavenging activity, antiplasmodial activity, toxicity, and antibacterial activity against pathogenic bacteria. The species' aerial part was extracted with methanol, followed by a liquid-liquid extraction against (*n*-hexane, dichloromethane, and ethyl acetate). The extracts' free radical scavenging activities were determined using DPPH and NO radicals. The antiplasmodial and toxicity assays were conducted using two *Plasmodium falciparum* strains (3D7 and W2) and the brine shrimp lethality test. In addition, antibacterial activity was determined using the well diffusion method. The results revealed that ethyl acetate depicted potential activities toward the assay. The ethyl acetate showed potential free radical scavenging activities with an IC₅₀ value of 51.63 ± 0.46 µg/mL (DPPH) and 60.03 ± 0.65 µg/mL (NO). The antiplasmodial activities showed that the ethyl acetate had potential activities among the extracts with an IC₅₀ value of 11.06 ± 0.45 µg/mL. However, all the extracts demonstrated nontoxic toward *Artemia salina* with LC₅₀ > 1000 µg/mL. Furthermore, the ethyl acetate demonstrated intermediate susceptibility against *B. cereus* ATCC 10876, *V. parahaemolyticus* ATCC 17802, *L. monocytogenes* ATCC 7644, and *S. Typhimurium* ATCC 14028 at a concentration of 500 µg/disc. According to these findings, the ethyl acetate extract of *S. palustris* (Burm.f.) Bedd is a promising source of natural antioxidants and antiplasmodial agents.

1. Introduction

Recent years have seen an increase in people's interest in traditional medicine as a result of its multiple modes of action, which have relatively few negative effects on humans. A plant with medicinal properties is a rich source of secondary metabolites, which are used in a variety of treatments. Because of this and considering the high level of biodiversity that exists in Indonesia, it is imperative that potential plant species, including ferns, be studied [1]. Several groups have conducted pharmacological research on ferns and related plants in response to reports of their

therapeutic efficacy, scientific curiosity, and the need for new medications. Pharmacological and ethnopharmacological studies of fern substances have revealed various pharmacological effects, including cytotoxicity, hepatoprotective activity, anti-inflammatory activity, and antimicrobial properties. Due to the need for new medications with such activities, pteridophytes and their secondary metabolites may have considerable medical value [2].

A fern belonging to the family *Blechnaceae*, *Stenochlaena palustris* (Burm.) Bedd. can be found scrambling high up trees or trailing along the ground. It is native to many tropical regions, including southern and northern India,

Malaysia, Polynesia, and Australia [3]. This particular species belongs to the family *Blechnaceae*, which contains seven species, including one endemic in the Province of Riau, *S. riauensis* [4]. In Malaysia, Thailand, the Philippines, and Indonesia, the reddish young fronds are harvested from the wild and consumed as vegetables [5].

This species has been used in traditional medicine in some countries to treat a variety of ailments. People in the Western Ghats of India, for example, use this species' fronds to treat fever, sore throat, and gastric ulcers. While the leaves and rhizomes are used as a cooling agent in the treatment of burns and ulcers [6]. In Malaysia, the plant is consumed raw or cooked with boiling water for the treatment of diarrhea, while in Sumatra, Indonesia, it is used as a mild laxative [7]. In addition, the paste is applied to ulcers, wounds, and bacterially infected skin in Sumatra Island, Indonesia, Borneo Island, Malaysia, and Indonesia. People in the central region of Papua New Guinea use the tender leaves of *S. palustris* (Burm.f.) Bedd. as a contraceptive and the plant's juice can aid in the treatment of fever [5, 8].

Although this species is edible and has medicinal properties, several pharmacology and phytochemistry studies have been published. According to Chai et al., crude water extracts of *S. palustris* (Burm.f.) Bedd. had higher antioxidant activity and polyphenol, flavonoid, and hydroxycinnamic acid contents [7, 9]. Chai and colleagues also revealed that water extract from the species acted as a potent α -glucosidase inhibitor compared to quercetin as a control [10]. Furthermore, *S. palustris* (Burm.f.) Bedd. leaf extract inhibited the food-borne pathogen *Aspergillus Niger* [11]. In addition, the methanolic extract of this species showed cholinesterase inhibitory activity against acetylcholinesterase and butyrylcholinesterase with IC_{50} of 121 and 19.8 $\mu\text{g/mL}$, respectively [12]. There is a paucity of information available regarding the phytochemistry of *S. palustris* (Burm.f.) Bedd. stenopaluside, cerebroside, lutein, and β -sitosterol-3-O- β -D-glucopyranoside were isolated from the leave of *S. palustris* (Burm.f.) Bedd [13]. In addition, stenopalustrosides A-E and kaempferol glucoside were isolated from species' leaves, and stenopalustrosides A-D showed significant antibacterial activities against Gram-positive strains [3]. Furthermore, kaempferol 3-O-(3''-O-E-p-coumaroyl)-(6''-O-E-feruloyl)- β -D-glucopyranoside and kaempferol 3-O-(3'', 6'' di-O-E-p-coumaroyl)- β -D-glucopyranoside isolated from the species showed cytotoxic properties against breast (MCF-7 and MDA-MB-231) and prostate (DU-145) cancer cell lines [14].

In this study, we reported the details of the extraction of *S. palustris* (Burm.f.) Bedd. producing less polar subextracts, as well as the free radical scavenging evaluation. In a previous study according to the findings of Arullappan et al., ethyl acetate extracted from the leaves of *S. palustris* (Burm.f.) Bedd. exhibited DPPH radical inhibitor activity with an IC_{50} value of 650 $\mu\text{g/mL}$ [5]. Even though the DPPH method is used generally in antioxidant activity, the results are not standardized in which direct comparison of antioxidant strength in both different plant extracts and pure compounds are difficult. Therefore, in this study, we report the antioxidant activity index based on the method by

Scherer and Godoy (2009) [15]. This analysis allows us to compare the antioxidant capacity of the extracts to that of the positive control. Moreover, this is the first time that radical scavenging activity using the nitric oxide radical that correlates with inflammation has been reported from the extracts of this species. A novel antiplasmodium and antibacterial activity from the species' extract for the first time. In addition, molecular chemotypes previously generated from this species were also discussed in correlation with the planta pharmacological claims.

2. Materials and Methods

2.1. Plant Material. Fresh fronds of *S. palustris* (Burm.f.) Bedd. were gathered at Science Park Universitas Riau (0.46587747854712624 and 101.38066586137639). Before the grinding process began, the species was air-dried and confirmed to be completely dry, as its weight remained constant. The soil samples were then placed in an airtight container and stored between 2 and 7°C until further analysis. A botanist (Dr. Nery Sofianty) has deposited a voucher specimen (No. 37/UN19.5.1.1.3/2019) with the Department of Biology at Universitas Riau.

2.2. Extraction. Maceration in methanol of up to 2 kg of ground samples for 1 × 24 hours was repeated three times until the maceration results were no longer dark green. The extract was filtered during the maceration process, and the filtrate was collected. The macerate was then concentrated using a 40°C rotary evaporator, yielding a crude extract. The crude methanol extract was then partitioned using a liquid-liquid extraction method. To obtain the *n*-hexane extract, the methanol extract was partitioned with water in a 9:1 methanol: water ratio before being partitioned with *n*-hexane in a 1:4 ratio. The methanol-water extract was then watered down to 40% (v/v) and partitioned with dichloromethane solvent in a 1:4 ratios to produce dichloromethane extract. The methanol-water extract was then evaporated to form a water extract, which was then partitioned in a 1:3 ratio with ethyl acetate to yield the ethyl acetate extract. The remaining water extract was evaporated to produce the finished product, which was water extract [16].

2.3. Antioxidant Activities

2.3.1. DPPH Radical Scavenging Activity. An antioxidant activity assay was carried out using the DPPH method (1,1-diphenyl-2-picryl hydrazyl). [1, 17, 18] Various extracts with a final concentration of 1000 $\mu\text{g/mL}$ were diluted using the two-fold dilution method (1000–31.25 $\mu\text{g/mL}$) in a 96-well clear polystyrene microplate. Fifty μL of the sample was mixed with 80 μL of 80 $\mu\text{g/mL}$ DPPH, and the mixture was incubated in the dark for 30 minutes. Absorbance was measured at 520 nm using a Berthold microplate reader. As a positive control, ascorbic acid was subjected to the same method. The assay was carried out in triplicate, and the data were reported as mean \pm standard deviation.

The % Inhibition value is calculated by the following formula:

$$\% \text{Inhibition} = \frac{(A_0 - A_s)}{A_0} \times 100\%. \quad (1)$$

IC₅₀ analysis was performed using GraphPad Prism 9 software, where A_0 is the absorbance of the DPPH radical solution without the sample and A_s is the absorbance of the sample with DPPH radical solution. The assay was carried out in triplicate, and the data were reported as mean \pm standard deviation.

The antioxidant activity index (AAI) is calculated as follows:

$$\text{AAI} = \frac{\text{DPPH}(\mu\text{g/mL})}{\text{IC}_{50}(\mu\text{g/mL})}. \quad (2)$$

2.3.2. Nitric Oxide Scavenging Activity. The nitric oxide (NO) scavenging activity of various extracts was determined by Oskoueian et al. [19]. In a 96-well flat-bottomed plate, sixty microliters of the sample that had been diluted two-fold were combined with sixty microliters of sodium nitroprusside dissolved in phosphate-buffered saline (PBS) at a concentration of ten millimoles per liter. This mixture was then incubated at room temperature for 150 minutes while exposed to light. In the end, an equal volume of the Griess reagent was added to each well to measure the NO content. Ascorbic acid was used as a positive control in this experiment. The NO scavenging activity was calculated according to the formula: $[(A_0 - A_1)/A_0] \times 100\%$, where A_0 was the absorbance of the control reaction and A_1 was the absorbance in the presence of the sample. The assay was carried out in triplicate, and the data were reported as mean \pm standard deviation. IC₅₀ analysis was performed using GraphPad Prism 8 software.

2.4. In Vitro Antiplasmodial Assays. In this analysis, chloroquine-sensitive (3D7) and chloroquine-resistant (W2) strains of *Plasmodium falciparum* were utilized. A total of 1 mg of the sample was dissolved in 100 μL of DMSO (stock solution, 10,000 $\mu\text{g}/\text{mL}$ concentration), and a serial dilution was made from the stock solution. The parasites used in this test were synchronous (ring stage) parasites with less than one percent parasitemia. Then, 198 μL of parasite was added (the final concentrations of test material were 100 $\mu\text{g}/\text{mL}$, 10 $\mu\text{g}/\text{mL}$, 1 $\mu\text{g}/\text{mL}$, 0.1 $\mu\text{g}/\text{mL}$, and 0.01 $\mu\text{g}/\text{mL}$). The plate was placed within the chamber, and gas was blended (O_2 5%, CO_2 5%, and N_2 90%). Forty-eight hours were spent incubating the chamber containing the plate at 37°C. The culture was then harvested, and a 20% Giemsa-stained blood layer formed. Under a microscope, the number of infected erythrocytes per 1000 healthy erythrocytes was determined for the performed blood test. DMSO was used as a negative control and artemisinin as a positive control (standard drug) to avoid false results. Then, the percentage of growth and inhibition was determined by analyzing the data. Based on the

percent inhibition results, a statistical analysis using a probit analysis in Minitab®19 was conducted to determine the IC₅₀ value or concentration of the test material that could inhibit the parasite's growth by up to 50%. The assay was carried out in triplicate, and the data were reported as mean \pm standard deviation.

2.5. Toxicity Assay. The brine shrimp lethality test was utilized to determine the toxicity level exhibited by the extracts (BSLT). After preparing ten vials, each containing 2 mL of seawater, for the purpose of conducting the test method, a two-fold dilution was performed to generate several different concentrations. After applying an aliquot (0.1 mL) containing approximately ten nauplii to each vial, the setup was allowed to continue for twenty-four hours. After twenty-four hours, the contents of each bottle were examined, and dead larvae were tallied. As a form of negative control, dimethyl sulfoxide, or DMSO, was utilized. The assay was carried out in triplicate, and the data were reported as mean \pm standard deviation. By plotting the median mortality percentage against the log of concentration, we could determine the concentration (LC₅₀) that resulted in fifty percent of the population dying as a direct result of being exposed to the extracts.

2.6. Antibacterial Activity

2.6.1. Microbial Strain. Seven microorganisms, namely, *Bacillus cereus* (ATCC 10876), *B. subtilis* (ATCC 19659), *Escherichia coli* (ATCC 8739), *Listeria monocytogenes* (ATCC 7644), *Staphylococcus aureus* (ATCC 6538), *Salmonella typhimurium* (ATCC 142028), and *Vibrio parahaemolyticus* (ATCC 17802) were used in this study. The microbial isolates were maintained on an agar slant at four °C in the Laboratory of Biochemistry, Department of Chemistry, Universitas Riau. Before any antimicrobial test, the strains were subcultured for 24 hours on a fresh agar plate.

2.6.2. Antibacterial Activity. Subsequently, 100 μL of nutrient broth containing pathogenic bacteria was inoculated in nutrient agar. The extracts were screened for their antibacterial activity against those pathogenic bacteria by using the agar-disc diffusion method with a cell suspension of about 1.5×10^6 CFU/mL obtained from a McFarland turbidity standard No 0.5. A 6 mm sterile filter paper disc was immediately placed into the inoculated media, and each disc was filled with 20 μL extracts with a final concentration of 500 $\mu\text{g}/\text{disc}$ with chloramphenicol (30 $\mu\text{g}/\text{disc}$) as a positive control, incubated overnight at 37°C. After 24 hours of incubation, the inhibition zone (in mm) was measured. The diameter (in millimeters) of the inhibition zone surrounding the well was measured using a caliper. All procedures were carried out in triplicate. Minitab®19 was used to analyze mean comparisons, and the data were presented as mean \pm SD.

2.7. Statistical Analysis. All tests were performed in triplicate and the obtained results are expressed as mean \pm SD. Data were analyzed using GraphPad Prism Version 9, and comparisons of the means were determined by one-way analysis of variance (one-way ANOVA) followed by the Tukey test. Values were significantly different if they fall below the significance level (0.05).

3. Results

3.1. Radical Scavenging Activity. This study's free radical scavenging activities were determined by using two chemical assays, namely, DPPH and nitric oxide radicals. The results showed that there was a significant difference between the extracts and the ethyl acetate extract depicted the highest inhibition toward DPPH radical with a percentage of inhibition value of $96.87\% \pm 0.67$, followed by dichloromethane at $95.21\% \pm 0.37$ at a concentration of $1000 \mu\text{g}/\text{mL}$, while the *n*-hexane extract had the lowest inhibition at $62.21\% \pm 1.09$.

In addition, the extracts' half-maximal inhibitory concentration (IC_{50}), along with ascorbic acid serving as a positive control, was evaluated in GraphPad Prism 9 using a dose-response curve (Table 1). The findings demonstrated that the ethyl acetate extract had the lowest IC_{50} among the extracts, with a value of $51.63 \pm 0.46 \mu\text{g}/\text{mL}$. Despite this, the findings showed that the ethyl acetate extract was still significantly less potent than ascorbic acid, which had an IC_{50} of $4.15 \pm 0.24 \mu\text{g}/\text{mL}$. In addition, the extracts' antioxidant activity index displayed a wide range of activity. According to Table 1, dichloromethane and ethyl acetate extracts have the highest antioxidant activity index, whereas *n*-hexane extracts have the lowest index. Even though ethyl acetate and dichloromethane had the highest index, the activity of these two compounds was regarded as being inferior to that of ascorbic acid.

Moreover, the radical scavenging activity of the extracts was examined concerning the nitric oxide (NO) radical, and the results showed that the extracts inhibited the NO radical in a manner dependent on the dose. On top of that, dichloromethane and ethyl acetate demonstrated high activity levels among the extracts, with their respective percentages of inhibition emerging at $83.40 \pm 0.45\%$ and $85.17 \pm 0.58\%$. Furthermore, the IC_{50} value of the extracts was analyzed in the same manner as IC_{50} for the DPPH assay. The ethyl acetate extract showed high activity with IC_{50} $60.30 \pm 0.65 \mu\text{g}/\text{mL}$, where the *n*-hexane possessed the lowest activity ($956.70 \pm 0.54 \mu\text{g}/\text{mL}$), and compared to the ascorbic acid, the extracts showed low activity toward NO radical.

3.2. In Vitro Antiplasmodial and Toxicity Assays. The antiplasmodial activity against chloroquine-sensitive 3D7 and chloroquine-resistant W2 strains of *P. falciparum* was evaluated in vitro, as was the toxicity using the brine shrimp lethality test. Each of the four extracts exhibited activity against *P. falciparum* in a dose-dependent manner and at $100 \mu\text{g}/\text{mL}$; ethyl acetate inhibited the growth of

P. falciparum by $69.85 \pm 0.22\%$, while dichloromethane extract showed a comparable activity level at $64.18 \pm 0.22\%$. The activity level observed with the ethyl acetate was about twice higher as the dichloromethane with IC_{50} of 11.06 ± 0.45 and $21.68 \pm 0.56 \mu\text{g}/\text{mL}$, respectively. However, these results show that the extracts exhibited lower activity than artemisinin, with an IC_{50} value of $0.0054 \pm 0.0013 \mu\text{g}/\text{mL}$. Furthermore, the extracts were applied with a brine shrimp lethality test to their toxicity level, and the findings showed that the extracts exhibited similar toxicity levels (Table 2).

3.3. Antibacterial Activity. The effect of the crude extracts on the growth of seven tested indicator bacteria is shown in Table 3. Ethyl acetate extract exhibited the best extract as it presented the highest inhibition against *B. cereus* ATCC 10876, *V. parahaemolyticus* ATCC 17802, *L. monocytogenes* ATCC 7644, and *S. Typhimurium* ATCC 14028, followed by the dichloromethane extract with greatest clear zone against *B. subtilis* ATCC 19659, and *S. aureus* ATCC 6538. The minor activity was shown by the *n*-hexane extract, and none of the crude extracts affected the growth of *E. coli* ATCC 10876.

4. Discussion

This study applied the solvent-solvent partition technique with a modified Kupchan partition to the methanol extract [20]. Separating a total MeOH extract from four large extracts, which are simplified mixes of different polarities, using only partitions made from nonmiscellaneous solvents, is the main objective of the Kupchan partition, a practical and straightforward method for initiating a purification procedure. It is possible for there to be significant differences in distribution between extracts of compounds with varying degrees of polarity [17], and then, the extracts were investigated for their antioxidant, antiplasmodial, and antibacterial properties.

An antioxidant is a substance that significantly retards or prevents the oxidation process. The antioxidant activity is indirectly measured by determining the rate at which oxidation processes are inhibited in the presence of an antioxidant. [21] Each extract was then analyzed for its antioxidant activity using the free radical (DPPH) scavenging method. DPPH (2, 2-diphenyl-1-picrylhydrazyl), an organic stable radical in crystalline form and solution, is often used to measure a compound's antiradical activity or extract. The ability of a compound or extract to get rid of free radicals is often linked to its ability to act as an antioxidant. [22].

Despite the widespread use of the DPPH method, the lack of standardization of the results makes it difficult to compare the antioxidant potency of various plant extracts and pure compounds. Scherer and Godoy, therefore, proposed a new antioxidant activity index (AAI). The AAI was calculated based on the mass of DPPH and the mass of the sample in the reaction, resulting in a constant for each sample regardless of the DPPH concentration and sample

TABLE 1: The IC₅₀ values of various extracts of *S. palustris* (Burm.f.) Bedd. in DPPH and nitric oxide scavenging activities.

Sample	IC ₅₀ (µg/mL)	DPPH	Nitric oxide
		Antioxidant activity index (AAI)	IC ₅₀ (µg/mL)
<i>n</i> -Hexane extract	233.60 ± 0.54	0.34 ± 0.044	956.70 ± 0.54
Dichloromethane extract	58.33 ± 0.24	1.37 ± 0.065	83.44 ± 0.22
Ethyl acetate extract	51.63 ± 0.46	1.55 ± 0.034	60.30 ± 0.65
Water extract	141.20 ± 0.75	0.57 ± 0.023	412.60 ± 0.47
Ascorbic acid	4.15 ± 0.24	19.28 ± 0.045	54.8 ± 0.76

used. The sample is considered to have a low level of antioxidant activity when the antioxidant index (AAI) is less than 0.50. When the AAI falls between 0.5 and 1.0, the sample's antioxidant activity is considered moderate. When the AAI is between 1.0 and 2.0, it is considered that the sample has a high level of antioxidant activity. [15] According to these standards, the ethyl acetate and dichloromethane extracts should be considered to have a high level of antioxidant activity.

The radical scavenging activities of these extracts might be the presence of secondary metabolites, specifically phenolics, and flavonoids. Flavonoids might have multiple properties for removing reactive oxygen and nitrogen species. Typically, the presence of an ortho-hydroxylation on the B-ring of the flavonoid molecule, the number of free hydroxyl groups, a C2-C3 double bond in the C-ring, or the presence of a 3-hydroxyl group is listed as a condition of antioxidant and antiradical activity. [23] According to findings from earlier research, *S. palustris* (Burm.f.) Bedd. contains acylated and nonacylated kaempferol glycosides, each of which possesses free radical scavenging activity toward the DPPH radical at IC₅₀ concentrations ranging from 120 to 400 µM. [14] On the basis of these findings, it is possible to conclude that the extracts of *S. palustris* (Burm.f.) Bedd., more specifically dichloromethane and ethyl acetate, can inhibit free radicals. These findings are consistent with a few other publications that have been done in the past. According to the findings of Arullappan et al., ethyl acetate extracted from the leaves of *S. palustris* (Burm.f.) Bedd., which originated in Malaysia, exhibited DPPH radical inhibitor activity with an IC₅₀ value of 650 µg/mL. In addition, the ethyl acetate extract of *S. palustris* (Burm.f.) Bedd. from Perak, Malaysia, demonstrated a high level of radical scavenging activity, and its IC₅₀ concentration was found to be 49.8 µg/mL, which is comparable to the results found in the report [5, 10].

Furthermore, to explore the potential of the various extracts toward free radicals, nitric oxide (NO) inhibitor was determined. Nitric oxide, also known as NO, is a potent pleiotropic mediator produced from the amino acid L-arginine by vascular endothelial cells, phagocytes, and specific brain cells. When NO combines with the superoxide radical, a highly reactive peroxy nitrite anion known as ONOO⁻ is produced. This is when NO's toxicity turns for the worse [24]. The reaction between the nitric oxide produced by sodium nitroprusside and oxygen results in the formation of nitrite. The nitrite ions diazotize with sulfanilic acid and couple with naphthyl ethylenediamine, forming a pink color

with a wavelength of 546 nm. [25] Table 1 shows the corresponding IC₅₀ values for NO scavenging activity, and the dichloromethane and ethyl acetate extracts had IC₅₀ values less than 200 µg/ml, indicating good NO-scavenging activity, whereas the *n*-hexane and water extracts had IC₅₀ values greater than 400 µg/ml, indicating poor NO scavenging activity. The NO scavenging values were classified based on Tsai et al. [26].

Moreover, the free radical scavenging activity of *S. palustris* (Burm.f.) Bedd. extracts, specifically the dichloromethane and ethyl acetate against the nitric oxide (NO) radical, showed suitable activities. This is due to the presence of kaempferol glycoside within species. Multiflorins A, a kaempferol glycoside, isolated from fern *Neochiropteris palmatopedata* showed a percentage inhibition of NO 52% at 20 µg/ml, and kaempferol was also reported to inhibit the production of NO in RAW 264.7 cells line at IC₅₀ 90.3 µM [27, 28].

Oxidative stress is thought to play a role in developing several diseases, including cancer, diabetes, and renal disease. This stress is brought on by the excessive production of nitric oxide (NO), which can result from infection or inflammation. As a result, removing NO radicals or inhibiting NO production by mitogen-activated cells may be promising indicators for discovering new treatments for those ailments. [26] Consequently, as a result of these studies, the extract of this species can further have its potential inhibition in the cell line of animal studies investigated. The presence of a NO inhibitor originating from this species has not previously been documented. As a consequence of this finding, additional information regarding the free radical scavenging activities of the species is provided.

In addition, it was discovered that all four extracts possessed varying degrees of activity against chloroquine-sensitive 3D7 strains of *P. falciparum* and chloroquine-resistant W2 strains (Table 2). According to the WHO guidelines and the fundamental requirements of anti-parasitic drug discovery, the activities of the extracts were separated into four categories based on their values for the IC₅₀ concentration: high activity (IC₅₀ ≤ 5 µg/ml), promising activity (5 µg/ml < IC₅₀ ≤ 15 µg/ml), moderate activity (15 µg/ml < IC₅₀ ≤ 50 µg/ml), and poor activity (IC₅₀ > 50 µg/ml). [29, 30] Therefore, the dichloromethane and ethyl acetate extracts have a moderate and promising antiplasmodial activity, respectively, whereas the *n*-hexane and water extracts were classified as having a poor activity. This might be due to the presence of kaempferol glucoside, such as kaempferol 3-O-α-L-

TABLE 2: Antiplasmodial activity and toxicity of *S. palustris* (Burm.f.) Bedd. extracts.

Sample	Antiplasmodial (IC ₅₀ ; µg/mL)	Toxicity (LC ₅₀ ; µg/mL)
<i>n</i> -Hexane extract	>100	>1000
Dichloromethane extract	21.68 ± 0.56	>1000
Ethyl acetate extract	11.06 ± 0.45	>1000
Water extract	62.19 ± 0.34	>1000
Artemisinin (control)	0.0054 ± 0.0013	—

TABLE 3: Antibacterial of *S. palustris* (Burm.f.) Bedd. extracts at a concentration of 500 µg/disc for Gram-negative and Gram-positive bacteria.

Microorganisms	Inhibition zone (cm)				
	<i>n</i> -Hexane	Dichloromethane	Ethyl acetate	Water extract	Chloramphenicol
<i>Bacillus subtilis</i>	0.00 ± 0 ^c	1.30 ± 0.03^b	1.11 ± 0.01 ^c	0.73 ± 0.02 ^d	3.18 ± 0.02 ^a
<i>Bacillus cereus</i>	0.00 ± 0 ^d	0.85 ± 0.02 ^{bc}	0.90 ± 0.01^b	0.73 ± 0.02 ^c	2.57 ± 0.04 ^a
<i>Streptococcus aureus</i>	0.75 ± 0.02 ^d	0.93 ± 0.03^b	0.85 ± 0.01 ^c	0.73 ± 0.02 ^d	2.91 ± 0.03 ^a
<i>Vibrio parahaemolyticus</i>	0.00 ± 0 ^d	0.86 ± 0 ^c	0.93 ± 0^b	0.82 ± 0 ^c	2.15 ± 0 ^a
<i>Listeria monocytogenes</i>	0.00 ± 0 ^c	0.00 ± 0 ^c	0.84 ± 0.03^b	0.00 ± 0 ^c	2.51 ± 0.05 ^a
<i>Salmonella typhi</i>	0.00 ± 0 ^c	0.82 ± 0.02 ^b	0.83 ± 0.02^b	0.77 ± 0.01 ^b	2.67 ± 0.03 ^a
<i>Escherichia coli</i>	0.00 ± 0 ^b	0.00 ± 0 ^b	0.00 ± 0 ^b	0.00 ± 0 ^b	2.95 ± 0.01 ^a

^{a-c} means in the same row with different lowercase letters differed significantly ($p < 0.05$). **Bold:** the highest inhibition on each indicator microorganism.

rhamnopyranoside, isolated from *S. palustris* (Burm.f.) Bedd. leave. This compound has been reported that possessed promising antiplasmodial activity against chloroquine-resistant *P. falciparum* with an IC₅₀ of 106 µM [14, 31]. This finding is supported by a previous report that determined the antimalarial activity of an ethanol extract of *S. palustris* (Burm.f.) Bedd. leaves against parasitemia and splenomegaly of *Plasmodium berghei* ANKA in infected BALB/c mice. The results indicated that the extract possessed good antimalarial activity at an ED₅₀ value of 77.05 mg/kg body weight [32].

The extracts were put through a lethality test using brine shrimp to determine their overall level of toxicity. The test is a straightforward and highly effective method for determining the toxic potential of bioactive substances or extracts. It is based on the ability of test compounds to kill shrimp, a relatively simple zoological organism (*Artemia Salina*) [33]. According to Meyer and the co-author, the plant extracts demonstrated toxicity with an LC₅₀ value of less than 1000 µg/ml; however, the plant extracts were regarded as highly toxic for the shrimp with an LC₅₀ value of less than 30 µg/ml [34]. The findings showed a nontoxic LC₅₀ value of >1000 µg/ml for all the extracts (Table 2). This result contradicts the previous report, which mentions that *S. palustris* (Burm.f.) Bedd. ethanolic extract growth in Bangladesh showed the LC₅₀ at 184.7 µg/ml using the BSLT method. [35] The toxicity of brine shrimp has been found to predict cytotoxicity and parasite activity. There was a strong correlation between BSLT toxicity and cytotoxicity against the 9 KB cell line (human nasopharyngeal carcinoma) and other tumors and *in vivo* murine leukemia. The BSLT detects potent anticancer activity, but its predictive ability to distinguish between potent, moderate, and weak anticancer compounds is limited. Thus, the BSLT provides a rapid screening for potent cytotoxins and a higher level of cancer discrimination. [36].

In general, the majority of the tested extracts exhibited weak antibacterial activity. As shown in Table 3, extracts at 500 µg/disc exhibited variable inhibitory activity against all bacteria, with inhibition zone diameters ranging from 0.7 to 1.5 cm. The responses of microorganisms to the different extracts varied. In this study, it was determined that all extracts inhibited Gram-positive bacteria more effectively than Gram-negative bacteria. As supported by these results, Gram-negative microorganisms are typically more resistant to antimicrobial agents than Gram-positive bacteria. This has long been attributed to an outer-membrane permeability barrier in Gram-negative bacteria, which prevents antimicrobial agents from reaching their targets within the bacterial cells [37]. Even though the antibacterial activity from the extract exhibited weak antibacterial, it is noteworthy that the finding from Liu et al. showed that stenopalustrosides A-D showed significant antibacterial activities against Gram-positive strains (*Bacillus cereus*, *Staphylococcus epidermidis*, *S. aureus*, and *Micrococcus luteus*) with MIC between 2 and 64 µg/mL [3]. Another report showed that methanol extract from *S. palustris* (Burm.f.) Bedd. exhibited various antibacterial activities against the pathogenic bacteria with a MIC range between 50 and 12.5 mg/mL [8].

The species, on the whole, demonstrated a wide range of biological activities throughout the subject of this investigation. According to the findings of the study, the species exhibit a potential activity by scavenging free radicals (DPPH and nitric oxide), and the ability to inhibit nitric oxide radicals in this species was reported for the first time. Aside from that, it was the first time that the antiplasmodium activity of this species was reported, and it has a good potential to inhibit the growth of both strains of *plasmodium*. On the other hand, this particular species does not exhibit significant antibacterial activity. The limitation of this study, however, is that we did not report the phytochemical compounds present in this species. Instead, we correlated the biological activities revealed in this study with

compounds already known to be present in the species based on previously published research.

5. Conclusions

The results of this study have demonstrated that an extract of *S. palustris* possesses promising antioxidant and antiplasmodial activities. The dichloromethane and ethyl acetate extracts showed significant levels of antiplasmodial and radical scavenging activities. The antibacterial activities of the extracts, on the other hand, showed only a weak level of activity against pathogenic bacteria. These findings provide a basis for further investigation into the isolating and analyzing the biological activity of secondary metabolites found in the extracts.

Data Availability

The datasets used and/or analyzed during this study are included in the article. Further clarification can be obtained from the corresponding author.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

R.H. and H.Y.T. were responsible for conceptualization, A.S.N. was responsible for methodology, R.H. and R.A. were responsible for validation, Y.H. was responsible for formal analysis, R.K. and M.A. were responsible for investigation, R.H. and A.S.N. wrote the original draft, N.F. and A.S.N. reviewed and edited the article, R.H. was responsible for visualization, H.Y.T. was responsible for supervision, R.K. and Y.H. were responsible for project administration, and R.H. was responsible for funding acquisition. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

The authors would like to thank the Center of Research and Community Development, University of Riau and Ministry of Research, Technology, and Higher Education of the Republic of Indonesia for their support (1635/UN19.5.1.3/PT.01.03/2022). This research was funded by Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, and Ministry of Research, Technology, and Higher Education, grant number 1635/UN19.5.1.3/PT.01.03/2022.

References

- [1] M. Almurdani, A. Zamri, T. T. Nugroho et al., "Antioxidant and antidiabetic activities of mepening (lithocarpus bancanus) leaves," *Pharmacognosy Journal*, vol. 12, no. 2, pp. 328–334, 2020.
- [2] H. Cao, T.-T. Chai, X. Wang et al., "Phytochemicals from fern species: potential for medicine applications," *Phytochemistry Reviews*, vol. 16, no. 3, pp. 379–440, 2017.
- [3] H. Liu, J. Orjala, O. Sticher, and T. Rali, "Acylated flavonol glycosides from leaves of *Stenochlaena palustris*," *Journal of Natural Products*, vol. 62, no. 1, pp. 70–75, 1999.
- [4] N. Sofiyanti, D. Iriani, D. Fitmawati, and A. A. Roza, "Stenochlaena riauensis (Blechnaceae), a new fern species from Riau, Indonesia," *Bangladesh Journal of Plant Taxonomy*, vol. 22, no. 2, pp. 137–141, 2015.
- [5] S. Arullappan, S. Sawai, L. A. Chee, M. Mahandan, and R. Shanmugavelan, "Phytochemical screening and evaluation of cytotoxic effect and antioxidant activity of fractions isolated from *Stenochlaena palustris* (Burm.f.) Bedd. Leaves," *Indian Journal of Pharmaceutical Education and Research*, vol. 51, no. 4s, pp. s735–s740, 2017.
- [6] A. Benjamin and V. Manickam, "Medicinal pteridophytes from the western Ghats," *Indian Journal of Traditional Knowledge*, vol. 6, no. 4, pp. 611–618, 2007.
- [7] T.-T. Chai, E. Panirchellvum, H.-C. Ong, and F.-C. Wong, "Phenolic contents and antioxidant properties of *Stenochlaena palustris*, an edible medicinal fern," *Botanical Studies*, vol. 53, no. 4, pp. 439–446, 2012.
- [8] Z. Zuraini, S. Sasidharan, S. R. Kaur, and M. Nithiyayini, "Antimicrobial and antifungal activities of local edible fern *Stenochlaena palustris*," (*Burm. F.*) *Bedd. Pharmacology online*, vol. 1, pp. 233–237, 2010.
- [9] G. Novita, S. Kusmardiyani, and I. Fidrianny, "Antioxidant activities from various extracts of different parts of kelakai (*Stenochlaena palustris*) grown in central Kalimantan-Indonesia," *Asian Journal of Pharmaceutical and Clinical Research*, vol. 9, no. Suppl 2, pp. 215–219, 2016.
- [10] T.-T. Chai, M.-T. Kwek, H.-C. Ong, and F.-C. Wong, "Water fraction of edible medicinal fern *Stenochlaena palustris* is a potent α -glucosidase inhibitor with concurrent antioxidant activity," *Food Chemistry*, vol. 186, pp. 26–31, 2015.
- [11] V. Sumathy, S. Jothy Lachumy, Z. Zuraini, and S. Sasidharan, "Effects of *Stenochlaena palustris* leaf extract on growth and morphogenesis of food borne pathogen, *Aspergillus Niger*," *Malaysian journal of nutrition*, vol. 16, no. 3, pp. 439–446, 2010.
- [12] N. J.-Y. Chear, K.-Y. Khaw, V. Murugaiyah, and C.-S. Lai, "Cholinesterase inhibitory activity and chemical constituents of *Stenochlaena palustris* fronds at two different stages of maturity," *Journal of Food and Drug Analysis*, vol. 24, no. 2, pp. 358–366, 2016.
- [13] H. Liu, J. Orjala, T. Rali, and O. Sticher, "Glycosides from *Stenochlaena palustris*," *Phytochemistry*, vol. 49, no. 8, pp. 2403–2408, 1998.
- [14] N. J.-Y. Chear, A. N. Fauzi, K.-Y. Khaw, S.-B. Choi, N. S. Yaacob, and C.-S. Lai, "Free radical scavenging and cytotoxic properties of acylated and non-acylated kaempferol glycosides from *Stenochlaena palustris*: a perspective on their structure-activity relationships," *Pharmaceutical Chemistry Journal*, vol. 53, no. 3, pp. 188–193, 2019.
- [15] R. Scherer and H. T. Godoy, "Antioxidant activity index (AAI) by the 2, 2-diphenyl-1-picrylhydrazyl method," *Food Chemistry*, vol. 112, no. 3, pp. 654–658, 2009.
- [16] R. Hendra, R. Khodijah, R. Putri et al., "Cytotoxicity and antiplasmodial properties of different *Hylocereus polyrhizus* peel extracts," *Medical Science Monitor Basic Research*, vol. 27, 2021.
- [17] R. Hendra, S. N. Gurning, U. P. A. Panjaitan, and H. Y. Teruna, "Antioxidant activity of an epiphyte fern in palm oil tree," *Journal of Physics Conference Series*, vol. 1655, 2020.

- [18] R. Hendra, L. Masdeatresa, R. Abdulah, and Y. Haryani, "Red dragon peel (*Hylocereus polyrhizus*) as antioxidant source," *AIP Conference Proceedings*, AIP Publishing LLC, College Park, MD, USA, 2020.
- [19] E. Oskoueian, N. Abdullah, R. Hendra, and E. Karimi, "Bioactive compounds, antioxidant, xanthine oxidase inhibitory, tyrosinase inhibitory and anti-inflammatory activities of selected agro-industrial by-products," *International Journal of Molecular Sciences*, vol. 12, no. 12, pp. 8610–8625, 2011.
- [20] S. M. Kupchan, G. Tsou, and C. W. Sigel, "Daticascacin, a novel cytotoxic cucurbitacin 20-acetate from *Datisca glomerata*," *Journal of Organic Chemistry*, vol. 38, no. 7, pp. 1420–1421, 1973.
- [21] F. Shahidi and Y. Zhong, "Measurement of antioxidant activity," *Journal of Functional Foods*, vol. 18, pp. 757–781, 2015.
- [22] S. B. Kedare and R. P. Singh, "Genesis and development of DPPH method of antioxidant assay," *Journal of Food Science & Technology*, vol. 48, no. 4, pp. 412–422, 2011.
- [23] S. Burda and W. Oleszek, "Antioxidant and antiradical activities of flavonoids," *Journal of Agricultural and Food Chemistry*, vol. 49, no. 6, pp. 2774–2779, 2001.
- [24] B. S. Taylor, Y.-M. Kim, Q. Wang, R. A. Shapiro, T. R. Billiar, and D. A. Geller, "Nitric oxide down-regulates hepatocyte-inducible nitric oxide synthase gene expression," *Archives of surgery*, vol. 132, no. 11, pp. 1177–1183, 1997.
- [25] D. M. Nagmoti, D. K. Khatri, P. R. Juvekar, and A. R. Juvekar, "Antioxidant activity free radical-scavenging potential of *Pithecellobium dulce* Benth seed extracts," *Free Radicals and Antioxidants*, vol. 2, no. 2, pp. 37–43, 2012.
- [26] P. J. Tsai, T. H. Tsai, C. H. Yu, and S. C. Ho, "Comparison of NO-scavenging and NO-suppressing activities of different herbal teas with those of green tea," *Food Chemistry*, vol. 103, no. 1, pp. 181–187, 2007.
- [27] J.-H. Yang, T. P. Kondratyuk, L. E. Marler et al., "Isolation and evaluation of kaempferol glycosides from the fern *Neocheiropteris palmatopedata*," *Phytochemistry*, vol. 71, no. 5-6, pp. 641–647, 2010.
- [28] H. S. Rho, A. K. Ghimeray, D. S. Yoo et al., "Kaempferol and kaempferol rhamnosides with depigmenting and anti-inflammatory properties," *Molecules*, vol. 16, no. 4, pp. 3338–3344, 2011.
- [29] D. A. Fidock, P. J. Rosenthal, S. L. Croft, R. Brun, and S. Nwaka, "Antimalarial drug discovery: efficacy models for compound screening," *Nature Reviews Drug Discovery*, vol. 3, no. 6, pp. 509–520, 2004.
- [30] O. Jansen, M. Tits, L. Angenot et al., "Anti-plasmodial activity of *Dicoma tomentosa* (Asteraceae) and identification of urospermal A-15-O-acetate as the main active compound," *Malaria Journal*, vol. 11, no. 1, p. 289, 2012.
- [31] M. I. Barliana, E. W. Suradji, R. Abdulah et al., "Anti-plasmodial properties of kaempferol-3-O-rhamnoside isolated from the leaves of *Schima wallichii* against chloroquine-resistant *Plasmodium falciparum*," *Biomedical Reports*, vol. 2, no. 4, pp. 579–583, 2014.
- [32] L. N. Azizah, P. Wardhani, and H. Arwati, "Antimalarial activity of ethanol extract of kelakai leaves (*Stenochlaena palustris*) to parasitemia and splenomegaly in BALB/c mice infected with plasmodium berghei ANKA," *JUXTA: Jurnal Ilmiah Mahasiswa Kedokteran Universitas Airlangga*, vol. 13, no. 1, pp. 1–5, 2022.
- [33] C. Wu, "An important player in brine shrimp lethality bioassay: the solvent," *Journal of Advanced Pharmaceutical Technology & Research*, vol. 5, no. 1, pp. 57–58, 2014.
- [34] B. Meyer, N. Ferrigni, J. Putnam, L. Jacobsen, D. Nichols, and J. McLaughlin, "Brine shrimp: a convenient general bioassay for active plant constituents," *Planta Medica*, vol. 45, no. 05, pp. 31–34, 1982.
- [35] S. L. Debnath, P. Kundu, M. F. Ahad, L. Saha, N. N. Biswas, and S. K. Sadhu, "Investigation of phytochemical and pharmacological assessment of ethanol extract of *Stenochlaena palustris*—an edible fern of Sundarbans," *Journal of Medicinal Plants Studies*, vol. 9, no. 3, pp. 226–232, 2021.
- [36] E. L. Ghisalberti, "Detection and isolation of bioactive natural products," *Bioactive Natural Products: Detection, Isolation, and Structural Determination*, CRC Press, Boca Raton, FL, USA, 2008.
- [37] F. Othman, *Chemical Constituents and Biological Activities of Flavonoids from Hydroponically Grown Pegaga (Centella Asiatica, Linn., Urban) Extracts*, Universiti Putra Malaysia Serdang, Seri Kembangan, Malaysia, 2003.