

ANALYSIS OF FACTORS THAT RELATED MATERNAL SEVERE PREECLAMPSIA TO THE ASPHYXIA OF NEW BORN BABY IN SOEBANDI HOSPITAL JEMBER REGENCY

Ida Srisurani Wiji Astuti, Medical Faculty, University of Jember; dr.rani82@yahoo.com

ABSTRACT

Background: Preeclampsia is a disorder of pregnancy characterized by high blood pressure and a large amount of protein in the urine. Including the literacy, the most horrible effect of maternal severe preeclampsia was the asphyxia of newborn baby. It could happen until 44 % birth. In 2012, Soebandi hospital noted that 423 mother with preeclampsia which 319 were severe preeclampsia. The purpose of this study was to determine the picture of severe preeclampsia with the asphyxia of newborn baby and analyze the factors that influence of it. That factors were quality of antenatal care, the number of births, and mother's body mass index. Methods: This study used retrospective analytic research using secondary data on maternal and neonatal care Soebandi hospital Jember Regency. The sample size of this research were 115 pregnant womens who were selected by purposive sampling on the 2015. Non-parametric chi-square test was used to analyze data. Results: The quality of antenatal care was taken from pearson chi square 1,875 with p value 0,171. Based on the number of births, p value was 0,231. Conversely, there were relationship that body mass index influenced severe preeclampsia and asphyxia of newborn baby (p value 0,025). It enabled the occurrence of dangerous pregnancy complications for mother and the infant. Conclusion: factor that influenced severe eclampsia and asphyxia in newborn baby is mother's body mass index. It means when the mother has high BMI (obesity), the complication of pregnancy can occur anytime like severe preeclampsia and asphyxia of the newborn baby. Quality of antenatal care and number of births do not prove relation between it.

Keyword: severe preeclampsia, asphyxia of the newborn baby

Introduction

An American study (Nawal,M. 2008) said that "Approximately 529,000 women die from pregnancy-related causes annually and almost all (99%) of these maternal deaths occur in developing nations. One of the United Nations' Millennium Development Goals is to reduce the maternal mortality rate by 75% by 2015. Causes of maternal mortality include postpartum hemorrhage, eclampsia, obstructed labor, and sepsis. Many developing nations lack adequate health care and family planning, and pregnant women have minimal access to skilled

labor and emergency care. Basic emergency obstetric interventions, such as antibiotics, oxytocics, anticonvulsants, manual removal of placenta, and instrumented vaginal delivery, are vital to improve the chance of survival."

Nationally, East Java province occupies the top five the rising trend in maternal mortality continues to occur in East Java. There were 487 cases in 2008, 535 cases in 2009, 598 cases in 2010 and 627 cases in 2011. The area of 'Tapal Kuda' contributes the highest maternal mortality rate, particularly Jember,

Banyuwangi, Situbondo, Bondowoso, and Lumajang. Based on data from Health Profile of East Java Province in 2011, Jember occupied the top rank of 38 regencies/cities in East Java. 54 cases of maternal death illustrate the poor state. It can be said that for four consecutive years (2008-2011), Jember Regency had not shifted from the first rank (Dinkes Propinsi Jatim, 2012).

In 2012, from January 1st until December 31th Soebandi hospital noted that 423 mothers with preeclampsia which 319 were severe preeclampsia (Adawiyah, 2014).

Perinatal outcome assessment could be done by using Apgar Score (AS). Less value AS which baby detected in the first minute did not endorse the out come in the future. A retrospective research concluding that AS in the early five minute still being important predictor for neonatal deaths, but it cannot use to know about long life outcome. In other data, amount 13.399 the premature baby (less than 26-36 week) revealed neonatal deaths higher (315/ 1000) with AS 0-3 in the first minute than (5/1000) with AS 5-7 in the early five minutes (Brian, M, et al. 2014).

Equal to the imunologic theory by Sudhaberata (2001), due to the mother on the first pregnant will be formed blocking antibodies. In other side, in the first pregnant being performed Human Leucocyte Antigen Protein G (HLA) which conducted maternal imunity. This can make mother rejects product of conception (placenta) or being intolerated between mother and placenta wich can conduct preeclampsia. Beside that, Desfauza (2008) claimed that any relationship significantly between quantity of give birth with asphyxia of the newborn baby. Data showed mother with 1 - >3 times give birth (60,5% asphyxia)

more higher risk than 2 - 3 times give birth (36,7% asphyxia).

Gestational nutritional status very important to reach well being health status for mother and baby. Women with low body mass index can get negative effect when labor such as preterm and low weight body of the baby (Papathakis, 2005). Conversely, the women with over BMI gets more high risk in gestational such as abortion, labor with sectio caesare, preeclampsia, trombosis, perinatal deaths, and macrosomia (Yu, 2006).

Methods

This study is a retrospective analytic research using secondary data on maternal neonatal care Soebandi hospital Jember regency. The sample size of this research were 115 pregnant womens who were selected by purposive sampling on the 2015.

Inclusion factors :

1. Mother has gotten medicinal check up on January-October 2015 with severe preeclampsia in Soebandi hospital.
2. Mother with aterm pregnant (37-42 weeks).
3. Mother with spontaneous labor by lies the head of baby.
4. Mother has medical record before the research was conducted.

Exclusion factors:

Everything that makes asphyxia of the newborn baby is exclusion factor, such us:

1. Mother → **abnormal bleeding**, postdate pregnant, longterm labor, and uterine rupture.
2. Placenta → **nuchal cord**, **short umbilical cord**, and prolapse baby's cord.
3. Baby → **preterm**, **congenital abnormality**, and positional abnormality.

Variable dependent in this study is the newborn baby with asphyxia from mother with severe preeclampsia. Variable independent are quality of antenatal care, the number of births, and mother's body mass index.

The analysis method used univariate analysis and bivariate analysis. Univariate methods consisted of samples frequency distribution. Meanwhile, the bivariate methods used non-parametric chi-square analysis to determine the effect between variables.

Results and discussion

Tabel 1. Distribution ANC on severe preeclampsia to asphyxia neonatorum 9

	Infant			
	Asphyxia		Non	
	Amount (person)	%	Amount (person)	%
Good	5	25%	15	75%
Bad	5	50%	5	50%
Total	10	33,3%	20	66,7%

The last research from Faiqoh (2014), declared about maternal severe preeclampsia occurs 42 (85,7%) because they did not obey midwifery advice when they do ante natal care (ANC). Only 7 (14,29%) mothers with severe preeclampsia do the advices from the medicine doctor or midwife. Conversely, the control group said that 45 (95,9%) mothers obey and only 2 (4,08%) did not obey.

Including Wiknjosastro (2008), give birth quantity is the one risk factor wich related preeclampsia. Primigravida has higher frequency than multigravida.

Tabel 2. Distribution of Body Mass Index 7

Kategori	Jumlah	Persentase
Thin	4	11,8%
Normal	11	32,4%
Fat	19	55,9%

Total	34	100,0%
-------	----	--------

Tabel 3. Analysis chi square of Body Mass Index 7

	Value	Df	Asymptotic Significance (2-sided)
Person Chi-Square	5,261 ^a	2	,072
Likelihood Ratio	7,594	2	,022
Linear-by-Linear Association	2,010	1	,156
N of Valid Cases	34		

Blomberg research (2013) noted which one thing that caused asphyxia neonatorum is body mass index of the mother when she was pregnant. Obesity in pregnant (BMI>25) caused 3 times higher risk than normal BMI. On the other side, mother with low BMI (18,5) had less risk in asphyxia neonatorum.

References

Adawiyah, R. 2014. *Studi Penggunaan Obat pada Penderita Eklamsi Berat di RSD dr.Soebandi Jember*. Tidak diterbitkan. Skripsi. Jember: Repository UNEJ.

Blomberg, M. 2013. Maternal Obesity, Mode of Delivery, and Neonatal Outcome. *Obstet Gynecol*. 122: 50-5.

Brian, M., Casey, Donald, D., Kenneth J. Leveno. the Continuing Value of the APGAR Score for Assessment of Newborn Infant. *NEJM*.

Desfaufa, E. 2008. *Faktor-faktor yang Mempengaruhi Terjadinya Asfiksia Neonatorum pada Bayi Baru Lahir yang Dirawat di RSU dr. Pirngadi Medan Tahun 2007*. Medan: Repository USU.

- Dinas Kesehatan Propinsi Jawa Timur. (2012). *Profil Kesehatan Propinsi Jawa Timur Tahun 2007-2011*. Surabaya.
- Faiqoh, E., Hendrawati, L. Y. 2014. Hubungan karakteristik Ibu, ANC dan Kepatuhan Perawatan Ibu hamil dengan Terjadinya Preeklamsia. *Jurnal Berkala Epidemiologi*, Vol. 2, No. 2 Mei 2014: 216-226.
- Habibi, B.Y. 2016. Hubungan Indeks Massa Tubuh Penderita Preeklamsia Berat terhadap Kejadian Asfiksia Neonatorum di RSD. Dr. Soebandi. Skripsi. Tidak diterbitkan. Jember: Repository UNEJ.
- Hashemi, A. 2015. Hubungan Paritas pada Penderita Preeklamsia Berat terhadap Kejadian Asfiksia Neonatorum di RSD dr. Soebadi Jember. Skripsi. Tidak diterbitkan. Jember: Repository UNEJ.
- Mufidah, A. D. 2015. Perbedaan Kualitas Antenatal Care pada Ibu dengan Preeklamsia Berat terhadap Kejadian Asfiksia Neonatorum di RSD dr. Soebandi Kabupaten Jember. Skripsi. Tidak diterbitkan. Jember: Repository UNEJ.
- Nawal, M Nour. (2008). An Introduction to Maternal Mortality. *Journal of reviews obstetrics and gynecology*. Vol. 1(2):77-81.
- Papathakis, P., Rollins, N. 2005. *HIV and Nutrition: Pregnant and Lactating Women*. South africa: WHO Department of Nutrition for Health and Development.
- Sudhaberata, K. 2001. Penanganan Preeklamsia Berat dan Eklamsia. Cermin Dunia Kedokteran. Nomor 133. Jakarta: Penerbit PT.Kalbe Farma.
- Wiknjosastro, H. 2008. Ilmu Kebidanan Edisi Keempat. Jakarta: PT. Bina Pustaka Sarwono Prawirohardjo.
- Yu, C.K.H., Teoh, T.G., Robinson, S. 2006. Obesity in Pregnancy. *BJOG An International Journal of Obstetric and Gynaecology*. 113: 1117-25.