Volume 9

Number 8

August 2018



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development Scopus coverage years: from 2010 to 2017 Publisher: R.K. Sharma, Institute of Medico-Legal Publications ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine: Public Health, Environmental and Occupational Health

CiteScore 2015-0.02 SJR 2015-0.105 SNIP 2015-0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof. Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohinee Delhi

NTERNATIONAL EDITORIAL ADVISORY BOARD

Dr. Abdul Rashid Khan B. Md Jagar Din, (Associate Professor) Department of Public Health Medicine, Penang Medical College, Penang, Malaysia

Dr. V Kumar (Consulting Physician) Mount View Hospital, Las Vegas, USA

Basheer A. Al-Sum,

Botany and Microbiology Deptt, College of Science, King Saud

University, Riyadh, Saudi Arabia

Dr. Ch Vijay Kumar (Associate Professor)

Public Health and Community Medicine, University of Buraimi, Oman

Dr. VMC Ramaswamy (Senior Lecturer)

Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur

Kartavya J. Vyas (Clinical Researcher) Department of Deployment Health Research,

Naval Health Research Center, San Diego, CA (USA)

Prof. PK Pokharel (Community Medicine) BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

Dr. Anju Ade (Associate Professor) Navodaya Medical College, Raichur, Kamataka

Dr. E. Venkata Rao (Associate Professor) Community Medicine,

Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.

Dr. Amit K. Singh (Associate Professor) Community Medicine,

VCSG Govt. Medical College, Srinagar - Garhwal, Uttarakhand

Dr. R G Viveki (Professor & Head) Community Medicine,

Belgaum Institute of Medical Sciences, Belgaum, Kamataka

Dr. Santosh Kumar Mulage (Assistant Professor) Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Kamataka

Dr. Gouri Ku. Padhy (Associate Professor) Community and

Family Medicine, All India Institute of Medical Sciences, Raipur

Dr. Ritu Goyal (Associate Professor)

Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar

Dr. Anand Kalaskar (Associate Professor)

Microbiology, Prathima Institute of Medical Sciences, AP

Dr. Md. Amirul Hassan (Associate Professor)

Community Medicine, Government Medical College, Ambedkar Nagar, UP

Dr. N. Girish (Associate Professor) Microbiology, VIMS&RC, Bangalore

Dr. BR Hungund (Associate Professor) Pathology, JNMC, Belgaum.

Dr. Sartaj Ahmad (Associate Professor),

Medical Sociology, Department of Community Medicine, Swami Vivekananda

Subharti University, Meerut, Uttar Pradesh, India

Dr Sumeeta Soni (Associate Professor)

Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

Dr Manoj Kumar Tripathi (Associate Professor), Political Science and public administration, Swami Vivekananda Subharti University. Meerut, Uttar Pradesh, India.

NATIONAL EDITORIAL ADVISORY BOARD

Prof. Sushanta Kumar Mishra (Community

Medicine) GSI Medical College - Raiahmundry Kamataka

Prof. D.K. Srivastava (Medical Biochemistry)

Jamia Hamdard Medical College, New Delhi

Prof. M Sriharibabu (General Medicine) GSL Medical College,

Rajahmundry, Andhra Pradesh

Prof. Pankaj Datta (Principal & Prosthodentist)

Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

Prof. Samarendra Mahapatro (Pediatrician)

Hi-Tech Medical College, Bhubaneswar, Orissa

Dr. Abhiruchi Galhotra (Additional Professor) Community and Family

Medicine, All India Institute of Medical Sciences, Raipur

Prof. Deepti Pruthvi (Pathologist) SS Institute of Medical Sciences &

Research Center, Davangere, Karnataka

Prof. G S Meena (Director Professor)

Maulana Azad Medical College, New Delhi

Prof. Pradeep Khanna (Community Medicine)

Post Graduate Institute of Medical Sciences, Rohtak, Haryana

Dr. Sunil Mehra (Paediatrician & Executive Director) MAMTA Health Institute of Mother & Child, New Delhi

Dr Shailendra Handu, Associate Professor, Phrma, DM (Pharma,

Dr. AC. Dhariwal: Directorate of National Vector Borne Disease

Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of India, Delhi

Print-ISSN: 0976-0245-Electronic-ISSN: 0976-5506, Frequency:

(Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website: www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Dev elopment. The journal does not guarantee directly or indirectly the quality or efcacy of any product or service featured in the advertisement in the journal, which are purely commercial

Editor

Dr. R.K. Sharma

Institute of Medico-legal Publications 501, Manisha Building, 75-76, Nehru Place, New Delhi-110019

Printed, published and owned by

Dr. R.K. Sharma

Institute of Medico-legal Publications 501, Manisha Building, 75-76, Nehru Place, New Delhi-110019

Published at

Institute of Medico-legal Publications

501, Manisha Building, 75-76, Nehru Place, New Delhi-110019



Swapna KG, Valsamma Joseph

Indian Journal of Public Health Research & Development

www.ijphrd.com

| Vo | olume 9, Number 8 August 2018 |
|----|--|
| 1. | Knowledge, Attitudes, and Practices of Pesticides Use Toward the Cholinesterase Enzyme in Vegetable Farmers Kusrini Wulandari, Wakhyono Budianto, Kuat Prabowo, Asep Tata Gunawan, Ismi Rajiani |
| 2. | A Cross-Sectional Study to Assess the Quality of Sleep and Level of Stress among First Year Medical Students in a Private Medical College in Chennai, Tamil Nadu |
| | Profile, Knowledge and Problems of Grass Roots Level Workers and Service Delivery by them in ICDS Blocks: A Cross Sectional Study |
| 4. | A Comparative Study on Normal Symptomatic Medication with Andrographis Paniculata (Nila-Vembu) Extract on Platelets Count in Thrombocytopenia Patient affected from Dengue Fever |
| | Evaluation of the Rationality of Claims Made in Drug Promotional Literature using World Health Organization Guidelines |
| | An Assessment of Water, Sanitation and Hygiene Practices in an Urban Slum of Visakhapatnam, Andhra Pradesh |
| | Comparative Prevalence and Pattern Analysis of Liver Abscesses in Alcoholic and Non-Alcoholic Patients 31 Anoop Chhabra, Bhuvnesh Yadav, Anil Kumar |
| | Effectiveness of House of Recovery Stunting (Hrs) to Reducing Stunting Child Below Two Years Old at Streams Area |
| 9. | Study to Assess the Effectiveness of Self Instructional Module on Neonatal Rearing Practices |
| | Incidence of Gestational Hypertension among Pregnant Women in the Rural Population of District Amritsar- A Community based Study |
| | Kuldip Raj, Aditya Paul, Kritika Bansal, Priyanka Devgun, Utkarsh Passi Comparison of the Psychosocial well-being of HIV Infected and HIV affected Adolescents Availing Services of State AIDS Control Society (SACS) in Kerala |

| 12. | Implementation of Family Planning Program Policy based on Culture in Jayawijaya Province of Papua54 Saparuddin Latu, Indar, Alimin Maidin, Darmawansyah, Sukri Palutturi |
|-----|--|
| 13. | Prevalence of Prediabetes among Traditional Population, Jenukuruba Tribal Children in Mysore District 59 <i>HS Rajani, D Narayanappa, K Jagadish Kumar, VG Manjunath</i> |
| 14. | Burnout Syndrome and Associated Factors among Dental Students |
| | Status of Public Health Care Delivery System– A Case Study of Nagaon and Nalbari Districts of Assam (India) |
| 16. | A Structural Equation Model of Tuberculosis (TB) Infection in Children |
| | A Correlation of National Health Insurance Self-Enrolled Member's Satisfaction on Healthcare Services with Premium Payment |
| | Study of Pleural Fluid to Serum Cholinesterase Ratio to Differentiate between Transudative and Exudative Pleural Effusions |
| 19. | A Study of Factors affecting Utilization of OT |
| | A Cross-Sectional Study of Prevalence of Hypertension among Women of Reproductive Age in Rural Community of Amritsar – A Community based Study |
| | Perception and Feedback of Final Year Medical Students on Teaching-Learning Methods in Andhra Medical College, Visakhapatnam |
| 22. | Social Correlates of Tobacco Consumption among the Medical Students in Meerut |
| 23. | Neuroretinal Rim Criteria Support an Ischemic Theory for Glaucomatous Optic Neuropathy112 Zainab Adel Hashim |
| 24. | Awareness of Breast Cancer and Breast Self-Examination among Female Non-Medical University Students |
| 25. | Knowledge and Awareness of Basic Life Support among Interns of a Dental College in Mangalore, India 124 <i>Karthik Shetty, Roma M, Mranali Shetty</i> |
| | Financial Feasibility of Selected Cardiology Surgeries Under Community Based Health Insurance Schemes in a Tertiary Care Hospital in South India |
| 27. | Effectiveness of an Organized Play Session among Hospitalized Children and Parents' Attitude |

| | Study of Bed Allocation and Utilisation in a Tertiary Care Teaching Hospital in South India |
|-------|---|
| | Evaluating Accuracy of Shade Matching in a Cross-Section of Dentists in a Dental Institute, Karnataka, South |
| | India |
| | Assessment of Competency based Medical Internship Training with 'Cumulative Grade Points Average System'- An Innovative Step Towards Meeting 'Vision 2015' of Medical Council of India |
| | Knowledge and Support of Family Members in the Management of Diabetes Mellitus among Diabetic Adults |
| | Athul Jayan, Neha Patil, Pray <mark>een, Priyal Jo</mark> vit <mark>a Noronha, Wilma Sandra D'</mark> Souza, Malathi G Nayak, Vinish <mark>V</mark> |
| | A Cross Sectional Study to Assess Relationship between Parental Locus of Control and Caries Experience in Preschool Children of Belgaum City |
| | Awareness Regarding Immunization among Mothers of Children from Northern Kerala Attending Pediatric Units, AIMS Kochi |
| | A Pre-screening Method for Cardiac Anomalies Detection using Phonocardiography Signals |
| | Explore the Feeding Practices on Health Outcome of Infants Among Mothers in Pediatric Wards at AIMS, Kochi |
| | Diosmetin Retrieves the Hippocampal Neuropathology on Alzheimer's-Like Ethanol-Induced Cognitive Impairment Rat Models |
| | Effectiveness of an Intervention Module in Improving the Knowledge, Attitude and Practices Regarding Diabetic Retinopathy in Diabetic Individuals in a Tertiary Care Teaching Hospital in Coastal Karnataka198 Tejaswi Pagadala, Kavitha T.C, Karan Giriyan, Keerthi Haripriya, Panchajanya Paul, Ajay Pal Singh, Rajesh Kamath |
| 38.] | Level of Activity and Obesity among High School Children |
| | Nitric Oxide Levels in Chronic Renal Failure Patients and Maintenance Hemodialysis in Comparison to Healthy Controls |
| | Salma Mahaboob R, E Prabhakar Reddy |
| 40. | Self-Esteem and Adolescents' Perception Regarding Maternal Parenting Style |

| | Lean Manufacturing Process Establishment for Lead Time Reduction in Pump Manufacturing Industry Us for Health Care Applications | |
|-----|--|-------|
| | J Logeshwaran, RM Nachiappan | 221 |
| | Evaluation of Teachers' Knowledge about the Early Detection for Mumps Diseases at the Primary School Al-Hilla City | |
| | Amean Ajeel Yasir, Ali F Abdul Hussein | |
| | Analysis Factor Influence with Waiting Time for Elective Surgery in General Surgical Outpatient Clinic Akhyarudin Noor, Izaak Zoelkarnain Akbar, Husaini, Roselina Panghiyangani, Lenie Marlinae | 233 |
| 44. | Occupational Stress of Nurses Working in the Academic-Related Sectors | . 238 |
| | The Effect of Service Quality and Operational Benevolence on Patient Satisfaction and Their Interest in Relation | |
| | Devy Halim, Agustinus Joh <mark>anes Djoha</mark> n <mark>, Roselina Panghiyan</mark> gani <mark>, Husaini, Lenie</mark> Marlinae | |
| | Message-Based Interaction (MBI) Mobile Phone Platform to Expedite Counselling and Education to HIV Patients in Developing Countries | 249 |
| 47. | Comparison of the Effects of Chlorhexidine Mouthwash with Jaftex in Periodontal Index | .255 |
| 48. | Identification of Health Service Requirements in HIV/AIDS Patients: A Phenomenological Study | . 260 |
| | Diagnosis of HCV Infection in Renal Chronic Infection Patients by Using ELSA and RT-PCR in Tikrit City | 265 |
| 50. | Impact of Dietitians in Errors Associated to Parenteral Nutrition Prescription | 271 |
| 51. | Critical Care Nurse's Capacities to Express a Caring Relationship with Patients in Teaching Hospitals | 277 |
| | A Strategic Framework for Effective Utilisation of eHealth Tools by Medical Doctors in Zimbabwe's Pub Hospitals | |
| | Self-Esteem, Sexual Values and Sexual Risk Behavior among Undergraduate Students in the Northern | |
| | Thailand | .289 |
| 54. | Mothers' Beliefs Toward Care of Neonatal Jaundice in Pediatric Teaching Hospital in Karbala City | 295 |
| | The Analysis of Behavioral Risk Factors on Periodontal Diseases in the Hospitalized Public Health Centre Cempaka Region Banjarbaru | |

| | The Effect of Supplementary Feeding Program for Chronic Energy Deficiency Pregnant Women on Hb Concentration, Muac, and Gestational Weight Gain in Indonesia | .306 |
|-----|---|------|
| | Henrick Sampeangin, Veni Hadju, Saifuddin Sirajuddin, Andi Imam Arundhana Thahir, Abdul Razak Thaha | |
| 57. | The Compliance of Fall Prevention Protocol through Patients with High Risk of Fall | .313 |
| | Effectiveness of an Educational Program on Nurses' Knowledge Concerning Nursing Management for Pat with Compound Fracture at Orthopedic Wards in Medical City Directorate | |
| | The Impact of Compensation, Career Development, and Organizational Culture Towards Nurses Commitmat Suaka Insan Hospital Banjarmasin | |
| | Evaluation of Teachers' Knowledge about the Early Detection for Chickenpox Diseases at the Primary Schin Al-Hilla City | |
| 61. | Influencing Factors of Marital Satisfaction in the Married Migrant Women from China | 337 |
| | Planning an Energy Management Program based on Hospital Electricity Consumption in a Tertiary Care Hospital in South India Shruti Billore, Kavitha T C, Namesh Malarout, Ashalata Pati, Aswathi Raj L, Biju Soman, Rajesh Kamath | .344 |
| 63. | Intention to Stay Model of Nurse Staff in Hospital | .351 |
| | Socio Demographic Profile in Relation to Care Givers of Persons with Permanent Disabilities Attending Rehabilitation Centre in H.D. Kote Taluk, Mysore District | .356 |
| | Comparative Evaluation of Aluminum Foil and Air/Water Spray as Temperature Control Method Using Direct Provisionalization | |
| 66. | Analysis of Environmental Health Risks Resulting from Exposure to Pollution Materials | .368 |
| | Comparison of Quality of Impression and Clinical Performance after Gingival Displacement with Magic F Cord and Gingival Retraction Cord - An in vivo Study | |

Analysis of Environmental Health Risks Resulting from Exposure to Pollution Materials

Isa Ma'rufi¹, Eri Witcahyo², Heru Santoso Wahito Nugroho³

¹Faculty of Public Health, University of Jember, ²Faculty of Public Health, University of Jember, ³Health Polytechnic of Ministry of Health at Surabaya

ABSTRACT

Analysis of Environmental Health Risks is an approach to calculate or predict the risks to human health, including factors identify of uncertainty, tracking on specific exposures, consider the inherent characteristics to the concern agent and characteristics of the specific target. The research was descriptive study by risk agent which were SO2, H2S, NO2, PM2.5 and PM10. The research was conducted in five zones Purwodadi sugar factory area at Magetan. Variables in the study were the identification of hazards, exposure analysis, the level of risk (RQ), and environmental health risk management. Data analysis technique used was descriptive. The results of research showed that the highest levels of risk agent SO2 levels was 0.33 mg/ m3, the highest levels of NH3 is 2.20 mg/ m3, highest levels of NO2 is 0.39 mg/ m3, highest levels of PM2.5 is 2.17 mg/m3 and PM10 at 1.76 mg/ m3. The distance value of the level of risk (RQ) for SO2, NH3 and NO2 was at ± 500 m from the source of the pollutants and showed above 1 mg/m3, it means the gas of NH3, PM2.5 and PM10 are highly risk to the health.

Keywords: Environmental health risks, Risk level, Sugar factory, Risk agent

INTRODUCTION

Analysis of Environmental Health Risk (ARKL) is a risk management tool that is used to protect the health of the community due to the effects of a bad environment. In Indonesia regualtions, ARKL is also a means of analysis of environmental health impact (ADKL) approach. Legal bases ARKL for ADKL are PerMenLH No 08/2006 on guidelines for the preparation of the Analysis of Environmental Impacts (AMDAL)(1), and KepMenKes No 876/Menkes/SK/VII/2001 on technical guidelines for ADKL(2).

ARKL that is used as ADKL approach is a great tool to get to know, understand, and predict the conditions and characteristics of the environment that could potentially pose a health risk as the basis for compiling or developing management and monitoring

Corresponding author: Heru Santoso Wahito Nugroho

E-mail: heruswn@gmail.com

Health Polytechnic of Ministry of Health at Surabaya

Jl.Pucang Jajar Tengah 56 Surabaya, Indonesia

environmental health risks. ARKL also constitute an adequate method to do a study of the health effects of the pollution cases in general(3).

According to Louvar & Louvar, ARKL a scientific framework for the review of the Environment and Health to solve the problems(4). US-EPA defines ARKL is the Scientific Evaluation of the potential health impacts Happens Certain substances exposure can because of or mixture specific conditions(5). While PPCs defines ARKL is the process of predicting the risk on a biota, system or (sub)population objectives, with all non uncertainty the accompanying, taxable income exposure posted agents specific, with notice characteristics of agents and objectives, specific (6).

Air is a mixture of gases, which consists of approximately 78% Nitrogen, 20% Oxygen, 0.93% Argon, 0.03% carbon dioxide and the rest consists of Neon, Helium, methane and hydrogen. The air is said to be "normal" and able to support human life in a composition like above. Whereas in case of adding other gases cause interference as well as changes in the composition, then said the air already polluted/

contaminated(7).

Purwodadi PG (sugar factory) in Karangrejo, Magetan as one sugar factory relics of the Dutch era, according to local residents the surrounding air polluted with particulate is emitted from chimney. It has been many years since the 1960s, and took place at a time when sugar factory ground, more or less during 6 months between May until October. Preliminary observations indicate that air pollution around Purwodadi PG felt already interfere with the health of the community, some citizens complained against their respiratory disorders, in addition to the presence of air pollution by the discovery of soot (former burnt leaves) above the ceilings of houses.

See the conditions then the need for environmental health impact analysis in the vicinity of Purwodadi PG to see the extent to which environmental quality resulting from pollution by emissions and liquid waste health effect on PG surrounding residents. This facts are needs to be done in terms of monitoring the air, liquid and solid wastes resulting from Purwodadi PG.

MATERIALS AND METHOD

Type of this research was descriptive, with risk agents were SO₂, NH₃, NO₂ PM_{2,5} and PM₁₀. Environmental health risk analysis method used was the wearing method approach to the international programme on chemical safety (IPCS) and the WHO which has been developed by Health Ministry and the University of Indonesia to be applied extensively in Indonesia. The research was carried out around the region of Purwodadi PG between ±500 m - 4000 m to see the environmental health risks due to air pollution motor vehicles. Variable in this research was the identification of hazards, identification of sources, analysis of exposure, dose-response analysis, risk characterization, risk management health environment. The characteristics of the risk represented by the level of risk (Risk Quotient) are a division between the intake (I) and inhalation reference concentration (RfC). In addition to determining the inhalation intake also needed Anthropometry parameters (weight and inhalation rate), the pattern of activity (time, frequency and duration of exposure) and so on. The level of risk is calculated by equation 1 and inhalation intake (I) calculated using equation 2(4).

Description:

I= Inhalation Intake (mg risk agent/kg west of individual agencies/day)

C= Risk Consentration agent in the air (mg risk agent/m3 air)

R= Inhalation rate (m3 air/hour)

tE = long exposure (hour/day)

fE = exposure freequension, 350 days/year for residensial default value

Dt= exposure duration, 30 year to default for resdensial population

Wb= west of individual agencies(kg)

tavg= The average time periode (Dt 365 days/years for noncarsinogenic, 70 years, 365 days/years for carsinogenic)

Environmental health risk management was carried out by four main elements, namely the evaluation of risks, control of emissions and exposures and monitoring of risks. Risk management was calculated by using the formula:

Maximum Concentration of safe human consumption (C.max).

Maximum consumption Rate that is safe for humans (R.max)

The assumptions used are RQ = 1

The calculation of the concentration of safe and secure consumption rate was:

Decision-making methods were:

If the initial concentration >max.secure concentration, it is necessary to take steps to reduce the concentrations of, improve environmental conditions, etc.

If the initial intake rate >max.secure intake rate, then the necessary measures to bring down the rate of intake by reducing dwell time / time said, etc.

RESULTS AND DISCUSSION

Sources of SO₂, NH₃, NO₂, PM_{2.5} and PM₁₀

Pollution source of SO₂, NH₃, NO₂, PM_{2.5} and PM₁₀ most (70%) derived from the production process Purwodadi PG and fraction (30%) comes from transportation the transport of raw sugar cane. Purwaningsih reported that the air quality at present milled simply applied a two-fold compared to when not milled, it is shown that most of the pollution comes from the production process(8).

Hazard Identification

Table 1. The Concentration of Gases and Particles around Purwodadi PG

| 77 | A of T of | Concentration (in mg/m ₃) | | | | | |
|------|-------------------------------------|---------------------------------------|-----------------|-----------------|------|------------|--|
| Zone | Area/ Location | SO ₂ | NH ₃ | NO ₂ | 2.5 | PIVI 10 | |
| 1 | ±500 m from the source of pollutan | 0.33 | 2.20 | 0.39 | 2.17 | 1.76 | |
| 2 | ±1000 m from the source of pollutan | 0.26 | 1.73 | 0.35 | 1.52 | 1.13 | |
| 3 | ±2000 m from the source of pollutan | 0.23 | 1.21 | 0.30 | 1.06 | 0.77 | |
| 4 | ±3000 m from the source of pollutan | 0.18 | 1.13 | 0.24 | 1.01 | 0.76 | |
| 5 | ±4000 m from the source of pollutan | 0.13 | 0.02 | 0.21 | 0.98 | 0.75 | |
| | Environmental Quality Standards | 0.26 | 1.36 | 0.92 | 0.26 | 1.00 | |

Based on the raw quality of environment against ambient air concentration measurements around the sugar factory, indicating that all of the ambient air at a distance of approximately 500 m and 1000 m from the sugar factory on top of BML, which means that the ambient air at a distance of 1 km of sugar factories are not healthy or polluted conditions. BML is the size limits or levels of living things, energy, substances, or components in the environment. BML is an instrument for managing the environment, used to find out the level of or decrease in intensity of reduction environment(9).

Exposure Analysis

Table 2. Default value for antropometric parametes

| Receptor | Inhalation rate (m ₃ /hour) | Exposure Frequency (hour) | Weight (kg) | Exposure Duration (year) | Frequency (day/year) |
|-------------|---|---|-------------|--------------------------|-------------------------|
| Children | 0.5a | 18 (6hour studying in the outside location) _c | 15a | 2° | 350a |
| Adult (IRT) | 0.83a | 24c | 55ь | 2c | 350a |
| Worker | 0.83a | 14 (10 hour working in the outside location) _c | 70a | 2c | 350a |

Source: aU.S.EPA 1990(5), bNukman et al (2005)(10), cauthor assumption

Table 3. Intake Risk Agent around Purwodadi PG

| | | Intake Ri | Intake Risk Agent | | | | | |
|------|--|-----------|-------------------|-----------------|--------|-----------|--|--|
| Zone | | SO_2 | NH ₃ | NO ₂ | PM 2.5 | PM_{10} | | |
| 1 | ±500 m from the source of pollutan | 0.0048 | 0.0318 | 0.006 | 0.0314 | 0.0255 | | |
| 2 | ± 1000 m from the source of pollutan | 0.0038 | 0.025 | 0.005 | 0.022 | 0.0164 | | |
| 3 | ±2000 m from the source of pollutan | 0.0033 | 0.0175 | 0.004 | 0.0153 | 0.0111 | | |
| 4 | ±3000 m from the source of pollutan | 0.0026 | 0.0164 | 0.003 | 0.0146 | 0.011 | | |
| 5 | ±4000 m from the source of pollutan | 0.0019 | 0.0078 | 0.003 | 0.0142 | 0.0109 | | |

Dose-Response Analysis

Table 4. Value of Dose-Response

| No. | Agent | RfC | Critical Effect and Sources of the Date (Referention) |
|-----|------------------|--------|---|
| 1. | SO ₂ | 2,6E-2 | Respiratory Disorders(3),(5),(10) |
| 2. | NO ₂ | 2E-2 | Respiratory Disorders(5),(11) |
| 3. | H ₂ S | 2E-3 | Olfactory Mocus Nasal lesion on rat inhalation bioassay subcronic (Brenneman, James, Gross, & Dorman, 2000) in IRIS(12) |
| 5. | TSP | 2,42 | Respiratory Disorders (3),(5),(10) |

Dose-response assessment was conducted to determine the toxicity of the quantitative values a brisk agent for any form of chemical species, expressed as a RfD (for drinking water and food) or RfC (for air) for non carcinogenic effects and the Slope Factor (SF) or Unit Risk (UR) to the effects of carcinogens (12).

on each segment of the population with the default values for inhalation rate, weight, exposure duration and frequency. As for the mean time period – average (t_{avg}) is 6 years (365 days/year) for the subpopulations of children and 30 years (365 days/years) to subpopulations of adults.

Risk Characteristic (Risk Level)

The characteristics of the risk is calculated based

Table 5. Risk Level (RQ) SO₂, NO₂, H₂S, and TSP for adult (IRT) with W_b 15 kg, 55 kg and 70 kg and f_E 350 days/years

| Zone | · · · · · | Adult | | 71 | | |
|------|--------------------------------------|-----------------|-----------------|-----------------|--------|------------------|
| Zone | Location | SO ₂ | NH ₃ | NO ₂ | 2.5 | PM ₁₀ |
| 1 | ±500 m from the source of pollutant | 0.1837 | 1.0612 | 0.282 | 1.3084 | 1.0612 |
| 2 | ±1000 m from the source of pollutant | 0.1447 | 0.8345 | 0.253 | 0.9165 | 0.6813 |
| 3 | ±2000 m from the source of pollutant | 0.128 | 0.5837 | 0.217 | 0.6391 | 0.4643 |
| 4 | ±3000 m from the source of pollutant | 0.1002 | 0.5451 | 0.174 | 0.609 | 0.4582 |
| 5 | ±4000 m from the source of pollutant | 0.0724 | 0.2605 | 0.152 | 0.5909 | 0.4522 |

The overall value of RQ for each risk agents in the location of the study population, there was some segmentation according to the indicate above 1 (RQ>1) and there are some which show brought 1 (RQ<1). It means some of the gas in the air that come from pollution emissions chimney Purwodadi PG takes control risk quotient for SO2 all under 1 (RQ<1), that is in the range 0.0724-0.1837. RQ NH₃ at a distance of approximately 500 m from the source of pollutants is more than 1 (>1), medium distance >500 m still under 1 (<1) RQ NO₂, SO₂ under 1 (RQ<1), RQ PM_{2.5} in a distance of approximately 500 m from the source of pollutants is more than 1 (>1), medium distance >500 m still under 1 (<1), and RQ PM₁₀ at a distance of approximately 500 m from the source of

pollutants is more than 1 (>1), medium distance >500 m still under 1 (<1).

Risk agent according to the it risk quotient the most dominant is NH $_3$, PM $_{2.5}$ and PM $_{10}$, so that it can be concluded that the people who live in an area with a diameter of Purwodadi PG \pm 500 m from the source of the pollutants risk of experiencing health problems. Aditama mention that the gases of SO $_2$, NH $_3$, NO $_2$, PM $_{2.5}$ and PM $_{10}$ can cause abnormalities in the respiratory tract if the pollutants in the ambient air of breathe. Gas that the most harmful to the lungs are SO $_2$ and NO $_2$, if this item smoked, then the various complaints in the lungs will arise with the name CNSRD (a non specific chronic

372

respiratory disease) such as asthma and bronchitis(13).

Research held by Groneberg Kloft et al. showed that SO_2 concentrations are low though continuously can irritate the respiratory tract and cause chronic cough or bronchitis (14).

NH₃ exposure in the low levels can cause pulmonary disorders if the exposure takes place in a long time, it is in accordance with the studies by Heederick et al., namely in the form of a cohort study on farmers exposed to ammonia 1.60 mg/m3(15). Cross sectional studies performed by Balla et al. show that male workers showed a relationship between exposure to ammonia gas with respiratory disorders such as asthma symptoms bronchial where the levels of ammonia in the interval 2.82-183.86 ppm(16).

Purwaningsih reported that respiratory disorders, such as cough, shortness of breath, fever, lethargy and perceived by respondents who live in the region of the dominant wind direction when the factory operates (milled) and at the end of a rolling pin. Respondents who live in the wind direction is not the dominant States that coughs, colds and Croup which is perceived is not affected by the pollutants Mojo PG. When associated with ambient air quality measurement results then it can be inferred that the pollutants Mojo PG respiratory tract disorders(8).

Risk Management

Various possibilities for risk management a risk agent needs to do. The first thing that is done is by means of lowering the value of concentration of each risk agent. For it takes a quantity value risk agent is needed so that the frequency of RQ=1 or RQ<1.

According to the information above, is not possible if it should reduce the exposure time of the daily and annual exposure because it is very small and is not realistic.

Altough the various risk management has been attempted and be scenario to cope with exposure risk agent to the population, but have not found a solution. Therefore, the last attempt is to evacuate in advance of the study on the location of residents to safety and avoid pollutants. If that intervention doesn't held, the possibilities of the population to accept

the health risks resulting from exposures were higher.

CONCLUSION

 SO_2 levels agent highest was 0.33 mg/m₃, the highest levels of NH $_3$ is 2.20, NO $_2$ levels highest 0.39, the highest levels of PM_{2.5} is 2.17 and the highest level of PM₁₀ is 1.76. The value of the level of risk (RQ) for SO_2 , NH₃, and NO₂ at a distance of approximately 500 m from the source of the pollutants shows over 1 (one), it means the gases NH₃, PM_{2.5} and PM₁₀ are very risky against health.

The most urgent control is recommended for controlling the environment, such as the blue sky program, encouraged the planting of plants forbidding residents to live in the territory and population of the sugar factory could be moved to a safer place.

Conflict-of-Interest: None

Source of Funding: Authors

Ethical Clearance: Taken from University of Jember.

REFERENCES

KemenKL. Regulation of the Minister of Environment No. 08, 2006 on Guidelines for the Preparation of Environmental Impact Analysis (Peraturan Menteri Negara Lingkungan Hidup Nomor 08 Tahun 2006 Tentang Pedoman Penyusunan Analisis Mengenai Dampak Lingkungan Hidup). Jakarta: KemenKL; 2006.

Kemenkes. Minister of Health Decree No. 876 / Menkes / SK / VIII / 2001 About ADKL Technical Guidelines (Keputusan Menteri Kesehatan No. 876 / Menkes/SK/VIII/2001 Tentang Pedoman Teknis ADKL). Jakarta: Kemenkes RI; 2001.

Rahman A. Environmental Health Risk Analysis (Study of Public Health Aspects in EIA Studies and Environmental Pollution Cases) (Analisis Risiko Kesehatan Lingkungan (Kajian Aspek Kesehatan Masyarakat dalam studi AMDAL dan Kasus-Kasus Pencemaran Lingkungan)). Depok: Pusat Kajian Kesehatan Lingkungan dan Industri, FKMUI; 2007.

Louvar JF, Louvar BD. Health and Environmental Risk Analysis: Fundamentals with Application. New Jersey: Prentice Hall; 1998.

US-EPA. Exposure Factors Handbook, EPA 600/8-

89/043. Cincinnati: US Environmental Protection Agency; 1990.

IPCS. Environmental Health Criteria XXXX: Principles for modeling dose-response for the risk assessment of chemicals (draft). Genewa: WHO and International Programme on Chemical Safety; 2004. Kastiyowati I. Impacts and Efforts to Address Air Pollution (Dampak dan Upaya Penanggulangan Pencemaran Udara). Jakarta: Puslitbangtek Balitbang Dephan; 2003.

Purwaningsih ED. Air Pollution and Its Effect on Health (Polusi Udara dan Pengaruhnya terhadap Kesehatan). Graduate Thesis. UGM: Yogyakarta; 2012.

KepmenKL. Assessment of Ambient Air Quality Standards (Pengkajian Baku Mutu Lingkungan Udara Ambien). Jakarta: KemenKL; 2011.

Nukman A, Rahman A, Warouw S, Ichsan M, Setiadi, Akib CR. Analysis and Management of Health Risks from Air Pollution: A Case Study in the Nine Big Cities with Solid Transportation (Analisis dan Manajemen Risiko Kesehatan pencemaran Udara: Studi Kasus di Sembilan Kota Besar Padat Transportasi). J Ekolog Kesehatan. 2005;4(2):270-289.

Kolluru RV. Health Risk Assessment: Principles and Practices. In: Risk Assessment and Management Handbook for Environmental, Health, and Safety Professionals (Kolluru RV, Bartell s, Pitblado R, Stricoff S, sds). New York: McGraw-Hill; 1996. IRIS. Integrated Risk Information System List of Subtances. 2012.

Aditama TY. Air Pollution and Health (Polusi Udara dan Kesehatan). Jakarta: Arcan; 1992.

Groneberg KB, Kraus T, Mark VA, Wagner U, Fischer A. Analyzing The Causes of Chronic Cough: Relation to Diesel Exhaust, Ozone, Nitrogen Oxides, Sulphur Oxides and Other Environmental Factors. Journal of Occupational Medicine and Toxicology. 2006.

Heederik D, Vogelzang p, Gulden J, Folgering H, Tielen M, Schayck V. Longitudinal Changes in Bronchial Responsiveness Associated with Swine Confinement Dust Exposure. Chest journal. 2000;117:1488-1495.

Balla, SG, Ali BA, Albar AA, Ahmed AO, Hasan AY. Bronchial Asthma in Two Chemical Fertilizer Producing Factories in Eastern Saudi Arabia. Int Journal Tuberc Lung Dis. 1998;2: 330-335.

Call for Papers/Article Submission

The editor invites scholarly articles that contribute to the development and understanding of all aspects of Public Health and all medical specialities. All manuscripts are double blind peer reviewed. If there is a requirement, medical statistician review statistical content. Invitation to submit paper: A general invitation is extended to authors to submit papers papers for publication in IJPHRD.

The following guidelines should be noted:

The article must be submitted by e-mail only. Hard copy not needed. Send article as attachment in e-mail.

The article should be accompanied by a declaration from all authors that it is an original work and has not been sent to any other journal for publication.

As a policy matter, journal encourages articles regarding new concepts and new information.

Article should have a Title

Names of authors

Your Affiliation (designations with college address)

Abstract

Key words

Introduction or back ground

Material and Methods

Findings

Conclusion

Interest of conflict

References in Vancouver style.

Please quote references in text by superscripting

Word limit 2500-3000 words, MSWORD Format, single file

All articles should be sent to : editor.ijphrd@gmail.com

Our Contact Info: Institute of Medico-Legal Publications

501, Manisha Building, 75-76, Nehru Place, New Delhi-110019, Mob: 09971888542, E-mail: editor.ijphrd@gmail.com Website: www.ijphrd.com



Indian Journal of Public Health Research & Development

CALL FOR SUBSCRIPTIONS

About the Journal

Print-ISSN: 0976-0245 Electronic - ISSN: 0976-5506, Frequency: Monthly

Indian Journal of Public Health Research & Development is a double blind peer reviewed international Journal. The frequency is half yearly. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, Public Health Laws and covers all medical specialities concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent, and south east Asia.

The journal has been assigned international standards (ISSN) serial number and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases.

Subscription Information

| Journal Title | N NV | Pricing of Journals | |
|---------------|------------|---------------------|-------------|
| IJPHRD | Print Only | Print+Online | Online Only |
| Indian | INR 7000 | INR 9000 | INR 5500 |
| Foreign | USD 450 | USD 550 | USD 350 |

Note for Subscribers

Advance payment required by cheque/demand draft in the name of "Institute of Medico-Legal Publications payable at New Delhi.

Cancellation not allowed except for duplicate payment.

Claim must be made within six months from issue date.

A free copy can be forwarded on request.

Send all payment to : Institute of Medico-Legal Publications

501, Manisha Building, 75-76, Nehru Place, New Delhi-110019, Mob: 09971888542, E-mail: editor.ijphrd@gmail.com, Website: www.ijphrd.com

