# Journal of

International

**Dental and Medical** 

## Research



2022 - Vol. 15 – No. 1 http://www.jidmr.com















Pages 392-402	
65. <b>Bibliometric Analysis of the 1</b> Nanda Rachmad Putra Gofur, Achmad Zam Zan	00 Most Cited Articles on Oral Biology Adhasy, Zamros Yuzadi Bin Mohd Yusof
Pages 403-411	
66.Craniofacial and upper Airwa Systematic Review	REVIEW y Morphology in Pediatric Obstructive Sleep Apnea Patients: A
Pages 412-421	nakom Saengphen, inn Sinsooniom
67.0zonotherapy as a Method of Diseases- Literature Review Michalina Zylkiewicz [Sd], Sylwia Kuderewska, C Full Text PDF	REVIEW Alleviating Symptoms of Oral Mucositis and other Selected Oral Grazyna Marczuk-Kolada
<b>MEDICINE</b> 68.Survival of Methicillin-Resist	EXPERIMENTAL ARTICLE ant Coagulase-Negative Staphylococci from Floor Dust
Plykaeow Chaibenjawong, Kannipa Tasanapak, Full Text PDF	Sutthirat Sitthisak, Thaniya Wongwan, Suttipalin Suwannakul
69. The Effects of Mobile Phones Hava Bektas Full Text PDF	EXPERIMENTAL ARTICLE on Diabetes and Appetite
70.Mother's Eating Behavior Dur Prevention in Madura, Indonesia Esti Yunitasari, Ribka Putri Sholecha, Ni Ketut A Full Text PDF	CLINICAL ARTICLE ring Pregnancy and Family Income with Malnutrition: Stunting (Mother's Eating and family Income with Stunting Prevention) It Armini, Retnayu Pradanie, Wahyuni Tri Lestari, Bih-O Lee
71.Functional Resources and Psy Northern University Artyom Borisovich Kiryanov, Mikhail Alekseevich	CLINICAL ARTICLE vchological Comfort of Students from Central Asia Studying at the Kunavin, Anna Andreevna Farkova
Pages 454-460 Editoria	Board of JIDMR

Prof. Dr. Izzet YAVUZ Editor-in-Chief and General Director

### **Advisory Board**

Prof. Dr. Refik ULKU Editor for Medicine Prof. Dr. Zulkuf AKDAG Editor for Biomedical Research Prof. Dr. Ozkan ADIGUZEL Associate Editor

Gajanan Kiran KULKARNI (CANADA) Betul KARGUL (TURKEY) Diah Ayu MAHARANI (INDONESIA) Francisco Cammarata-Scalisi (Venezuela) Myroslav Goncharuk-Khomyn (UKRAINE)

Ferranti WONG (UK) Zeki AKKUS (TURKEY) Michele CALLEA (ROME, ITALY) Zelal ULKU (TURKEY) Moschos A. PAPADOPOULOS ( Lindawati S KUSDHANY (INDO Yasemin YAVUZ (TURKEY) Yuliya MAKEDONOVA (RUSSIA Noriah Nik HUSSEIN (MALAYSI

## **Editorial Board**

Abdel Fattah BADAWI (EGYPT) Abdurrahman ONEN (TURKEY) Ahmet YALINKAYA (TURKEY) Ahmet DAG (TURKEY) Ali Al-ZAAG (IRAQ) Ali BUMIN (TURKEY) Ali FADEL (EGYPT) Ali GUR (TURKEY) Ali Kemal KADIROGLU (TURKEY) Ali Riza ALPOZ (TURKEY) Ali Riza Tunçdemir (TURKEY) Allah Bakhsh HAAFIZ (USA) Alpaslan TUZCU (TURKEY) Alpen ORTUG (TURKEY) Armelia Sari WIDYARMÁN (INDONESIA) Ashish AGGARWAL (INDIA) Ayse GUNAY (TURKEY) Aziz YASAN (TURKEY) Awiruth KLAISIRI (THAILAND) Balasubramanian MADHAN (INDIA) Benik HARUTUNYAN (ARMENIA) Betul KARGUL (TURKEY) Betul URREHMAN (UAE) Bugra OZEN (TURKEY) Carlos Menezes AGUIAR (BRAZIL) Cemil SERT (TURKEY) Chiramana SANDEEP (INDIA) Christine Bettina STAUDT (SWITZERLAND) Cihan AKGUL (TURKEY) Claudia DELLAVIA ( ITALY ) Diah Ayu MAHARANI (INDÓNESIA) **Dinesh Rokaya (NEPAL)** Edoardo BAUNER (ROMA) Emmanuel Joao N. Leal da SILVA (BRAZIL) Emin Caner TUMEN (TURKEY) Emrullah BAHSI (TURKEY) Ertunc Dayı (TURKEY) Fadel M. ALI (EGYPT) Fahinur ERTUGRUL (TURKEY) Feral OZTURK (TURKEY) Feridun BASAK (TURKEÝ) Ferranti WONG (UNITED KINGDOM) Feyzi Çelik (TURKEY) Feyzullah Uçmak (TURKEY) Figen SEYMEN (TURKEY) Filippo BATTELLI (ITALY) Filiz Acun KAYA (TURKEY) Flavio Domingues Das NEVES (BRAZIL) Folakemi OREDUGBA (NIGERIA) Francesca De Angelis (ITALY) Gajanan Kiran KULKARNI (CANADA) Gamze AREN (TURKEY) Gauri LELE (INDIA) Gonul OLMEZ (TURKEY) Gulsen YILMAZ (TURKEY) Gulten UNLU (TÜRKEY)

Guvenc BASARAN (TURKEY) Guven ERBIL (TURKEY) Halimah AWANG (MALAYSIA) Halit AKBAS (TURKEY) Heloisa Fonseca MARÁO (BRAZIL) Hilal TURKER (TURKEY) Huseyin ASLAN (TURKEY) Igor BELYAEV (SWEDEN) Ilhan INCI (ZURICH) Ilker ETIKAN (TURKEY) Isil TEKMEN (TURKEY) Isin ULUKAPI (TURKEÝ) Jalen DEVECIOGLU KAMA (TURKEY) Kemal CIGDEM (TURKEY) Kemal NAS (TURKEY) Kewal KRISHAN (INDIA) King Nigel MARTYN(HONG KONG, CHINA) Kursat ER (TURKEY) Levent ERDINC (TURKEY) Luca TESTARELLI (ROMÉ) Lucianne Cople MAIA (BRAZIL) Luciane Rezende COSTA (BRAZIL) Marri Sai ARCHANA (INDIA) Manoj KUMAR (INDIA) Marcelo Rodrigues AZENHA (BRAZIL) Marcia Cancado FIGUEIREDO (BRAZIL) Marco MONTANARI (ITALY) Margaret TZAPHLIDOU (GREECE) Maria Elisa Oliveira dos SANTOS (BRAZIL) Medi GANIBEGOVIC (BOSNIA and HERZEGOVINA) Mehmet DOGRU (TURKEY) Mehmet Emin ERDAL (TURKEY) Mehmet Sinan DOGAN (TURKEY) Mehmet Ünal (TURKEY) Mehmet Zulkuf AKDAG (TURKEY) Meral ERDİNC (TURKEY) Michele CALLEA (ITALY) Mohamed TREBAK (USA) Mohammad Khursheed Alam (KSA) Mohammad SUHRON (INDONESIA) Mohammed Mustahsen URREHMAN (UAE) Moschos A. PAPADOPOULOS (GREECE) Mostaphazadeh AMROLLAH (IRAN) M.S. Rami REDDY (INDIA) Muhammad FAHIM (INDIA) Mukadder ATMACA (TURKEY) Murat AKKUS (TURKEY) Murat SOKER (TURKEY) Mustafa KELLE (TURKEY) Mustafa ZORTUK (TURKEY) Muzeyyen YILDIRIM (TURKEY) Neval Berrin ARSERIM (TURKEY)

Nezahat AKPOLAT (TURKEY) Nihal HAMAMCI (TURKEY) Nik Noriah Nik HÜSSEIN (MALAYSIA) Nicola Pranno (ROME) Nurten AKDENIZ (TURKEY) Nurten ERDAL (TÜRKEY) Orhan TACAR (TURKEY) Ozant ONCAG (TURKEY) Ozgur UZUN (TURKEY) Ozkan ADIGUZEL (TURKEY) Rafat Ali SIDDIQUI (PAKISTÁN) Refik ULKU (TURKEY) Sabiha Zelal ULKU (TURKEY) Sabri BATUN (TURKEY) Sadullah KAYA (TURKEY) Saul Martins PAIVA (BRAZIL) Sedat AKDENIZ (TURKEY) Seher GUNDUZ ARSLAN (TURKEY) Selahattin ATMACA (TURKEY) Selahattin TEKES (TURKEY) Serdar ERDINE (TURKEY) Serdar ONAT (TURKEY) Sergio Adriane Bezerra DE MOURA (BRAZ Serhan AKMAN (TURKEY) Sertac PEKER (TURKEY) Seyed Amir Danesh Sani (USA) Sevit Burhanedtin ZİNCİRCİOĞLU (TURKE Shailesh LELE (INDIA) Sinerik N. AYRAPETYAN (ARMENIA) Smaragda KAVADIA (GREECE) Sossani SIDIROPOULOU (GREECE) Stefano Di CARLO (ROME) Sunit Kr. JUREL (INDIA) Stephen D. SMITH (USA) Susumu TEREKAWA (JAPAN) Suha TURKASLAN (TURKEY) Suleyman DASDAG (TURKEY) Taskin GURBUZ (TURKEY) Ufuk ALUCLU (TURKEY) Ugur KEKLIKCI (TURKEY) Xiong-Li YANG (CHINA) Vatan KAVAK (TURKEY) Yasar YILDIRIM (TURKEY) Yasemin YAVUZ (TURKEY) Yavuz SANISOGLU (TURKEY) Yu LEI (USA) Yuri LIMANSKI (UKRAINE) Zafer C. CEHRELI (TURKEY) Zeki AKKUS (TURKEY) Zeynep AYTEPE (TURKEY) Zuhal KIRZIOGLU (TURKEY) Zurab KOM

#### The Toothpaste's Containing Coffee bean Skin Demonstrated the Specific Physicochemical Properties, Inhibited the Growth of Streptococcus mutans and Increased the Viability of Neutrophil

I Dewa Ayu Ratna Dewanti<sup>1</sup>, Ristya Widi Endah Yani<sup>2</sup>

1. Department of Biomedical Science, Faculty of Dentistry, Jember University, Jember, Indonesia.

2. Department of Public Dental Health Faculty of Dentistry, Jember University, Jember, Indonesia.

#### Abstract

Coffee bean skin contains amino acids, proteins, flavonoids, catechins equivalent to the seeds and leaves, which is thought to be used as toothpaste. As a basic ingredient of toothpaste, coffee bean skin must have good quality.

This study to analyze the Physicochemical properties (organoleptic, pH, homogeneity, spreadability, viscosity), the inhibition to S. mutans and cell viability of toothpaste's contained Robusta coffee bean skin.

Concentrations of toothpaste were 5%, 10%, 15%. Organoleptic test observed shape, color and odor, pH used a pH meter, viscosity test used vicotester VT 03. Spreadability test by measuring the diameter of toothpaste that was smear on the glass. Inhibition was carried out by measuring the diameter of the barriers against S. mutans. Cell viability used neutrophil cells. The cells were placed in 24-well microtiter plate, treated according to group. Viability was stained with Trypan Blue. Cells were counted under an inverted microscope with magnification 400 times from an average of 4 fields of view. The data obtained were ANOVA followed by LSD test and descriptive. Toothpaste preparation was cream colored, soft texture, homogeneous, smells of coffee, has a pH of 7 in accordance with the pH requirements for toothpaste preparations according to SNI 12-3524-1995, namely 4.5-10.5. Inhibition and cell viability tests showed significant differences, but not significant between 15% concentration with commercial toothpaste (ANOVA and LSD analysis, 0.05<p).

Toothpaste's contain coffee bean skin has physico-chemical properties, inhibited to S. mutans and increase cell viability equivalent to commercial toothpaste.

Experimental article (J Int Dent Med Res 2022; 15(1): 66-71) Keywords: Toothpaste, coffee bean skin, organoleptic, spreadability, pH, cell viability. Received date: 11 January 2022 Accept date: 14 February 2022

#### Introduction

Brushing your teeth with toothpaste is important to prevent dental and oral infections. Toothpaste serves to clean teeth, reduce plaque formation, prevent caries, clean and polish tooth surfaces, eliminate or reduce bad breath, give a fresh taste to the mouth and maintain oral health. Toothpaste products will continue to change their chemical composition due to producer competition. Toothpaste contains antibacterial properties of various substances with different abilities to inhibit the growth of germs in the oral

\*Corresponding author: I Dewa Ayu Ratna Dewanti Faculty of Dentistry of Jember University. Kalimantan street 37 Jember. Indonesia E-mail : <u>idewadewanti@yahoo.com</u> cavity, as well as a number of ingredients with specific purposes to solve certain problems. The many choices of toothpaste with various ingredients make it difficult for patients to choose the right toothpaste<sup>1.2</sup>. On the other hand, toothpaste products on the market can cause side effects such as canker sores, allergies, fluorosis, thyroid disorders to cancer. Some whitening products have been reported to have a pH as low as 4.0, while others have been reported to have a pH of 7.5. It has been the higher the peroxide reported that concentration, the more acidic the pH of the bleaching product.26 Some office whitening products containing 35% hydrogen peroxide may have a low pH<sup>3</sup>. The use of natural ingredients such as coffee bean husks is expected to minimize side effects.

Coffee bean skin contains amino acids, proteins, flavonoids, catechins equivalent to

Volume  $\cdot$  15  $\cdot$  Number  $\cdot$  1  $\cdot$  2022

seeds and leaves (results of previous research). The skin of coffee beans like coffee beans contains an antioxidant, Trigonelline which is useful for protecting teeth. Although coffee drinkers generally have a slightly black tooth color due to these substances, this compound will protect teeth against S. mutans<sup>4.5</sup>. On the other hand, coffee does not interfere with the results after teeth whitening, so there is no need for dietary restrictions<sup>6</sup>. Previously we also researched about coffee plants, including: Steeping from green and black robusta coffee beans increase viability of Peripheral Blood Mononuclear Cells (PBMC) and salivary leukocytes which is induced by streptococcus mutans,Robusta Coffee Beans (Coffea canephora) Decrease IL-1α (Interleukin-1 $\alpha$ ) Expression and Increase the Number of Fibroblasts in Healing Process in Dental Pulp in Wistar Rat, the effect of steeping robusta coffee beans on monocytes:expression of IL-1ß and TNF-α streptococcus mutans<sup>7-9</sup>. against Meanwhile, as a basic ingredient for toothpaste, coffee bean rind must have qualities regarding physical properties and biocompatibility that are equivalent to or better than toothpastes on the market. Therefore, this study aimed to analyze the quality (organoleptic, pH, homogeneity, spreadability, viscosity), microorganism inhibition and cell viability of Robusta coffee bean skin toothpaste.

#### Materials and methods

Research tools include: mortar and stamper, pH meter, rotary evaporator, a set of macerator tools, viscometer, digital caliper, autoclave, petri dish, incubator, ose needle, pH indicator paper, pH meter, glass object, microtiter plate, inverted microscope, micro pipette. The ingredients used were DMSO. Robusta coffee bean skin, 96% ethanol, menthol, calcium carbonate. sodium CMC, NaCL. sodium benzoate, sodium lauryl sulfate, saccharin, sorbitol, NA media, Streptococcus mutans isolate, Histopaque (Sigma), ficolhipaque ( HBSS (Hank's Balanced Sigma). Salt Solution/Gibco), fungizone (Sigma), Trypan Blue, heparin, RPMI (Sigma).

Before making toothpaste, the coffee bean skin was screened for phytochemicals using the TLC method (Thin Layer Chromatography for flavonoids, polyphenols, triterpenoids, alkaloids, anthraquinones),

#### 1. Making Coffee Bean Skin Toothpaste

Sodium CMC is sprinkled over hot water as much as 20 times the amount of Na CMC, let for 30 minutes, then grind until stand homogeneous. Menthol is dissolved in ethanol. The extract was diluted with ethanol, added sorbitol and menthol solution, added Na CMC dispersion, stirred until homogeneous. Added calcium carbonate, sodium benzoate, sodium lauryl sulfate and saccharin. stirred until toothpaste is formed. Coffee bean skin toothpaste used with concentrations of 5%, 10%, 15%.

2. Organoleptic Test

Organoleptic test observed shape, color and odor.

3. pH measurement

The paste preparation (1 gram) was dissolved with 10 ml of distilled water and measured using an OHAUSS pH meter.

4. Viscosity test

Viscosity test using vicotester VT 03.

5. Spreadability test

Spreading power by measuring the diameter of toothpaste smeared on a glass (1 gram), covered with a glass and then given a load of 200 g.

6. Inhibitory Test.

The method used is disc diffusion, by measuring the diameter of the resistance against, mutans (pure isolate cultured in the S. Biomedical Lab, Faculty of Dentistry, University microorganisms of Jember). The were standardized with standard Mc. Farland 0.5 and obtained a concentration of 1.5x108CFU/mL. A paper disk (6mm) was inserted into the sample, then placed in a petri dish containing the solidified media. Incubated for 24 hours at a temperature of 35°C-37°C in an incubator, then observed the area of bacterial growth inhibition using a digital caliper (Balouir et al., 2016). David and Stout (1971) in Ouchari et al., 2019, classified the zone of inhibition (ZOI): >20 mm (very strong); 10-20 mm (strong), 5-10 mm (medium); and <5 mm (no response).

7. Cell viability test

Peripheral blood collection of healthy people as much as 6 cc is then mixed with anticoagulant (heparin). Layer 3 ml of histopaque on the falcon, 3 ml of ficoll on top of histopaque, 6 ml of blood on it carefully. ficoll-hypaque centrifugation, take the monocyte layer, add 1:1

Volume · 15 · Number · 1 · 2022

Journal of International Dental and Medical Research ISSN 1309-100X	Toothpaste's Containing Coffee bean Skin
http://www.jidmr.com	I Dewa Ayu Ratna Dewanti and Ristya Widi Endah Yani

HBSS and pipette. Centrifugation, add HBSS, fungizone and penstripe Cells were placed on a 96-well microtiter plate 8 x 105 cells/well for 45 minutes at 37°C and washed 4 times with medium. Cells were placed in 24-well microtiter plates, treated with S. mutans (100µl/well) + toothpaste (5%, 10%, 15%), except C- (S. mutans only), then incubated for 5 hours. was stained with Trypan Blue. Live cells (in white) were counted under an inverted microscope with magnification 400 times from an average of 4 fields of view.

The data obtained were ANOVA followed by LSD test and descriptive.

#### Results

Analysis of the levels of polyphenols  $(393.0166 \pm 85.5224 \text{ mg GAE/g} \text{ and flavonoids} (0.788592 \pm 0.114787 \text{ mb QE/g})$  using 425 nm wavelength spectrophotometry with comparison of Quercetin (flavonoid) and gallic acid (polyphenol).

The results of the physicochemical test: the preparation of coffee bean skin toothpaste has a soft, homogeneous texture, smells of coffee, has in accordance with the pH a pH of 7, requirements for toothpaste preparations according to SNI 12-3524-1995, namely 4.5-10.5. All toothpastes are homogeneous. The viscosity of the coffee bean skin toothpaste is 200 dPa, according to the standard (200-500 dPa's). The spreadability that meets the standards is 15% (5) coffee bean husk toothpaste, while 5% (4) and 10% (4.5) gels are good between 5 to 7 cm.

sample name	Organoleptic
Coffee bean skin Toothpaste 5 %	Shape: Toothpaste
	Color: light brown
	Smell: Mint coffee scent
	Shape: Toothpaste
Coffee bean skin	Color: Brown
Toothpaste 10 %	Smell: Coffee, mint
	Shape: Toothpaste
Coffee bean skin Toothpaste 15 %	Color: dark brown
	Smell: Aroma of coffee,
	mint
Commercial toothpaste	Shape: Toothpaste
	White color
	Smell: Mint

 Table 1. Organoleptic of Coffee bean skin.

Volume · 15 · Number · 1 · 2022

Inhibition against S. mutans, there was a significant difference from the Anova analysis and LSD test (p 0.05). All groups can inhibit S. mutans except C-. The largest diameter was found at C+ followed by T1, T2 and T3, categorized as having a strong inhibitory power (>10). The higher the concentration of coffee bean rind toothpaste, the larger the diameter of the barrier (figure 1).







The results of Anova and LSD analysis showed cell viability, C-(S. mutans) had more dead cells (dark in color), while C+(commercial toothpaste) had the most live cells (light/white in color). However, the number of live C+ cells with T1( Coffee bean skin Toothpaste 5 %), T2(Coffee bean skin Toothpaste 10 %), T3(Coffee bean skin Toothpaste 15 %) was not significantly different (Figures 2 and 3).



**Figure 2**. Cell viability was depicted with white cells (red arrows), while dead cells were dark (black arrows).



**Figure 3.** Bar chart of the cell viability of the coffee bean shell toothpaste.

#### Discussion

The preparation of coffee bean skin toothpaste is soft, homogeneous, has a distinctive coffee smell, has a pH of 7 in accordance with the pH requirements for toothpaste preparations according to SNI 12-

3524-1995, namely 4.5-10.5, pH so that it does not irritate the oral mucosa. The smell of this coffee can be a special attraction, because in this world there are many coffee lovers, so we hope that many people will like it. Meanwhile, the dispersion power that most meets the standards similar to commercial toothpaste is coffee bean husk toothpaste with a concentration of 15%, meaning that the ability to spread toothpaste when applied to a toothbrush is good.

Coffee bean skin toothpaste with 5%, concentrations of 10%, 15%, and commercial toothpaste had a strong ability to inhibit S. mutans bacteria, although it was not as strong as commercial toothpaste. The higher the concentration of the coffee bean husk, the stronger the ability to inhibit the growth of S. mutans. This is presumably because of its bioactive content, the higher the concentration, the higher the bioactive content. Antioxidants (caffeic acid, chlorogenic acid, coumaric acid, sinapat acid, chlorogenic acid, flavonoids, melanoidin, faran, pitol and maltol) react directly with free radicals and form stable molecules. Chemical content of coffee such as flavonoids, xanthine, antioxidants, alkaloids, polyphenols act anti-inflammatory, antibacterial; as platelet aggregation inhibits the growth of Streptococcus mutans<sup>7-11</sup>. The content of alkaloids in coffee include caffeine and trigonelin. Caffeine is a purine alkaloid, has antimicrobial activity, which inhibits esterase enzymes along with DNA and RNA polymerase, inhibits cellular respiration, plays a role in DNA intercalation, causes damage and lysis<sup>12-15</sup>. Trigonelin which is a pyridine alkaloid can disrupt the stability of the bacterial cytoplasmic membrane which causes an imbalance in the metabolic function of bacteria so that the growth of bacteria inhibited<sup>16</sup>. Flavonoids are antioxidants that have antibacterial and antifungal effects because they contain phenol groups. Phenol groups can coagulate proteins, and reduce the surface tension of microbial cells<sup>13.17</sup>. In addition, the lipophilic nature of flavonoids interferes with microbial membranes, which will inhibit bacteria from forming a defense system<sup>18</sup>. Triterpenoids react with porins (trans membrane proteins) on the outer membrane of the bacterial cell wall, forming a strong polymeric bond, resulting in the destruction of the porin<sup>7-9.19</sup>. The ability to maintain cell viability by coffee bean rind toothpaste, especially at a concentration of 15%, is thought to be due to the presence of

 $Volume \cdot 15 \cdot Number \cdot 1 \cdot 2022$ 

Ca2+ flavonoids. Flavonoids activate in mitochondria which makes cells able to produce ATP so they can survive<sup>20</sup>. In addition, flavonoids have the function of preventing precipitation, damage to cell synthesis, protein denaturation, damage to cell metabolism, increasing the work of Transforming Growth Factor-beta 1 (TGF-1) which is the main factor to stimulate fibroblast proliferation. This antioxidant content is thought to work through suppression. on the production (Nitric Oxide). NO This suppression of mechanism can go through several possibilities. First, it may inhibit the action of the cytokineinduced NO synthase (iNOS) enzyme by controlling iNOS mRNA. Another possibility is to inhibit arginine transport through the CAT-2 mRNA (Cationic Amino Acid Transporter-2 mRNA) control mechanism. This can occur because it is suspected that monocyte cell damage is caused by the effects of nitric oxide (NO) free radicals. Usually NO can be produced by cells exposed to pathogens, which is caused by the formation of various cytokines and endotoxins of pathogenic bacteria that are damaging to cells. K attack pathogens, NO will be synthesized in large quantities. NO which is synthesized in large quantities will cause impaired function and cell proliferation (cytostatic)<sup>21.22</sup>.

#### Conclusions

Toothpaste's contain coffee bean skin has physico-chemical properties, inhibited to S. mutans and increase cell viability equivalent to commercial toothpaste.

#### Acknowledgments

Thanks to LP2M University of Jember for supporting this research through internal grants.

#### **Declaration of interest**

There is no conflict of interest between all authors and all involved in the research or writing of articles.

#### References

 Vranic Edina, Amela Lacevic, Aida Mehmedagic, Alija Uzunovic. Formulation Ingredients for Toothpastes and Mouthwashes. Bosnian JournaL of Basic Medical Sciences. 2004;4(4):51-58.

Volume  $\cdot$  15  $\cdot$  Number  $\cdot$  1  $\cdot$  2022

- Maldupa Ilze, Anda Brinkmane, Inga Rendeniece, Anna Mihailova. Evidence based toothpaste classifi cation, according to certain characteristics of their chemical composition. Stomatologija, Baltic Dental and Maxillofacial Journal. 2012;14:12-22.
- Price Richard B.T., DDS, MS, MRCD(C), FDS, RCS(Edin), Mary Sedarous, B.Sc., Gregory S. Hiltz, B.Sc.(Hon.), DDS. The pH of Tooth-Whitening Products. J Can Dent Assoc 2000;66:421-6. https://www.cda-adc.ca/jcda/vol-66/issue-8/421.html.Tooth-Whitening Products. J Can Dent Assoc 2000; 66:421-6. https://www.cda-adc.ca/jcda/vol-66/issue-8/421.html.
- Anila PC Namboodiripad and Sumathi Kori. Can coffee prevent caries?. J Conserv Dent. 2009;12(1):17–21.
- Najmeh Akhlagh, Marziye Sadeghi, Fataneh Fazeli, Shiva Akhlaghi, Maryam Mehnati, and Masuod Sadeghi. The antibacterial effects of coffee extract, chlorhexidine, and fluoride against *Streptococcus mutans* and *Lactobacillus plantarum*: An *in vitro* study. Dent Res J (Isfahan). 2019; 16(5):346–353.
- Câmara João Victor Frazão, Luine de Paiva Pereira Santos de Souza, Daniel Otero Amaral Vargas, Isabel Ferreira Barbosa, Gisele Damiana da Silveira Pereira. Effect of Tooth Enamel Staining by Coffee Consumption During at-Home Tooth Bleaching with Carbamide Peroxide. Rev Odontol UNESP. 2020;49:e20200024. https://doi.org/10.1590/1807-2577.02420.
- Dewanti I Dewa Ayu Ratna, I Dewa Ayu Susilawati, Pujiana Endah Lestari, Ristya Widi Endah Yani, Erawati Wulandari, Roedy Budirahardjo, Dyah Setyorini & Sunlip Wibisono. Robusta Coffee (Coffeacanephora) Decreasing IL-1α (Interleukin-1α) Expression and Increasing the Number of Fibroblasts in Healing Process in Dental Pulp in Wistar Rats. J. Math. Fund. Sci. 2019;51(1):68-76.
- Dewanti I Dewa Ayu Ratna, Pujiana Endah Lestari, Roedy Budirahardjo, Dyah Setyorini, RistyaWidi Endah Yani, Sunlip Wibisono, Maizirwan Mel. The effect of steeping robusta coffee beans on Monocytes: Expression Of IL-1β and TNF-α Against Streptococcus mutans. Coffee Science, Lavras. 2019;14(4):477-483.
- Dewanti I-Dewa A.R., Roedy Budirahardjo, Ristya WE. Yani, Pujiana E. Lestari, Dyah Setyorini, Eri H. Jubhari, Sunlip Wibisono,. Steeping from green and black robusta coffee beans increase viability of Peripheral Blood Mononuclear Cells (PBMC) and salivary leukocytes which is induced by streptococcus mutans. Journal of Dentomaxillofacial Science (J Dentomaxillofac Sci). December 2019;4(3):154-158.
- Namboodiripad P., K. Srividya: Can Coffee Prevent Caries? -An In-Vitro Study. The Internet Journal of Dental Science. 2009;7(2):1-6. DOI: 10.5580/101c.
- 11. Mulato Śri, Edy Suharyanto, 2015. Coffee, Brewing & Health. Indonesian Coffee and Cocoa Research Center. Jember.
- Aniszewski Tadeusz . 2007. Alkaloids Secrets of Life Alkaloid Chemistry, Biological Significance, Applications and Ecological Role. First edition. Elsevier Radarweg 29, PO Box 211, 1000 AE Amsterdam, The Netherlands The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, UK.
- Septiadi T, Pringgenies D. dan Radjasa O.K.. Phytochemical Test and Antifungal Activity of Keling Sea Cucumber Extract (Holoturia atra) from Bandengan Beach, Jepara against Candida albicans. Journal of Marine Research. 2013;2(2):76-84.
- Novelli Silvia, Canuti Lorena, Canini Antonella. Identification of Alkaloid's Profile in Ficus benjamina L. Extracts with Higher Antioxidant Power. American Journal of Plant Sciences. January 2. 2015;5(26): 4029-4039.
- Tanauma H, Citraningtyas G, Lolo W. Antibacterial activity of robusta coffee bean extract (Coffea canephora) against Escherichia coli. PHARMACON Pharmaceutical Scientific Journal. 2016; 5(4):2302-2493.
- Nuhu A. Bioactive micronutrients in coffee: Recent analytical approaches for characterization and quantification. ISRN Nutrition. 2014;Article ID 384230:1-13.
- 17. Waluyo L. General Microbiology, Revised Edition. UPT, Muhammadiyah University Publisher Malang; 2007.
- 18. Lutfiyanti R, Widodo F, Eko N. dan Dewi. Antifungal Activity

Bioactive Compounds Gelidium latifolium Extract against Candida albicans. Journal of Fishery Products Processing and Biotechnology. 2012;1(1):1-8.

- Das Annada and Kaushik Satyaprakash, "Antimicrobial properties of natural products: A review," Pharma Innov. J.2018;7(6):532–537.
- Kicinska Anna and Wieslawa Jarmuszkiewicz. Flavonoids and Mitochondria: Activation of Cytoprotective Pathways?. Molecules. 2020 Jul;25(13): 3060. Published online 2020 Jul 4. doi: 10.3390/molecules25133060. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7412508/.
- Akhlaghi M, Bandy B, Review article: mechanisms of flavonoid protection against myocardial ischemia– reperfusion injury. Journal Molecullar and Cellular Cardiology. 2009;46:309–317.
- 22. Halliwell B, Gutteridge JMC,. 2015. Free Radicals In Biology And Medicine. Ed ke-4. Oxford, UK: Oxford University Press.