

ORIGINAL ARTICLE

pISSN: 1907-3062 / eISSN: 2407-2230

Detection of immunogenic protein from salivary gland of *Aedes albopictus*

Rike Oktarianti¹, Rochmatul Nuryu Khasanah¹, Syubbanul Wathon¹,
and Kartika Senjarini^{1*}

ABSTRACT

BACKGROUND

Dengue virus is transmitted by several species of *Aedes* mosquitoes, with *Aedes albopictus* as secondary vector. During blood feeding, these vectors inject saliva into the vertebrate hosts. The saliva contains anticoagulant, anti-inflammatory and immunogenic factors. The objective of this research was to detect immunogenic proteins from *Ae. albopictus* salivary glands reacting with sera of people living in dengue endemic areas.

METHODS

The identification of immunogenic proteins of *Ae. albopictus* salivary gland used one-dimensional gel electrophoresis (sodium dodecyl sulfate polyacrylamide gel electrophoresis), and western blot analysis, respectively. To determine the immunogenic nature of the candidate proteins, the antigens from the salivary gland of *Ae. albopictus* were reacted with sera from healthy persons, dengue hemorrhagic fever (DHF) patients, and neonates, each of the groups comprising 10 samples.

RESULTS

The protein profiles of *Ae. albopictus* salivary glands showed 13 bands with molecular weights from 16 kDa up to 97 kDa, i.e. 16, 17, 26, 28, 31, 32, 45, 55, 60, 67, 73, 76, and 97 kDa. According to western blot analysis result, the 31 kDa proteins were recognized in all endemic population sera, both in DHF patients and healthy persons. In contrast, protein bands of 47 and 67 kDa were only recognized by the sera of DHF patients.

CONCLUSION

Three immunogenic proteins of 31, 47 and 67 kDa were detected from *Ae. albopictus* salivary glands. These immunogenic proteins may be developed as candidate biomarkers for bite exposure to *Ae. albopictus* and as vector-based DHF vaccines.

Keywords : Immunogenic protein, salivary gland, *Aedes albopictus*

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Jember University

Correspondence :

*Kartika Senjarini
Department of Biology,
Faculty of Mathematics and Natural
Sciences, Jember University
JL. Kalimantan No.37, Tegalboto,
Sumbersari, Jember, East Java,
Indonesia
Phone/Fax: +6287857703792
Email: senjarini@unej.ac.id
ORCID ID: 0000-0001-7041-1719

Date of first submission, March 5, 2021

Date of final revised submission, November 8, 2021

Date of acceptance, November 20, 2021

This open access article is distributed under a Creative Commons Attribution-Non Commercial-Share Alike 4.0 International License

Cite this article as: Oktarianti R, Khasanah RN, Wathon S, Senjarini K. Detection of immunogenic protein from salivary gland of *Aedes albopictus*. Univ Med 2021;40:234-42. doi: 10.18051/UnivMed.2021.v40.234-242.



INTRODUCTION

Dengue hemorrhagic fever (DHF) often affects people in tropical and subtropical areas.⁽¹⁾ Areas at risk of dengue infection comprise 128 countries, including Indonesia.⁽²⁻⁴⁾ The latest data show that about two-thirds of the world's population has the potential to be infected by dengue, with the number of cases reaching 100-390 million each year.^(5,6) Dengue hemorrhagic fever (DHF) is caused by infection with the dengue arbovirus which has four serotypes, i.e. dengue virus (DENV)-1, DENV-2, DENV-3, and DENV-4.^(7,8) The dengue virus infects the human body through transmission mediated by *Aedes* mosquitoes (*Aedes aegypti* and *Aedes albopictus*).⁽⁹⁾ *Ae. aegypti* is the primary vector because its habitat is around settlements so that it often comes into contact with humans. *Ae. albopictus* acts as a secondary vector because its habitat is in gardens or forests far from human settlements. At present *Ae. albopictus* has adapted to rural and suburban environments.⁽¹⁰⁾ Dengue virus transmission to human hosts occurs when the vector is blood feeding.⁽¹¹⁾ Successful transmission of the dengue virus occurs due to the presence of important proteins in *Aedes* salivary glands.⁽¹²⁾ *Aedes* saliva has several proteins able to act as antihemostatic, vasodilator, anti-inflammatory, and immunomodulatory substances which play a role in the successful transmission of the dengue virus.⁽¹³⁾ Antihemostatic substances play a role in preventing coagulation of host blood when mosquitoes are blood feeding. Vasodilator substances can cause vasodilation of blood capillaries. Immunomodulators are substances that can suppress the host immune response so that it can increase the transmission of pathogens into the host by modulating the host immune response so as to facilitate their transmission into the host.^(12,14)

Immunomodulatory proteins in arthropod saliva cause a shift in the immune response towards T helper type 2 (Th2), characterized by increased production of Th2 cytokines such

as interleukin-4 (IL-4) and IL-10 and decreased production of Th1 cytokines such as interferon gamma (IFN- γ) and IL-12.⁽¹³⁾ Repeated exposure can increase production of specific antibodies in the host against salivary proteins. This is because when the host is repeatedly exposed, there will be an increase in the proliferation of B cells into plasma cells and memory cells which will cause the production of immunoglobulin G (IgG). The increase in IgG production will trigger the host's immunity against pathogens. IgG production can also be used as a marker for the presence of immunogenic proteins in mosquito saliva.^(15,16)

Previous research identified two immunogenic proteins from *Ae. aegypti* salivary glands of 31 and 56 kDa, respectively.⁽¹⁷⁾ Both of these proteins were recognized in sera samples of healthy persons and DHF patients living in DHF endemic areas, but not in sera samples of healthy persons in non-endemic areas and in infants (neonates). This shows that persons living in endemic areas develop specific antibodies. Because *Ae. albopictus* is also known to be a potential vector of DHF, it is suspected that the salivary glands of *Ae. albopictus* contain immunogenic proteins similar to those in *Ae. aegypti*. Therefore, further research is needed to detect the immunogenic protein fraction in the salivary glands of *Ae. albopictus*. The immunogenic protein fraction can be used as a candidate target for the development of vector-based DHF vaccines and as a biomarker of exposure to *Ae. albopictus*. Therefore, the aim of this study was to detect the immunogenic proteins from the salivary glands of *Ae. albopictus*.

METHODS

Research design

A laboratory study was conducted from January to November 2020 at the Biotechnology Laboratory, Department of Biology, Faculty of Mathematics and Natural Sciences (FMIPA), Jember University.

Research subjects

Thirty sera samples were taken from healthy persons, DHF patients, and infants (neonates) living in Jember, consisting of 10 samples each. The inclusion criteria of healthy sample sources were persons aged 15-50 years who never had DHF, whereas DHF patients as sample source were persons aged 15-50 years diagnosed as having DHF by a doctor. The exclusion criteria of sample sources were persons under 15 years old or more than 50 years old and non-residents of Jember.

The study commenced with the rearing of *Ae. albopictus* mosquitoes and was then continued by the identification of *Ae. albopictus* based on morphological characters, isolation of the salivary glands, extraction of salivary gland proteins, protein profile analysis using sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE), and finally western blot analysis to identify immunogenic proteins.

Rearing of *Ae. albopictus*

Rearing was conducted at the Animal Care Unit, University of Jember, at room temperature (27°C). Rearing was begun with the landing collection of *Ae. albopictus* mosquitoes using aspirators in gardens around the University of Jember. Male mosquitoes were fed 10% sucrose solution, while female mosquitoes were fed by blood feeding.

Species identification based on morphological characters

Identification of species based on morphological characters was carried out prior to isolation of the salivary glands. *Ae. albopictus* and *Ae. aegypti* are easily distinguished by the mesonotum, mesepimeron, and femur. The mesonotum in *Ae. aegypti* has rows of white hairs forming a white line pattern, namely two curved white lines like a lyre and two parallel lines, while in *Ae. albopictus* it has only one white line. The mesepimeron of the two mosquitoes is very different: *Ae. aegypti* has a collection of white scales forming two separate

white patches, whereas *Ae. albopictus* has a single white patch. *Ae. aegypti* has rows of white hairs that form a long white stripe on the anterior femoral surface of the middle leg, while *Ae. albopictus* does not have such a femoral stripe.⁽¹⁰⁾

Isolation and extraction of *Ae. albopictus* salivary gland proteins

Isolation of the salivary glands was by microdissection using a dissection needle, in which the investigator's left hand gently presses on the thoracic part and the right hand slowly pulls on the head, so that the 6 clear salivary glands appear.⁽¹⁸⁾ The salivary glands that have been obtained are then separated from the fat body or other tissues until they are clean and then put into a microtube containing 10 µl of 1 mM phenylmethylsulfonyl fluoride (PMSF) in phosphate buffered solution (PBS) in the ratio of 1:1 for the 10 pairs of salivary gland samples and stored at -20°C until used. Salivary gland protein extraction was carried out by adding 10 µl of sample buffer into a microtube containing 10 pairs of salivary gland samples + 10 µl of 1 mM PMSF in PBS. The extraction process was followed by boiling using a thermoshaker at a temperature of 95°C for 4 minutes. After boiling, the salivary gland protein extract was centrifuged to obtain the supernatant.

Blood sera preparation

Thirty serum samples were taken from healthy persons, DHF patients, and infants (neonates) living in Jember, East Java, a region within the endemic range for dengue, each group comprising 10 samples. These samples were collected in 4 months from July-October 2020. The serum samples of DHF patients were taken from the Dokterku Clinic and Citra Husada Hospital, Jember. Neonatal serum samples were taken from the Umi Soemarno Clinic, Jember. Blood samples were taken from the brachial veins in the upper arm, except for neonatal sera which were taken from the portion of the umbilical cord that remained with the baby after birth. Each blood

sample had a volume of 3 ml and was stored in a vacutainer without anti-coagulant. The blood samples were allowed to stand for 30 minutes, then the top clear layer was taken and centrifuged at 3200 rpm for 15 min at 4°C. The resulting supernatant after centrifugation constituted the serum. The serum samples were stored at -20°C until used.

Protein profile analysis of *Ae. albopictus* salivary glands

The salivary gland protein profile of *Ae. albopictus* was analyzed using SDS-PAGE. Electrophoresis of the protein samples was done using 12% separating gel and 4% stacking gel. Each well in the gel contained 20 µl of protein sample (10 pairs of salivary gland samples in 10 µl of PMSF in PBS + 10 µl of sample buffer). Electrophoresis was carried out at a constant voltage of 150 V for 100 minutes in a running buffer. Then the gel underwent a coloring process using coomassie brilliant blue dye solution (staining) for 60 minutes. The washing process (destaining) was performed 3 times using tris-buffered saline (TBS) every 30 minutes to remove the dye solution that was not bound to protein.

Identification of the immunogenic protein fraction from *Ae. albopictus* salivary glands

Identification of immunogenic protein fractions was carried out by western blot analysis, with several modified steps using the semi-dry method.⁽¹⁹⁾ Proteins in the polyacrylamide gel from SDS-PAGE analysis were transferred to the polyvinylidene fluoride (PVDF) membrane for 1 hour with a constant current of 100 mA. The PVDF membranes were cut according to the size of the polyacrylamide gel then immersed in methanol for 1 minute. The polyacrylamide gel, PVDF membrane, 2 sheets of Whatman filter paper, and fiber pad were immersed in the transfer buffer prior to western blot analysis. The PVDF membrane, SDS-PAGE acrylamide gel, filter paper, and blotting paper (fiber pad) were arranged sequentially.

The immunoblotting process was followed by a reaction between the salivary gland proteins of *Ae. albopictus* on the membranes and primary antibodies derived from sera of healthy persons in DHF endemic areas, as well as DHF patients and neonates, that were dissolved in 5% skim milk in TBS (1: 250) then incubated overnight at 4°C. The membrane was washed with 3 TBS times for 5 minutes each. The secondary antibody used was goat – anti human IgG alkaline phosphatase conjugate (ROCKLAND, USA) which was dissolved in 5% skim milk in TBS (1: 2500). The membranes that had been treated with secondary antibodies were then incubated for 2 hours. Visualization was done by adding the nitro blue tetrazolium-(5-bromo-4-chloro-3-indolyl-phosphatase) (NBT-BCIP) alkaline phosphatase substrate (ROCKLAND, USA) to the brackets and incubating in a dark room for 1 minute until immunogenic protein bands appeared. The membranes were washed with distilled water to stop the reaction. Positive western blot results were shown by bands on the PVDF membrane, indicating the presence of binding between the antigens from the *Ae. albopictus* salivary gland proteins and primary antibodies.

Ethical clearance

All participants gave written, informed consent to take part in the study. The collecting protocol was approved by the Ethical Committee of Medical Research, Faculty of Dentistry, University of Jember under No. 1034/UN25.8/KEPK/DL/2020

RESULTS

The structure of the *Ae. albopictus* salivary glands consists of two parts connected by the salivary duct. Each part has three lobes, namely two lateral lobes (L) and one medial lobe (M). Each of the lateral lobes consists of two parts, namely the proximal lobe (PL) and distal lobe (DL). The structure of the salivary glands can be seen in Figure 1.





Figure 1. Structure of salivary glands of *Ae. albopictus* (A) and *Ae. aegypti* (B). Salivary duct (sd), lateral lobe (L), proximal lobe (PL), distal lobe (DL), medial lobe (M). 50x magnification on the Nikon SMZ735 microscope. Camera: Xiaomi Note 6 Pro.

The protein profile analysis of salivary gland extract of *Ae. albopictus* was conducted by using SDS PAGE. The results showed that the salivary gland proteins from *Ae. albopictus* have 13 protein bands with molecular weights of 16, 17, 26, 28, 31, 32, 45, 55, 60, 67, 73, 76, and 97 kDa, respectively, as can be seen in Figure 2.

Analysis of the immunogenicity of the salivary gland proteins of *Ae. albopictus* was performed by western blot analysis of the cross-reaction between the antigens (salivary gland proteins) and the sera of healthy persons living in endemic areas, DHF patients, and neonates (as negative controls). The 31 kDa immunogenic protein can be recognized by all endemic population sera, both the sera of DHF patients and healthy persons. Meanwhile, the 47 and 67 kDa protein bands were only recognized by the sera of DHF patients (Figure 3).

DISCUSSION

The results of SDS-PAGE analysis of the salivary gland protein extracts from *Ae. albopictus* showed 13 protein bands, which are thought to have certain roles such as in anti-host homeostasis and are involved in the metabolism of the mosquito's body. The role of the 97 kDa and 79 kDa proteins is unknown. The 60-70 kDa proteins are thought to contain apyrase which functions to inhibit platelet aggregation. (16,20-22) The proteins with molecular weights ranging from 40 to 55 kDa are serpins which function as



Marker: Prestained Blue Eye PS-104 (M)
Figure 2. Salivary gland protein profile of *Ae. albopictus* (Samsung A50 camera)




Figure 3. Results of the immunogenicity analysis of the *Ae. albopictus* salivary gland proteins in individual sera (a, b, c and d) of DHF patients, healthy persons, and neonates (Samsung A50 camera)

Indonesian abbreviations: Pasien DBD [Demam Berdarah Dengue] = Dengue Hemorrhagic Fever patients
Sehat Endemik = healthy endemic [sera]

protease inhibitor and anticoagulant. Approximately 500 serpins are currently known, consisting of between 350 and 400 amino acids.⁽²³⁾ The 31 kDa protein is a D7 protein that acts as vasodilator, immunomodulatory and anticoagulant substance.⁽¹⁷⁾ The 27-29 kDa bands contain immunogenic proteins and are present in all *Aedes* mosquitoes.^(21,22) The 17 kDa band has

D7 family proteins and lectins, which function in opsonization and as antimicrobials. These proteins control microbial growth on sucrose in the mosquito's midgut.⁽²⁴⁾

Apyrase (ATP-diphosphohydrolase) contained in the 60-70 kDa bands is an enzyme released by *Ae. albopictus* and other hematophagous arthropods during blood feeding

that functions as platelet aggregation inhibitor.⁽²⁵⁾ The 67 kDa molecular weight protein in our study is thought to contain apyrase and α -glucosidase. The latter acts as an allergen that will induce IgE in humans. This enzyme is also released when mosquitoes suck blood and sugar so that the sugar and blood are not mixed in the mosquito's mid gut.^(26,27) The apyrase of *Ae. albopictus* is in the molecular weight range of 61 kDa to 68 kDa.^(16,20,28,29)

Western blot analysis of the *Ae. albopictus* salivary gland proteins showed 3 immunogenic proteins of 31, 47 and 67 kDa, respectively. The 47 kDa band contains the main protein thought to be a serpin which functions as an anticoagulant.⁽²³⁾ In addition, serpins in the body of insects play a role as proteinase inhibitors. Serine is a proteinase that breaks down the polypeptide bonds produced by pathogens such as viruses. The action of serine is inhibited by a serine-protease inhibitor, namely serpin.⁽³⁰⁾ Serpins that enter the host are thought to inhibit the action of serine so that the transmission of the dengue virus runs smoothly. DHF patients also activate B cells to produce more antibodies, so the serum of DHF patients are able to recognize more immunogenic proteins. Samples taken from each individual presumably reflect the various responses of the individuals to the different antigens carried by the mosquitoes.

From the western blot analysis results in Figure 3, it is apparent that the immunogenicity analysis of sera of the endemic population shows different bands for each individual. This is presumably because of the differing responses of each individual to exposure to antigens carried by mosquitoes. The intensity of exposure to *Aedes* salivary antigens in endemic areas will influence the human immune system to form specific antibodies (IgG) against these antigens.^(15,31)

Exposure to antigens for the first time (primary) will activate the T cells and B cells. The B cells will differentiate and proliferate to produce IgG in small amounts. The T cells will be activated by the antigens to become memory cells upon secondary or multiple exposure to the

antigens. Re-exposure (secondary) to the same antigen will produce more IgG.⁽³²⁾ Apyrase, serpins, and the D7 family are thought to be the most immunogenic proteins in the salivary glands, so that even low titers can trigger the formation of antibodies in the host body. These three proteins are thought to be immunogenic because they can modulate the immune response of the host.

The band of 31 kDa is the D7 immunogenic family which facilitates the process of blood feeding. The D7 family functions as a vasodilator, immunomodulator, and anticoagulant.⁽³³⁾ The protein of 47 kDa is serpin, that acts as anticoagulant and anti-inflammatory,^(23,25,26) where as the band of 67 kDa is apyrase, that acts as allergen and anticoagulant. The abovementioned three proteins are thought to trigger modulation of the host's immune response because they comprise proteins that act as allergens and immunomodulators. Allergens will react with IgE and cause inflammation in the host.^(26,27) The immunomodulators function to modulate the host's immune response so that they can increase the transmission of pathogens into the host.⁽²⁴⁾

Neonate serum samples were not recognized as containing immunogenic proteins, presumably because there was no binding between antigens of salivary gland proteins and serum antibodies. This is in accordance with previous research which stated that no immunogenic protein was recognized by neonate sera samples.⁽³³⁾ Neonate samples were obtained from the umbilical cord of the baby at birth, thus the neonate samples had never been directly exposed to *Ae. albopictus* salivary gland proteins so that no immunogenic proteins were identified. The three immunogenic proteins of 31, 47 and 67 kDa from the salivary glands can be used as biomarker protein candidate for bite exposure to *Ae. albopictus* and can be developed into a vector-based DHF vaccine. Therefore, further studies are needed to determine the identities of these proteins and their biological functions in dengue virus transmission and blood feeding using transcriptomic or proteomic analysis. The limitation of this study was the sera samples were

only taken from one DHF endemic region i.e. Jember, East Java - Indonesia. Therefore, further investigation in the several endemic regions is needed to find out the relevancy of the data”.

CONCLUSIONS

This study demonstrated 3 immunogenic proteins that were identified from the salivary glands of *Ae. albopictus* of 31, 47 and 67 kDa, respectively. The 31 kDa protein fraction could be recognized by all endemic population sera, both the sera of endemic DHF and healthy patients. In contrast, the 67 and 47 kDa fractions were only recognized by the sera of DHF patients. These immunogenic proteins can be used as biomarker candidates for bite exposure to *Ae. albopictus* and can be developed into a vector-based DHF vaccine.


CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

ACKNOWLEDGEMENT

This study was financially supported by "Hibah Penelitian Pendukung IDB" - Internal Financial (PNBP) of Jember University.

CONTRIBUTORS

RO conceived and planned the experiments and wrote the manuscript. RNK carried out the experiment and contributed to the interpretation of the results. SW contributed to sample preparation and interpretation of the results. KS conceived the original idea and supervised the project. All authors have read and approved the final manuscript. 

REFERENCES

1. Hadinegoro SRS. The revised WHO dengue case classification: Does the system need to be modified? *Paediatr Int Child Health* 2012;32:33–8. doi: 10.1179/2046904712Z.00000000052.
2. Sandra T, Sofro MA, Suhartono S, Martini M, Hadisaputro S. Faktor yang berpengaruh terhadap kejadian demam berdarah dengue pada anak usia 6-12 tahun. *J Ilm Permas J Ilm STIKES Kendal* 2019;9:28–35. <http://dx.doi.org/10.32583/pskm.9.1.2019.28-35>.
3. Zhang H, Lui R. Releasing Wolbachia-infected *Aedes aegypti* to prevent the spread of dengue virus: a mathematical study. *Infect Dis Model* 2020;5:142–60. <https://doi.org/10.1016/j.idm.2019.12.004>.
4. Koh C, Allen SL, Herbert RI, Mcgraw EA. The transcriptional response of *Aedes aegypti* with variable extrinsic incubation periods for dengue virus. *Genome Biol Evol* 2018;10:3141–51. <https://doi.org/10.1093/gbe/evy230>.
5. Ahammad F, Tengku Abd Rashid TR, Mohamed M, Tanbin S, Fuad FAA. Contemporary strategies and current trends in designing antiviral drugs against dengue fever via targeting host-based approaches. *Microorganisms* 2019;7:296. <https://doi.org/10.3390/microorganisms7090296>.
6. Sun B, Zhang X, Zhang H, et al. Genomic epidemiological characteristics of dengue fever in Guangdong province, China from 2013 to 2017. *PLoS Negl Trop Dis* 2020;14: e0008049. <https://doi.org/10.1371/journal.pntd.0008049>.
7. Liu Y, Zhang F, Liu J, et al. Transmission-blocking antibodies against mosquito C-type lectins for dengue prevention. *PLoS Pathog* 2014;10: e1003931. <http://dx.doi.org/10.1371/journal.ppat.1003931>.
8. Londono-Renteria B, Troupin A, Conway MJ, et al. Dengue virus infection of *Aedes aegypti* requires a putative cysteine rich venom protein. *PLoS Pathog* 2015;11: e1005202. <http://dx.doi.org/10.1371/journal.ppat.1005202>.
9. Wan SW, Lin CF, Wang S, et al. Current progress in dengue vaccines. *J Biomed Sci* 2013;20:37. <https://doi.org/10.1186/1423-0127-20-37>.
10. Rahayu DF, Ustiawan A. Identifikasi *Aedes aegypti* dan *Aedes albopictus*. *Balaba J Litbang Pengendali Penyakit Bersumber Binatang Banjarnegara* 2013;9:7–10.
11. Sun P, Nie K, Zhu Y, et al. A mosquito salivary protein promotes flavivirus transmission by activation of autophagy. *Nat Commun* 2020;11: 260. <http://dx.doi.org/10.1038/s41467-019-14115-z>.
12. Guerrero D, Cantaert T, Missé D. Aedes mosquito salivary components and their effect on the immune response to Arboviruses. *Front Cell Infect*

- Microbiol 2020;10:1–11. <http://dx.doi.org/10.3389/fcimb.2020.00407>.
13. Fontaine A, Diouf I, Bakkali N, et al. Implication of haematophagous arthropod salivary proteins in host-vector interactions. *Parasites Vectors* 2011;44: 187. <http://dx.doi.org/10.1186/1471-2164-13-614>.
 14. Martin-Martin I, Smith LB, Chagas AC, et al. *Aedes albopictus* D7 salivary protein prevents host hemostasis and inflammation. *Biomolecules* 2020;10:1–17. doi: 10.3390/biom10101372.
 15. Manning JE, Morens DM, Kamhawi S, Valenzuela JG, Memoli M. Mosquito saliva: the hope for a universal arbovirus vaccine? *J Infect Dis* 2018; 218:7-15. doi: 10.1093/infdis/jiy179.
 16. Doucoure S, Cornelie S, Patramool S, et al. First screening of *Aedes albopictus* immunogenic salivary proteins. *Insect Mol Biol* 2013;22:411–23. <http://dx.doi.org/10.1111/imb.12032>.
 17. Oktarianti R, Senjarini K, Hayano T, Fatchiyah F, Aulanni'am. Proteomic analysis of immunogenic proteins from salivary glands of *Aedes aegypti*. *J Infect Public Health* 2015;8:575–82. <http://dx.doi.org/10.1016/j.jiph.2015.04.022>.
 18. Schmid MA, Kauffman E, Payne A, Harris E, Kramer LD. Preparation of mosquito salivary gland extract and intradermal inoculation of mice. *Bio Protoc* 2017;7:e2407. doi: 10.21769/BioProtoc.2407.
 19. Liu Y, Zhang F, Liu J, et al. Western Blot comparison of wet transfer and semi-dry transfer methods. *Parasites Vectors* 2011;10:54–64. <http://dx.doi.org/10.1016/j.jiph.2015.04.022>.
 20. Oktarianti R, Wathon S, Indrasari IF, Fitriani NR, Senjarini K. Platelet aggregation in vitro analysis of 67 kDa immunogenic fraction from *Aedes albopictus* salivary gland. *Bioedukasi* 2020;18:47–52. <http://dx.doi.org/10.19184/bioedu.v18i2.18891>.
 21. Wasinpiyamongkol L, Patramool S, Thongrunkiat S, et al. Protein expression in the salivary glands of dengue-infected *Aedes aegypti* mosquitoes and blood-feeding success. *Southeast Asian J Trop Med Public Health* 2012; 43:1346-57.
 22. De Carvalho IL, Rocha DK, Almeida APG. Immune reactivity to dengue and *Aedes albopictus* mosquitoes in the population from Macao, China, before dengue occurrence. *In Vivo (Brooklyn)* 2011;25:625–31.
 23. Calvo E, Mizurini DM, Sá-Nunes A, et al. Alboserpin, a factor Xa inhibitor from the mosquito vector of yellow fever, binds heparin and membrane phospholipids and exhibits antithrombotic activity. *J Biol Chem* 2011;286: 27998–8010. doi: 10.1074/jbc.M111.247924.
 24. Luplertlop N. *Aedes* mosquito salivary immune peptides: boost or block dengue viral infections. *J Coast Life Med* 2014;2:163–8. doi:10.12980/JCLM.2.2014C1219.
 25. Masoud HMM, Helmy MS, Darwish DA, Abdel-Monsef MM, Ibrahim M.A. Apyrase with anti-platelet aggregation activity from the nymph of the camel tick *Hyalomma dromedarii*. *Exp Appl Acarol* 2020;80:349–61. doi: 10.1007/s10493-020-00471-9.
 26. Juhn J, Naeem-Ullah U, MacIel Guedes BA, et al. Spatial mapping of gene expression in the salivary glands of the dengue vector mosquito, *Aedes aegypti*. *Parasites Vectors* 2011;4:1–13. <http://dx.doi.org/10.1186/1756-3305-4-1>.
 27. Peng Z, Caihe L, Beckett AN, Guan Q, James AA, Simons FER. rAed a 4: a new 67-kDa *Aedes aegypti* mosquito salivary allergen for the diagnosis of mosquito allergy. *Int Arch Allergy Immunol* 2016;170:206–10. DOI: <https://doi.org/10.1038/s41467-019-14115-z>
 28. Wichit S, Ferraris P, Choumet V, Missé D. The effects of mosquito saliva on dengue virus infectivity in humans. *Curr Opin Virol* 2016;21:139–45. doi: 10.1016/j.coviro.2016.10.001.
 29. Dong F, Fu Y, Li X, Jiang J, Sun J, Cheng X. Cloning, expression, and characterization of salivary apyrase from *Aedes albopictus*. *Parasitol Res* 2012;110:931–7. doi: 10.1007/s00436-011-2579-x.
 30. Meekins DA, Kanost MR, Michel K. Serpins in arthropod biology. *Semin Cell Dev Biol* 2017; 62:105–19. doi: 10.1016/j.semcdb.2016.09.001.
 31. Doucoure S, Mouchet F, Cornelie S, et al. Evaluation of the human IgG antibody response to *Aedes albopictus* saliva as a new specific biomarker of exposure to vector bites. *PLoS Negl Trop Dis* 2012;6:e1487. doi:10.1371/journal.pntd.0001487.
 32. Bratawijaya K, Rengganis I. Basic immunology. 11th ed. Jakarta: Publishing Agency of the Faculty of Medicine, University of Indonesia; 2014.
 33. Oktarianti R, Senjarini K, Fatchiyah F, Aulanni'am. Immunogenic protein from salivary gland of *Aedes aegypti* against to human sera. *Adv Nat Appl Sci* 2014;8:101–7.