

Series: Advances in Health Sciences Research

Proceedings of the 11th International Dentistry Scientific Meeting (IDSM 2017)

The International Dentistry Scientific Meeting 2017 (IDSM 2017) is an annual dentistry scientific event held by Faculty of Dentistry, Universitas Indonesia for dental students and dentists from all over the world. Formerly known as Dentistry Scientific Meeting in its past 9 years, DSM has broadened its scale to International level since last year as the 10th International Dentistry Scientific Meeting. This year marks the event's 11th year in providing continual seminars and courses that accelerate knowledge and skills of dental students and professionals. IDSM 2017 chose "Clinical and Research Updates in Dentistry" as the topic of the event in order to increase people's interests in dental research.

The event itself was successfully held on September 16-17th, 2017 at IMERI FKUI, Central Jakarta. In this forum, participants could take part in scientific programs such as seminars, hands on, poster presentation, and courses given by outstanding international and national speakers. There was also a dental exhibition throughout the event which was hosted by some of the industry's top manufacturers and suppliers.

There were 78 papers presented at the conference including the proceedings, which are intended to provide information to undergraduates and post-graduates students as well as general practitioners and researchers on current issues of dentistry. Each contributed paper was refereed before being accepted for publication. The double-blind peer reviewed was used in the paper selection. Then, two reviewers reviewed each of the papers. From all papers submitted, 45 papers were accepted successfully for publication based on their interest, relevance, innovation and application to the oral health sciences.

Finally, but not least, we would like to express our highest gratitude towards the

speakers, the steering committee, organizing committee and members of reviewers for their hard work and kind assistance in reviewing the papers.

The Editors

Linda Kusdhany

Izzet Yavuz

Michele Callea

Atlantis Press

A professional publisher of scientific books, journals and proceedings, Atlantis Press offers world-class services, fast turnaround and personalised communication. Our proceedings and journals are published in Open Access on our platform, generating millions of downloads each month.

For more information, please send us a message at:

contact@atlantis-press.com

Copyright © 2006-2018 Atlantis Press [Privacy Policy](#) [Terms of use](#)

▶ PROCEEDINGS

▶ JOURNALS

▶ BOOKS

▶ SEARCH

▶ PUBLISH WITH US

▶ ABOUT

▶ CONTACT

PREMIUM PAPER

Series: Advances in Health Sciences Research

Proceedings of the 11th International Dentistry Scientific Meeting (IDSM 2017)

131 authors

Achmad, Harun

The Efficacy of Chitosan Toothpaste Based White Shrimp (*Litopenaeus Vannamei*) to Decrease Streptococcus Mutans Colonies in Children with Early Childhood Caries

Agusnar, Harry

In Vitro Chitosan Hydrogel Based Tetracycline Cytotoxicity Test on Fibroblast Viability

Amalia, Martina

An in Vitro Evaluation of the Antimicrobial Effectiveness of Alginate-Based Metronidazole Gel Against *Aggregatibacter Actinomycetemcomitans*

Andrew, Andrew

In Vitro Chitosan Hydrogel Based Tetracycline Cytotoxicity Test on Fibroblast Viability

Anggono, Jessica

Congenital Missing Teeth

Anggraini, Cynthia M.

Malocclusion in Mouth-Breathing Children Caused by Nasal Obstruction

Ariella, Aida

Genetics of Crouzon Syndrome

Arina, Yuliana M.D

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Atika, Vella

Body Mass Index and Its Relation to Blood Pressure in Preclinical Students of the Faculty of Dentistry, Padjadjaran University

Auerkari, Elza Ibrahim

Molecular Genetics of Peutz-Jegher Syndrome

Auerkari, Elza Ibrahim

Genetics of Cherubism

Auerkari, Elza Ibrahim

Molecular Genetics and Epigenetics of Ankyloglossia

Auerkari, Elza Ibrahim

Genetics of Crouzon Syndrome

Auerkari, Elza Ibrahim

Effect of GHR variation on mandibular height

Auerkari, Elza Ibrahim

Congenital Missing Teeth

Auerkari, Elza Ibrahim

Ectodermal Dysplasia

Auerkari, Elza Ibrahim

The Genetics of Asthma

Auerkari, Elza Ibrahim

Genetic and Epigenetic Aspects of Crouzon's Syndrome

Bangun, Hakim

An in Vitro Evaluation of the Antimicrobial Effectiveness of Alginate-Based Metronidazole Gel Against Aggregatibacter Actinomycetemcomitans

Budiardjo, Sarworini B.

Malocclusion in Mouth-Breathing Children Caused by Nasal Obstruction

Chaira, Muhammad Abu

Dekiler-Box Sterility Test as an Innovation in Multifunction Sterilization Equipment for Dental Instruments

Chray, Mengkheng

Pilot evaluation of the therapeutic effect of Silver Diamine Fluoride (SDF) in Arresting dental caries in the primary teeth of Cambodian slum children

Da, Sreykhouch

Pilot evaluation of the therapeutic effect of Silver Diamine Fluoride (SDF) in Arresting dental caries in the primary teeth of Cambodian slum children

Dinh Phan, Nhat Dinh

Povidone-Iodine as Subgingival Irrigation in Chronic Periodontitis Treatment

Durward, Callum

Pilot evaluation of the therapeutic effect of Silver Diamine Fluoride (SDF) in Arresting dental caries in the primary teeth of Cambodian slum children

Ermawati, Tantin

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Ervina, Irma

An in Vitro Evaluation of the Antimicrobial Effectiveness of Alginate-Based Metronidazole Gel Against *Aggregatibacter Actinomycetemcomitans*

Ervina, Irma

The Effect of Scaling and Root Planing on Total Antioxidant Status of Chronic Periodontitis Patients

Ervina, Irma

In Vitro Chitosan Hydrogel Based Tetracycline Cytotoxicity Test on Fibroblast Viability

Ervina, Irma

In Vitro Chitosan Hydrogel Based Metronidazole Cytotoxicity Test on 3t3 Fibroblast Viability

Fasda, Bachuroh

Dekiler-Box Sterility Test as an Innovation in Multifunction Sterilization Equipment for Dental Instruments

Fauziah, Eva

Respiratory Rate as a Physiological Response to Dental Anxiety

Firdaus, Nikita Syahrussiami

Genetics of Cherubism

Gitasari, Nur Ismah

Body Discomfort Due to Musculoskeletal Disorders among Dental Practitioners in Primary Healthcare Center in Bandung City.

Goh, Victor

The Effect of Light-Curing Duration and Different Composite Shades on Color Stability of Composite Resins

Grafiyanti, Stevany

Radiographic Evaluation of Dental Implants in the Periodontal Clinic Rumah Sakit Khusus Gigi dan Mulut (RSKGM) by the Faculty of Dentistry, Universitas Indonesia from 2009 to 2014

Hak, Muhammad Syafrudin

Anthropometric Evaluation of Lip Symmetry following Cronin Method Labioplasty for Unilateral Complete Cleft Lip and Palate Cases

Hak, Muhammad Syafrudin

Evaluation of Upper and Lower Lip Growth following Cronin Method Labioplasty in Unilateral Complete Cleft Lip and Palate Cases

Hak, Muhammad Syafrudin

Evaluation of Nostril Symmetry Post-Cronin Labioplasty and Tajima Rhinoplasty in Unilateral Cleft Lip and Palate Cases

Halim, Arief Johanes

Effect of GHR variation on mandibular height

Hartanto, Wilson

Facial Profile Analysis by Photometry on W. R. Supratman 1 & 2 Chinese High School Students in Medan

Herawati, Dahlia

Vitamin-D and Dexamethasone Stimulation of Estrogen Receptor-?' and Androgen Receptor in Osteoblasts of Aggressive Periodontitis

Ho, Ting Khee

Patient's Self-Assessment on Orofacial Aesthetic

Ho, Ting Khee

The influence of different barometric pressure conditions on bond strength of glass fiber post cemented with resin cement

Ilyas, Syafruddin

In Vitro Chitosan Hydrogel Based Metronidazole Cytotoxicity Test on 3t3 Fibroblast Viability

Indiarti, Ike S.

Malocclusion in Mouth-Breathing Children Caused by Nasal Obstruction

Irmaleny, Irmaleny

The Acidity Level (pH) of Saliva of People in Living in High Radon Level Areas and Its Correlation with the Prevalence of Caries

Irmaleny, Irmaleny

Salivary Flow Rate in Human Population with High Radon Exposure and Its Correlation to Caries

Iskandar, Teuku Nolly

Evaluation of Nostril Symmetry Post-Cronin Labioplasty and Tajima Rhinoplasty in Unilateral Cleft Lip and Palate Cases

Jamal, Nur Syuhadah

The Effect of Light-Curing Duration and Different Composite Shades on Color Stability of Composite Resins

Katthika, Vassa Khema

Ectodermal Dysplasia

Khairani, Saufi

The Effect of Scaling and Root Planing on Total Antioxidant Status of Chronic Periodontitis Patients

Khairunnisa, Putri

The Efficacy of Chitosan Toothpaste Based White Shrimp (*Litopenaeus Vannamei*) to Decrease Streptococcus Mutans Colonies in Children with Early Childhood Caries

Kholifa, Mahmud

Comparative Effectiveness of Angsana Latex (*Pterocarpus indicus*) and

Chlorhexidine 0.2% Against Growth of Streptococcus mutans**Khorn, Shunhour****Pilot evaluation of the therapeutic effect of Silver Diamine Fluoride (SDF) in Arresting dental caries in the primary teeth of Cambodian slum children****Krisnamurthy, Krisnamurthy****Differences of Salivary Superoxide Dismutase Levels in Gingivitis and Periodontitis Patients: Periodontal Installation, Faculty Of Dentistry, Universitas Sumatera Utara****Latief, Fourier D.E****Virulence of periodontopathogens Aggregatibacter actinomycetemcomitans and Porphyromonas gingivalis: micro-computed tomography and inflammatory cytokine analysis****Ling, Goo Chui****The Effect of Light-Curing Duration and Different Composite Shades on Color Stability of Composite Resins****Loo, Chu Mian****Patient's Self-Assessment on Orofacial Aesthetic****Louisa, Marie****Esthetic Frenectomy Using Bilateral Pedicle Approach****Lubis, Hilda Fitria****Facial Profile Analysis by Photometry on W. R. Supratman 1 & 2 Chinese High School Students in Medan****Mardiana, Mardiana****The Efficacy of Chitosan Toothpaste Based White Shrimp (Litopenaeus Vannamei) to Decrease Streptococcus Mutans Colonies in Children with Early Childhood Caries****Masulili, Sri Lelyati C****Biologic Width Considerations in Crown Lengthening Surgical Procedures: A Case Report****Md Razi, Mohd Najib****The influence of different barometric pressure conditions on bond strength of**

glass fiber post cemented with resin cement

Mohd Sadek, Nurhanani

The Effect of Light-Curing Duration and Different Composite Shades on Color Stability of Composite Resins

Monica, Monica

Molecular Genetics of Peutz-Jegher Syndrome

Murdiastuti, Kwartarini

Effect of Collagen -Activated Platelet-Rich Plasma to Synthesis of Extracellular Matrix in Fibroblast of Periodontal Ligament

Murdiyanto, Dendy

Dekiler-Box Sterility Test as an Innovation in Multifunction Sterilization Equipment for Dental Instruments

Nadiah, Nisa

Salivary Flow Rate in Human Population with High Radon Exposure and Its Correlation to Caries

Nazarudin, Tifani

Dekiler-Box Sterility Test as an Innovation in Multifunction Sterilization Equipment for Dental Instruments

Nazruddin, Nazruddin

Evaluation of the Depth of the Curve of Spee, Overjet, and Overbite in Class I, Class II, and Class III Malocclusion Among Patients at University of North Sumatera Dental Hospital

Nguyen, Anh Thi Mai

Povidone-Iodine as Subgingival Irrigation in Chronic Periodontitis Treatment

Nguyen, An Thi Khanh

Evaluation and Comparison of Patient Feedback on Chlorhexidine Vs Anti-Calculus Mouthrinse After 6 Months of Compliance

Nguyen, Phuc Anh

Factors Associated with Gingivitis in 10-Year-Old School Children in Vietnam

Ong, Hsiao Ting

Patient's Self-Assessment on Orofacial Aesthetic

Permatasari, Dhea Ferrani

The Acidity Level (pH) of Saliva of People in Living in High Radon Level Areas and Its Correlation with the Prevalence of Caries

Pham, Thuy Anh Vu

Evaluation and Comparison of Patient Feedback on Chlorhexidine Vs Anti-Calculus Mouthrinse After 6 Months of Compliance

Pham, Thuy Anh Vu

Evaluation and Comparison of Patient Feedback on Chlorhexidine Vs Anti-Calculus Mouthrinse After 6 Months of Compliance

Pham, Thuy Anh Vu

Factors Associated with Gingivitis in 10-Year-Old School Children in Vietnam

Pham, Thuy Anh Vu

Correlation Between Obesity And Periodontitis In Vietnamese Patients

Phanestu, Tiffany Henry

Surface Roughness of Nanofiller Composite Resin after Immersion in Black Tea

Praharani, Depi

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Pujiastuti, Peni

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Putri, Ardhita Rosiana

Dekiler-Box Sterility Test as an Innovation in Multifunction Sterilization Equipment for Dental Instruments

Putri, Astrinia Ristia

The Genetics of Asthma

Putriyanti, Faradina

Clinical Evaluation of Dental Implants Performed in the Universitas Indonesia Dental Training Hospital Periodontal Clinic from 2009 to 2014

Ramadhany, Yunita Feby

The Efficacy of Chitosan Toothpaste Based White Shrimp (*Litopenaeus Vannamei*) to Decrease Streptococcus Mutans Colonies in Children with Early Childhood Caries

Rasni, Hilma

In Vitro Chitosan Hydrogel Based Metronidazole Cytotoxicity Test on 3t3 Fibroblast Viability

Rizali, Ervin

The Acidity Level (pH) of Saliva of People in Living in High Radon Level Areas and Its Correlation with the Prevalence of Caries

Samnieng, Patcharaphol

Preventive Oral Health Care in a School for Children with Special Health Care Needs

Sari, Desi S.

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Sari, Kartika Indah

Body Mass Index and Its Relation to Blood Pressure in Preclinical Students of the Faculty of Dentistry, Padjadjaran University

Sari, Lili Nur Indah

Molecular Genetics and Epigenetics of Ankyloglossia

Setyanto, Darmawan B.

Malocclusion in Mouth-Breathing Children Caused by Nasal Obstruction

Sinaga, Chrisnawati

Differences of Salivary Superoxide Dismutase Levels in Gingivitis and Periodontitis Patients: Periodontal Installation, Faculty Of Dentistry, Universitas Sumatera Utara

Sireerat, Kittichai

Preventive Oral Health Care in a School for Children with Special Health Care Needs

Siregar, Nasytha Vikarina

Genetic and Epigenetic Aspects of Crouzon's Syndrome

Sitam, Suhardjo

The Acidity Level (pH) of Saliva of People in Living in High Radon Level Areas and Its Correlation with the Prevalence of Caries

Soehardjo, Soehardjo

Salivary Flow Rate in Human Population with High Radon Exposure and Its Correlation to Caries

Soeroso, Yuniarti

Clinical Evaluation of Dental Implants Performed in the Universitas Indonesia Dental Training Hospital Periodontal Clinic from 2009 to 2014

1 2 >

Atlantis Press

A professional publisher of scientific books, journals and proceedings, Atlantis Press offers world-class services, fast turnaround and personalised communication. Our proceedings and journals are published in Open Access on our platform, generating millions of downloads each month.

For more information, please send us a message at:

contact@atlantis-press.com

Copyright © 2006-2018 Atlantis Press [Privacy Policy](#) [Terms of use](#)

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ SEARCH



Series: Advances in Health Sciences Research

Proceedings of the 11th International Dentistry Scientific Meeting (IDS M 2017)

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Authors

Melok A. Wahyukundari, Desi S. Sari, Peni Pujiastuti, Depi Praharani, Yuliana M.D Arina, Tantin Ermawati, Fourier D.E Latief

Corresponding author

Desi S. Sari

Keywords

Periodontal disease, crude protein of *A. actinomycetemcomitans*, *P. gingivalis* lipopolysaccharides, inflammatory cytokine, μ -CT

Abstract

The objective of this study is to determine the difference between *A. actinomycetemcomitans* and *P. gingivalis* virulence as the cause of periodontal disease. Thirty male Wistar rats were divided into six groups. Ten rats were induced by 10 μ g/mL *P. gingivalis* lipopolysaccharides (LPS), and the other ten rats were induced by 10 μ g/mL crude proteins of *A. actinomycetemcomitans*. The injection was performed three times for six weeks in the interdental area between the first and the second mandibular right molar by using a 30gauge needle. On the third and seventh days, five rats from each group were decapitated and observed with micro-computed tomography (μ -CT) to determine the severity of alveolar bone re-sorption. The immunohistochemical method was used to find the expression of macrophage inflammatory cytokines (IL-1 β and TNF- α). The ANOVA

test was performed to determine the difference among each group. The $\mu\text{-CT}$ image observation of each group revealed that the severity of alveolar bone resorption was different in each group. In the group that was induced by the crude proteins of *A. actinomycetemcomitans*, resorption was more advanced than those in the LPS and the control group. A significant increase ($p < 0.05$) was found in IL-1 β and TNF- α expressions after induction in both groups on the third and seventh days, with the highest cytokine expression observed on the seventh day. The expressions of IL-1 β and TNF- α in the group that was induced by the crude proteins of *A. actinomycetemcomitans* were significantly higher than those in the LPS group. The crude proteins of *A. actinomycetemcomitans* showed a greater level of virulence than the *P. gingivalis* LPS in either alveolar bone resorption or inflammatory cytokine expression.

⊕ [Download article \(PDF\)](#)

Part of series

[Advances in Health Sciences Research](#)

Publication date

May 2018

ISBN

978-94-6252-513-9

ISSN

2468-5739

DOI

[doi:10.2991/idsm-17.2018.42](https://doi.org/10.2991/idsm-17.2018.42) [How to use a DOI?](#)

Copyright

© The authors.

This article is distributed under the terms of the Creative Commons Attribution License 4.0, which permits non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited. See for details:

<https://creativecommons.org/licenses/by-nc/4.0/>

Atlantis Press

A professional publisher of scientific books, journals and proceedings, Atlantis Press offers world-class services, fast turnaround and personalised communication. Our proceedings and journals are published in Open Access on our platform, generating millions of downloads each month.

For more information, please send us a message at:

contact@atlantis-press.com

Copyright © 2006-2018 Atlantis Press [Privacy Policy](#) [Terms of use](#)

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ SEARCH

- ▶ PUBLISH WITH US
- ▶ ABOUT
- ▶ CONTACT

PREMIUM P...

Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis

Melok A. Wahyukundari¹, Desi S. Sari^{1*}, Peni Pujiastuti¹, Depi Praharani¹,
Yuliana M.D Arina¹, Tantin Ermawati², Fourier D.E. Latief³

¹Departement of Periodontology, Faculty of Dentistry, University of Jember, Jember, Indonesia

²Departement of Biomedic, Faculty of Dentistry, University of Jember, Jember, Indonesia

³Physics of Earth and Complex Systems, Faculty of Mathematics and Natural Sciences, Bandung
Institute of Technology, Bandung, Indonesia

*Email: desi_sari.fkg@unej.ac.id

Abstract. The objective of this study is to determine the difference between *A. actinomycetemcomitans* and *P. gingivalis* virulence as the cause of periodontal disease. Thirty male Wistar rats were divided into six groups. Ten rats were induced by 10 ug/mL *P. gingivalis* lipopolysaccharides (LPS), and the other ten rats were induced by 10 ug/mL crude proteins of *A. actinomycetemcomitans*. The injection was performed three times for six weeks in the interdental area between the first and the second mandibular right molar by using a 30gauge needle. On the third and seventh days, five rats from each group were decapitated and observed with micro-computed tomography (μ -CT) to determine the severity of alveolar bone resorption. The immunohistochemical method was used to find the expression of macrophage inflammatory cytokines (IL-1 β and TNF- α). The ANOVA test was performed to determine the difference among each group. The μ -CT image observation of each group revealed that the severity of alveolar bone resorption was different in each group. In the group that was induced by the crude proteins of *A. actinomycetemcomitans*, resorption was more advanced than those in the LPS and the control group. A significant increase ($p < 0.05$) was found in IL-1 β and TNF- α expressions after induction in both groups on the third and seventh days, with the highest cytokine expression observed on the seventh day. The expressions of IL-1 β and TNF- α in the group that was induced by the crude proteins of *A. actinomycetemcomitans* were significantly higher than those in the LPS group. The crude proteins of *A. actinomycetemcomitans* showed a greater level of virulence than the *P. gingivalis* LPS in either alveolar bone resorption or inflammatory cytokine expression.

Keywords: Periodontal disease, crude protein of *A. actinomycetemcomitans*, *P. gingivalis* lipopolysaccharides, inflammatory cytokine, μ -CT

1 Introduction

Periodontal disease is a chronic infectious disease with a multi-bacterial factor, and it is characterized by the destruction of periodontal ligaments and alveolar bone. The severity of periodontal disease has been established to be dependent on the

interactions between the microbial challenge and the host immune inflammatory responses. Some gram-negative bacteria, such as *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*, have been consistently detected as normal constituents of the microbial flora in periodontal disease. *P. gingivalis* and *A. actinomycetemcomitans* are important periodontal pathogens [1,2]. All of the periodontopathogenic bacteria have virulence factors, such as lipopolysaccharides (LPS) and crude proteins, which can stimulate inflammatory cytokine expression [3]. LPS have been primarily shown to stimulate monocytes to produce cytokines, such as TNF α and IL-1. The balance between the local levels of cytokines stimulated in response to periodontopathogenic bacteria and their products is important in determining the outcome of an immune response to a given pathogen [4]. IL-1 is a highly pro-inflammatory cytokine. IL-1 is a multifactorial cytokine that can activate many cell types with strong inflammatory features. The IL-1 release is typically observed to occur shortly after LPS stimulation. IL-1 affects nearly all cell types and usually in concert with other cytokines or mediator molecules. The broad biological effects of IL-1 result from its central role in the regulation of many different genes that regulate cytokines, cytokine receptors, acute phase reactants, growth factors, extracellular matrix components, and adhesion molecules [5,6]. Clinical studies found that the amount of IL-1 β is much higher in the gingival crevicular fluid (GCF) in periodontal pockets or in the underlying inflamed gingival tissue than in the healthy sites. Furthermore, IL-1 β levels are higher in active periodontitis sites than in stable sites [7].

TNF- α is released by monocytes and macrophages in a rapid burst after LPS stimulation. TNF- α induces bone and extracellular matrix resorption by activating the osteoclasts. The local cellular effects of TNF- α include the adhesion of polymorph nuclear leukocytes (PMNs) to endothelial cells, degranulation of PMNs, activation of phagocytosis, and intercellular adhesion molecule (ICAM) -1 expression. The amount of TNF- α was demonstrated to be at a high level in the gingival crevicular fluid and diseased periodontal tissues. Furthermore, experimental studies have shown a central role for TNF- α in alveolar bone resorption [6,8]. The current study aims to determine the difference between *A. actinomycetemcomitans* and *P. gingivalis* virulence as periodontopathogenic bacteria in periodontal disease.

2 Material and Methods

This experiment was conducted at the Microbiology and Biomedical Laboratory of the Faculty of Dentistry at University of Jember and the Faculty of Medicine at Brawijaya University, the Bioscience Laboratory of Oral and Dental Hospital at University of Jember, and the Faculty of Physics of Earth and Complex Systems of the Faculty of Mathematics and Natural Sciences, Bandung Institute of Technology.

2.1 Preparation of *P. gingivalis* LPS

One milligram of *P. gingivalis* LPS (Invivo Gen, San Diego, USA) was dissolved in 2 mL Dulbecco's phosphate-buffered saline, homogenized, and stored at 4 °C before usage.

2.1 Culture and Preparation of Crude Proteins of *A. actinomycetemcomitans* Toolbars

A *actinomycetemcomitans* Y4 (serotype b) strains were grown in brain-heart infusion media with the addition of 1% yeast extract at 37 °C with 10% level of CO₂ using a CO₂ generating GasPack sachet for 7-10 days [9]. A *actinomycetemcomitans* of the local isolate/serotype b was isolated using the method of Westphal and Jann. The crude proteins were isolated using the phenol method. About 500 ml of bacterial suspension was centrifuged at a rotation of 15,000 g (6000 rpm) for 1 h. The pellets obtained were resuspended in 10 ml deionized water, which was heated at 72 °C-75 °C. The suspension was then vortexed for 20 s and placed in a water bath at a temperature of 72 °C-75 °C with the mouth of test tube opened. Subsequently, 10 ml of the 88% phenol solution (previously heated at 72 °C-75 °C) was added, vortexed, and incubated for 15 min in a water bath [10].

3 Experimental Model of Periodontitis

Thirty male *Rattus norvegicus* of Wistar strains aged 12-14 weeks and weighing 200-250 g were used in this study. This research was approved by the Ethics and Advocacy Unit of the Faculty of Dentistry, Gadjah Mada University (No.00748 / KKEP / FKG-UGM / EC / 2016). Anesthesia was performed before induction using a combination of intra-muscular ketamine and xylazine. Chronic periodontitis conditioning was conducted by the induction of the *P. gingivalis* LPS and the crude proteins of *A. actinomycetemcomitans*. These substances were injected at the interproximal gingiva between the first and second molars of the buccal aspect of a mandible. Each experimental animal was given *P. gingivalis* LPS and crude proteins of *A. actinomycetemcomitans*, with a 10 µl volume and 0.5 mg/ml concentration, with a 1 cc/ml tuberculin syringe (Terumo) with a 30 G (BD) needle administered three times a week for six weeks [11,12]. The rats were divided into six groups, in which five rats were induced by the crude proteins of *A. actinomycetemcomitans* and decapitated on the third day post-injection and another five rats on the seventh day post-injection. The other five rats were induced by LPS from *P. gingivalis* and decapitated on the third day following the injection, and another five were decapitated on the seventh day post-injection. The control group, which consisted of five rats, was decapitated on the third and seventh days after six weeks.

3.1. Micro-Computed Tomography (µ-CT)

The microstructural analysis of the mandibles was conducted using the X-ray µ-CT imaging technique. Three samples were scanned: six weeks of the control sample, six weeks of the induced crude proteins of the *A. actinomycetemcomitans* sample, and six weeks of the induced *P. gingivalis* LPS sample. The samples were scanned using the Bruker SkyScan 1173 High-energy X-ray µ-CT (Bruker Micro-CT, Kostich, Belgium) using the primary parameters listed in Table 1. The samples were scanned using different settings. However, the adjustment of the parameters was made on the basis of the primary acquisition standard that the intensity of the detected X-ray should be in the min-max range of 30%-90%.

Table 1. Scanning parameters for the three scanned samples

Parameters	Control	Crude protein of <i>A. Actinomycetemcomitans</i>	<i>P. gingivalis</i> LPS
Source Voltage (kV)	60	60	80
Source Current (uA)	100	50	50
Filter	Al 1.0 mm	Al 1.0 mm	-
Exposure (ms)	700	125	125
Rotation Step (deg)	0.2	0.2	0.2
Image Pixel Size (µm)	12.114213	24.231851	24.231851
Camera Binning	1x1 (2240 x 2240 pix)	4x4 (560 x 560 pix)	4x4 (560 x 560)

Following the scanning process, reconstruction of the projection images was conducted using the NRecon software (using the GPUReconServer kernel). Subsequently, 3D visual analyses were conducted using the CTVox software. Evaluation of the resorption was performed by measuring the distance between the cemento-enamel junction (CEJ) and the processus alveolaris. The measurement can be taken using DataViewer or CTAn software [13].

3.2. Decapitation and Immunohistochemical (IHC) Assay

The animal subjects were asphyxiated using either inhalation on the third and seventh days postinjection. Subsequently, the right mandibular bone of the first and second molars was soaked in a 10% buffered formalin fixation solution for at least 8 h before decalcification. The specimen was immersed in an EDTA solution. After the decalcification process was completed, the sample for IHC analysis was examined to investigate the expression of IL-1 and TNF- α . The expression of TNF- α was disclosed in the rodent mandibular tissue with the TNF- α monoclonal antibody (LS Bio, USA). The method of TNF- α expression painting was performed according to the UltraTek HRP Anti-Polyvalent (DAB) Staining System (Scytek, USA). The IL- β expression was detected in the rodent mandibular tissue using the TNF- α monoclonal antibody (LS Bio, USA). The method of IL1 β expression painting was based on the UltraTek HRP Anti-Polyvalent (DAB) Staining System (Scytek, USA). The expressions of TNF- α and IL1 β were observed under an Olympus CX-31 light microscope at 400x magnification [11,14].

3.3. Statistical analysis

Results are presented as the mean and the standard deviation (SD). The ANOVA test was performed to determine the significance of the results between the selected groups at a significance level of 95%. The significant difference was analyzed by T-test.

4 Results

Reconstruction of the projection images produced a set of 8-bit grayscale images, which are 2D images of the structure in the transaxial slices. The grayscale images contain information on the intensity level (grey level) of the structure of the sample, which can be commonly related to the density of the sample's structure. These images can be visualized in 3D using CTVox (see Fig. 1). For the sample that was induced by the crude proteins of *A. actinomycetemcomitans* for six weeks, the resorption was so severe (see the bottom inset of Figure 1.B) that the measurement of the CEJ-processus alveolaris distance was difficult to conduct using the grey scale images. Thus, we used the maximum intensity projection (MIP) image of the sample to produce the one shown in Figure 1.B. The result of the measurement of the CEJ-processus alveolaris distance is presented in Table 2.




Fig 1. (A) Description of the μ -CT of a six-week control rat. The arrows indicate the CEJ and the processus alveolaris. (B) Description of the μ -CT of a rat that was induced by the crude proteins of *A. actinomycetemcomitans* for six weeks. (C) Description of the μ -CT of a rat that was induced by *P. gingivalis* LPS for six weeks

Table 2. Results of the resorption measurement of the alveolar bone of the Wistar rats. The sequence of the measurements of the samples in Figure 1 begins from left to right.

Measurement point	1	2	3	4	5	6	7
CEJ-processus alveolaris distance of the control rat (μm)	420	576	370	525	447	343	293
CEJ-processus alveolaris distance of the crude proteins of <i>A. actinomycetemcomitans</i> (μm)	2400	2472	2160	2040	2088	1440	1560
CEJ-processus alveolaris length of <i>P. gingivalis</i> LPS (μm)	1503	1285	1042	1163	1018	800	969

The μ -CT analysis showed that alveolar bone resorption occurred in the rats that were induced by *P. gingivalis* LPS and the crude protein of *A. actinomycetemcomitans* for six weeks. The resorption indicated a decrease in the apical margin of the gingiva in the molar rat tooth area compared with the control sample.

4.1. TNF- α Expression Analysis Using IHC Methods

Fig 2 shows that TNF- α expression in tissue with crude proteins of *A. actinomycetemcomitans* and *P. gingivalis* LPS did not affect the TNF- α expression in the control group on either the third day or the seventh day. By contrast, for the treatment group on the 7th day, the injection of the crude proteins of *A. actinomycetemcomitans* and *P. gingivalis* LPS increased the TNF- α expression unlike on the third

day. The result of the statistical analysis concluded a significant increase in the TNF- α expression on the seventh day post-injection of the crude proteins of *A. actinomycetemcomitans*. *P. gingivalis* LPS and the control group exhibited no significant differences. The analysis was using IHC Methods (Fig 3-5)




Fig. 2. Bar chart analysis showing a significant increase in the TNF- α expression on the seventh day compared with that on the third day and post-injection control ($p=0.02$) six weeks after the induction by the crude proteins of *A. actinomycetemcomitans*. (*: Significant, NS: not significant)




Fig. 3. TNF- α expression on macrophages in the periodontal ligament tissues using IHC methods in the control group and the induction group of crude proteins of *A. actinomycetemcomitans*. (A) Control group on the third day post-injection. (B) Control group on the seventh day post-injection. (C) Group treated with the crude proteins of *A. actinomycetemcomitans* on the third day post-injection. (D) Group treated with the crude proteins of *A. actinomycetemcomitans* on the seventh day post-injection. (E) Group treated with *P. gingivalis* LPS on the third day post-injection. (F) Group treated with *P. gingivalis* LPS on the seventh day post-injection (400x magnification).

4.2. IL-1 β analysis using IHC methods




Fig. 4. Bar chart analysis showing a significant increase in the IL-1 β expression on the seventh day compared with that on the third day and post-injection control ($p=0.01$) six weeks after the induction by the crude proteins of *A. actinomycetemcomitans*. (*: Significant, NS: non-significant)




Fig. 5. IL-1 β expression on macrophages in periodontal ligament tissues observed using IHC methods in the control group and the group treated with the crude proteins of *A. actinomycetemcomitans*. (A) Control group on the third day post-injection. (B) Control group on the seventh day post-injection. (C) Group treated with the crude proteins of *A. actinomycetemcomitans* on the third day post-injection. (D) Group treated with the crude proteins of *A. actinomycetemcomitans* on the seventh day post-injection. (E) Group treated with *P. gingivalis* LPS on the third day post-injection. (F) Group treated with *P. gingivalis* LPS on the seventh day post-injection (400x magnification).

4 Discussion

In the present study, we established a new model of periodontitis in rats through the induction of the crude proteins of *A. actinomycetemcomitans* and *P. gingivalis* LPS. In six weeks, evidence of periodontal disease, including increased inflammatory cell infiltration, enhanced inflammatory cytokine expression, and stimulated osteoclastogenesis, indicated a significant alveolar bone loss. Based on the results, the virulence level of *A. actinomycetemcomitans* in the expression of increased inflammatory cytokines was higher than that of *P. gingivalis*. *A. actinomycetemcomitans* produces collagenase that can destroy type 1 collagen, thus causing the collagen degradation in periodontal tissue. These bacteria also have several other virulence factors that accelerate periodontal disease progression. Virulence affects the intensity and potential of the pathogen in host destruction as well as their defense mechanisms toward the host immune system [11,15].

The crude proteins from *A. actinomycetemcomitans* are a type of protein that can be found inside the bacteria cell, both in endotoxin and in exotoxin form. *A. actinomycetemcomitans* produces two exotoxins, namely, leukotoxin and cytolethal distending toxin. These toxins can cause cell death, inhibit growth, and stimulate an inflammatory response such as bone resorption. Target cells with receptors involved in leukotoxin induction experience cell lysis, which is related to the function of receptor-related lymphocytes 1 (LFA-1) [16,17].

P. gingivalis LPS, which are important pathogenic bacteria, are closely associated with the inflammatory destruction of periodontal tissues. They induce the release of cytokines and local factors from inflammatory cells, stimulate osteoclastic cell differentiation, and cause alveolar bone resorption. *P. gingivalis* LPS stimulate the expression of inflammatory cytokines, as inflammatory cytokines such IL-1 and IL-6 are involved in the pathogenesis of periodontal disease [18].

A study Roger et al, on a rat periodontitis model with *Escherichia coli* endotoxin or using a wire ligature-induced method showed 50% more severe bone destruction on the 7-14 days after induction. Conversely, eight weeks after the induction of the LPS of *A. actinomycetemcomitans*, the alveolar bone destruction was even more severe than that of the induction of the LPS of *E. coli* induction. Proinflammatory cytokine production occurs in the pathway of toll-like receptors (TLRs), which have been categorized as a family of innate immune recognition receptors that detect conserved microbial patterns and endogenous ligands. These receptors, along with CD14-expressing immune cells, play a fundamental role in innate immune signaling and lead to inflammatory responses by activating several transcription factors, including the nuclear factor- κ B (NF- κ B) [11].

LPS, proinflammatory glycolipids and a main component of the cell wall of all Gram-negative bacteria, are shown to be capable of inducing bone resorption in vitro. LPS initiate signal transduction through the toll-like receptor (TLR)-4, which causes the activation of macrophages, lymphocytes, and endothelial cells to release proinflammatory cytokines, such as TNF- α , IL-1 β , and IL-6. These proinflammatory cytokines can induce osteoclast formation and activation. Moreover, LPS can promote osteoclast differentiation, fusion, survival, and activation independence of IL-1, TNF- α , and RANKL. In these cytokines, TNF- α has been reported to be upstream of other cytokines and plays a central role in inflammatory reaction. TNF- α

exerts its biological effects by binding to the TNF type 1 receptor (TNFR1) and type 2 receptor. These receptors are expressed in the lineages of both osteoclasts and osteoblasts. TNFR1 has already been shown to be critical for LPS-induced osteoclastogenesis in vivo. 19,20 LPS promote the survival of differentiated osteoclasts in vitro in the presence of OPG or IL-1 receptor antagonists, or in cells derived from TNF- α receptor-deficient mice, through the activation of both the NF- κ B and P13K pathways. Therefore, LPS can promote osteoclast survival both indirectly through the stimulation of cytokines and directly through the activation of survival pathways [21].

TNF- α and/or IL-1 β and/or IL-6 levels in blood were found to be high in rats with ligature-induced periodontitis. In the LPS rat periodontitis model, the increase in the TNF- α levels in blood was also reported, although this model could be difficult to interpret because LPS from different Gram-negative bacteria could elicit different host responses. Proinflammatory cytokines expression, primarily T helper 1-type TNF- α , was significantly elevated on the first 30 days post-infection, but the level decreased in the last 30 days post-infection [22].

6 Conclusions

Crude proteins of *A. actinomycetemcomitans* showed a higher level of virulence than *P. gingivalis* LPS in either alveolar bone resorption or inflammatory cytokine expression.

References

1. Takasaki K, Fujise O, Miura M, Hamachi T, Maeda K. *Porphyromonas gingivalis* displays a competitive advantage over *Aggregatibacter actinomycetemcomitans* in co-cultured biofilm. *J Periodont Res*. 2013;48(3):286-92.
2. Carranza FA. Bone loss and patterns of bone destruction. *Carranza's clinical periodontology*, 2002. 2002.
3. Grover HS, Saini R, Bhardwaj P, Bhardwaj A. Cytokines and Other Inflammatory Mediators in Periodontal Health and Disease. *Indian J Oral Health Res*. 2016;2(1):12.
4. Kulkarni C, Kinane DF. Host response in aggressive periodontitis. *Periodontol 2000*. 2014;65(1):79-91.
5. Yücel ÖÖ. Inflammatory cytokines and the pathogenesis of periodontal disease. *Immun Res*. 2015;11(2):1.
6. Cekici A, Kantarci A, Hasturk H, Van Dyke TE. Inflammatory and immune pathways in the pathogenesis of periodontal disease. *Periodontol 2000*. 2014;64(1):57-80.
7. How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: an overview of periodontopathic pathogen below the gum line. *Frontiers Microbio*. 2016; 7:53.
8. Vahabi S, Sattari M, Taheraslani M, Bagheban AA. Correlation between Interleukin-1 β , Interleukin-6 and Tumor Necrosis Factor- α and Clinical Parameters in Chronic and Aggressive Periodontal Disease. *J Periodontol Implant Dent*. 2011; 3:51-6.
9. Izawa A, Ishihara Y, Mizutani H, Kobayashi S, Goto H, Okabe E, Takeda H, Ozawa Y, Kamiya Y, Sugita Y, Kubo K. Inflammatory bone loss in experimental periodontitis induced by *Aggregatibacter actinomycetemcomitans* in interleukin-1 receptor antagonist knockout mice. *Infection and immunity*. 2014 May 1;82(5):1904-13.
10. Westerman RB, He Y, Keen JE, Littledike ET, Kwang J. Production and characterization of monoclonal antibodies specific for the lipopolysaccharide of *Escherichia coli* O157. *J clin microbio*. 1997;35(3):679-84.

11. Rogers JE, Li F, Coatney DD, Rossa Jr C, Bronson P, Krieder JM, Giannobile WV, Kirkwood KL. Actinobacillus actinomycetemcomitans lipopolysaccharide-mediated experimental bone loss model for aggressive periodontitis. *J periodontol.* 2007;78(3):550-8.
12. Jin J, Zhang X, Lu Z, Li Y, Lopes - Virella MF, Yu H, Haycraft CJ, Li Q, Kirkwood KL, Huang Y. Simvastatin inhibits lipopolysaccharide - induced osteoclastogenesis and reduces alveolar bone loss in experimental periodontal disease. *J periodontal res.* 2014;49(4):518-26.
13. Latief FD, Sari DS, Fitri LA. Applications of Micro-CT scanning in medicine and dentistry: Microstructural analyses of a Wistar Rat mandible and a urinary tract stone. In *Journal of Physics: Conference Series* 2017;884(1) 012042.
14. Hendrijantini N, Kresnoadi U, Salim S, Agustono B, Retnowati E, Syahrial I, Mula-wardhana P, Wardhana MP, Pramono C, Rantam FA. Study biocompatibility and osteogenic differentiation potential of human umbilical cord mesenchymal stem cells (hUCMSCs) with gelatin solvent. *J Biomed Sci Engineering.* 2015;8(07):420.
15. Yücel ÖÖ. Inflammatory cytokines and the pathogenesis of periodontal disease. *Immunome Res.* 2015;11(2):1.
15. Raja M, Fajar Ummer CP. Aggregatibacter actinomycetemcomitans—A tooth killer? *Journal of clinical and diagnostic research: JCDR.* 2014;8(8): ZE13.
16. Francis PM, Manickam R, Venkatesan BP. Crude protein extract of Actinobacteria exhibits antibacterial activity against Salmonella typhi. *Int J Curr Microbiol App Sci.* 2014;3(7):319-26.
17. Bostanci N, Belibasakis GN. Porphyromonas gingivalis: an invasive and evasive opportunistic oral pathogen. *FEMS Microbiol Letters.* 2012;333(1):1-9.
18. Guo C, Hou GQ, Li XD, Xia X, Liu DX, Huang DY, Du SX. Quercetin triggers apoptosis of lipopolysaccharide (LPS)-induced osteoclasts and inhibits bone resorption in RAW264. 7 cells. *Cellular Physiol Biochem.* 2012;30(1):123-36.
19. Hou GQ, Guo C, Song GH, Fang N, Fan WJ, Chen XD, Yuan L, Wang ZQ. Lipopolysaccharide (LPS) promotes osteoclast differentiation and activation by enhancing the MAPK Pathway And COX-2 Expression In RAW264. 7 Cells. *Int j molecular med.* 2013 Aug 1;32(2):503-10.
20. Bilezikian JP, Raisz LG, Martin TJ, editors. *Principles of bone biology.* Academic Press; 2008 Sep 29.
21. Nemeč A, Petelin M, Pavlica Z. Systemic inflammatory host response to periodontopathogenic bacteria in the oral cavity: from experimental to clinical studies. *Microbial pathogens and strategies for combating them: science, technology and education.* 2013:1778-86.