

IMPACT FACTOR 4.927

Indexed in: PubMed

A Composite Electrolyte of Novel in Situ Growth of ZIF-8 in Porous Epoxy Matrix

Volume 27 · Issue 21 | November (I) 2022



mdpi.com/journal/molecules ISSN 1420-3049

Editorial Board

Prof. Dr. Farid Chemat grade Editor-in-Chief Université d'Avignon et des Pays du Vaucluse, 84029 Avignon, France

Dr. Sylvain Caillol Section Editor-in-Chief Institut Charles Gerhardt Montpellier (ICGM), University of Montpellier, CNRS, ENSCM, 34095

Dr. Giuseppe Cirillo Section Editor-in-Chief Department of Pharmacy, Health and Nutritional Sciences, University of Calabria, 87036 Rende, Italy

Prof. Dr. Roman Dembinski Section Editor-in-Chief Department of Chemistry, Oakland University, 146 Library Drive, Rochester, MI 48309-4479, USA

Prof. Dr. Mara G. Freire Section Editor-in-Chief CICECO - Aveiro Institute of Materials, Chemistry Department, University of Aveiro, Aveiro, Portugal

Prof. Dr. James Gauld Section Editor-in-Chief Department of Chemistry and Biochemistry, University of Windsor, Windsor, ON N9B 3P4, Canada

Prof. Dr. Henryk H. Jeleń
Section Editor-in-Chief
Faculty of Food Science and Nutrition, Poznań University of Life Sciences, Wojska Polskiego 31, 60-624 Poznań, Poland

Prof. Dr. Ashok Kakkar

Section Editor-in-Chief

Department of Chemistry, McGill University, 801 Sherbrooke St. West, Montreal, QC H3A 0B8, Canada

Prof. Dr. Axel Klein Section Editor-in-Chief Department für Chemie, Institut für Anorganische Chemie, Universität zu Köln, Greinstraße 6, 50939 Köln, Germany

Prof. Dr. R. Daniel Little Section Editor-in-Chief Department of Chemistry and Biochemistry, University of California, Santa Barbara, CA 93106, USA

Prof. Dr. Mauro Maccarrone Section Editor-in-Chief 1. Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, L'Aquila, Italy

Prof. Dr. M. Gilles Mailhot

Section Editor-in-Chief

Institut de Chimie de Clermont-Ferrand, Université Clermont Auvergne - CNRS, F-63000 Clermont-Ferrand, France

 Dr. Diego Muñoz-Torrero
 Section Editor-in-Chief
 Laboratory of Medicinal Chemistry, Faculty of Pharmacy and Food Sciences, Institute of Biomedicine (IBUB), University of Barcelona, Av. Joan XXIII, 27-31, E-08028 Barcelona, Spain

Prof. Dr. Lucia Panzella

Section Editor-in-Chief

Department of Chemical Sciences, University of Naples "Federico II", Via Cintia 4, I-80126 Naples, Italy

Dr. Joselito P. Quirino

Section Editor-in-Chief

Australian Centre of Research on Separation Science, School of Physical Science, University of Tasmania, Hobart, Tasmania, Australia

Prof. Dr. Thomas J. Schmidt

Section Editor-in-Chief

Institute of Pharmaceutical Biology and Phytochemistry, University of Münster, Corrensstrasse 48, D-48149 Münster, Germany

Prof. Dr. Vadim A. Soloshonok Section Editor-in-Chief Department of Organic Chemistry I, University of Basque Country, San Sebastian, Spain

Prof. Dr. Michal Szostak Section Editor-in-Chief Department of Chemistry, Rutgers University, 73 Warren St, Newark, NJ 07102, USA

Prof. Dr. Mark von Itzstein Section Editor-in-Chief Institute for Glycomics, Gold Coast Campus, Griffith University, Gold Coast, QLD 4222, Australia

Prof. Dr. Anan Yaghmur Section Editor-in-Chief Department of Pharmacy, Kobenhavns Universitet, Copenhagen, Denmark

Prof. Dr. Fernando Albericio Associate Editor IRB Barcelona - Institute for Research in Biomedicine, Barcelona, Spain

Prof. Dr. Maurizio Battino grade Associate Editor Department of Odontostomatologic and Specialized Clinical Sciences, Sez-Biochimica, Faculty of Medicine, Università Politecnica delle Marche, Via Ranieri 65, 60100 Ancona, Italy

Dr. John A. Beutler

Associate Editor

Molecular Targets Laboratory, Bldg 560-1, Room 11-86, Frederick National Laboratory for Cancer Research, Frederick, MD 21702-1201, USA

Prof. Dr. Yi Tang

Associate Editor

Department of Chemistry, Shanghai Key Laboratory of Molecular Catalysis and Innovative Materials, Laboratory of Advanced Materials and Collaborative Innovation Center of Chemistry for Energy Materials, Fudan University, Shanghai 200433, China

Prof. Dr. Curt Wentrup

Associate Editor

School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia

Prof. Dr. Santiago Aparicio Associate Section Editor-in-Chief Department of Chemistry, University of Burgos, 09001 Burgos, Spain

Prof. Dr. Nigel D. Browning

Associate Section Editor-in-Chief

Pacific Northwest National Laboratory, School of Engineering & School of Physical Sciences, University of Liverpool, Liverpool L69 3GQ, UK

Prof. Dr. Santiago Garcia-Granda Associate Section Editor-in-Chief

Faculty of Chemistry, University of Oviedo, 33006 Oviedo, Spain

Prof. Dr. Ivan Gitsov

Associate Section Editor-in-Chief The Michael M. Szwarc Polymer Research Institute, State University of New York – ESF, Syracuse, NY 13210, USA

Prof. Dr. Eun Kyoung Seo Associate Section Editor-in-Chief College of Pharmacy, Ewha Womans University, Seoul 03760, Korea

Dr. Pierre Pichat Section Editor-in-Chief Photocatalyse et Environnement, CNRS/ Ecole Centrale de Lyon (STMS), 69134 Ecully CEDEX, France

Prof. Dr. M.-H. Whangbo Section Editor-in-Chief

1. Department of Chemistry, North Carolina State University, Raleigh, NC 27695-8204, USA

2. State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter (FJIRSM), Chinese Academy of Sciences (CAS), Fuzhou 350002, China

3. State key Laboratory of Crystal Materials, Shandong University, Jinan 250100, China

Prof. Dr. Roland J. Pieters
Section Editor-in-Chief
Department of Chemical Biology & Drug Discovery, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, PO Box 80082, Utrecht, The Netherlands

Prof. Dr. Rudy J. Richardson
Section Editor-in-Chief
Molecular Simulations Laboratory, EHS/Toxicology, University of Michigan, 1415 Washington Heights, Ann Arbor, Michigan 48109-2029 USA

Prof. Dr. Daniela Rigano Associate Editor Department of Pharmacy, University of Naples "Federico II", Via D. Montesano, 49, 80131 Naples, Italy

Dr. Jean Jacques Vanden Eynde Associate Editor Formerly head of the Department of Organic Chemistry (FS), University of Mons-UMONS, 7000 Mons, Belgium

Contents

- Phenolic Compounds as Phytochemical Tracers of Varietal Origin of Some Autochthonous Apple Cultivars Grown in Serbia. by Nikola Horvacki, Filip Andrić, Uroš Gašić, Dejan Đurović, Živoslav Tešić, Milica Fotirić Akšić and Dušanka Milojković-Opsenica. Molecules 2022, 27(21), 7651; https://doi.org/10.3390/molecules27217651 - 07 Nov 2022
- Efficient Synthesis of Fluorescent Coumarins and Phosphorous-Containing Coumarin-Type Heterocycles via Palladium Catalyzed Cross-Coupling Reactions. by Rumen Lyapchev, Ana I. Koleva, Iskra Z. Koleva, Kristian Subev, Ivelina Madzharova, Kristina B. Simeonova, Nevena Petkova-Yankova, Bernd Morgenstern, Vesela Lozanova, Petar Y. Petrov andRositca D. Nikolova. Molecules 2022, 27(21), 7649; https://doi.org/10.3390/molecules27217649 - 07 Nov 2022
- Isolation of a New Polysaccharide from Dandelion Leaves and Evaluation of Its Antioxidant, Antibacterial, and Anticancer Activities. by Mo Li, Henan Zhang, Xinyu Hu, Yumeng Liu, Yanfeng Liu, Meijun Song, Rina Wu andJunrui Wu. Molecules 2022, 27(21), 7641; https://doi.org/10.3390/molecules27217641 - 07 Nov 2022
- Biodereplication of Antiplasmodial Extracts: Application of the Amazonian Medicinal Plant Piper coruscans Kunth. by Pedro G. Vásquez-Ocmín, Jean-François Gallard, Anne-Cécile Van Baelen, Karine Leblanc, Sandrine Cojean, Elisabeth Mouray, Philippe Grellier, Carlos A. Amasifuén Guerra, Mehdi A. Beniddir, Laurent Evanno, Bruno Figadère and Alexandre Maciuk. Molecules 2022, 27(21), 7638; https://doi.org/10.3390/molecules27217638 - 07 Nov 2022
- Dietary Polyphenols as Prospective Natural-Compound Depression Treatment from the Perspective of Intestinal Microbiota Regulation. by Xuanpeng Wang, Jing Yu and Xin Zhang. Molecules 2022, 27(21), 7637; https://doi.org/10.3390/molecules27217637 07 Nov 2022
- Flavonoids from Sedum japonicum subsp. oryzifolium (Crassulaceae). by Takayuki Mizuno, Nahoko Uchiyama, Seiji Tanaka, Takahisa Nakane, Hari Prasad Devkota, Kazumi Fujikawa, Nobuo Kawahara and Tsukasa Iwashina. Molecules 2022, 27(21), 7632; https://doi.org/10.3390/molecules27217632 - 07 Nov 2022
- Amino-Acid-Conjugated Natural Compounds: Aims, Designs and Results. by Hanggara Arifian, Rani Maharani, Sandra Megantara, Amirah Mohd Gazzali and Muchtaridi Muchtaridi. Molecules 2022, 27(21), 7631; https://doi.org/10.3390/molecules27217631 07 Nov 2022
- Antibacterial Effect of Cell-Free Supernatant from Lactobacillus pentosus L-36 against Staphylococcus aureus from Bovine Mastitis. by Gengchen Wang and Hong Zeng Molecules 2022, 27(21), 7627; https://doi.org/10.3390/molecules27217627 - 07 Nov 2022
- Quercetin's Effects on Glutamate Cytotoxicity. by Kade Riche and Natalie R. Lenard. Molecules 2022, 27(21), 7620; https://doi.org/10.3390/molecules27217620 07 Nov 2022
- Assessment of Cucurbita spp. Peel Extracts as Potential Sources of Active Substances for Skin Care and Dermatology. by Katarzyna Gaweł-Bęben, Karolina Czech, Marcelina Strzępek-Gomółka, Marcin Czop, Monika Szczepanik, Anna Lichtarska and Wirginia Kukula-Koch. Molecules 2022, 27(21), 7618; https://doi.org/10.3390/molecules27217618 - 06 Nov 2022
- Ultrasound-Assisted Water Extraction of Gentiopicroside, Isogentisin, and Polyphenols from Willow Gentian "Dust" Supported by Hydroxypropyl-β-Cyclodextrin as Cage Molecules. by Miloš S. Jovanović, Nemanja Krgović, Katarina Šavikin and Jelena Živković. Molecules 2022, 27(21), 7606; https://doi.org/10.3390/molecules27217606 - 06 Nov 2022
- Synthesis, Structure and Molecular Docking of New 4,5-Dihydrothiazole Derivatives Based on 3,5-Dimethylpyrazole and Cytisine and Salsoline Alkaloids. by Marat K. Ibrayev, Oralgazy A. Nurkenov, Zhanar B. Rakhimberlinova, Altynaray T. Takibayeva, Irina V. Palamarchuk, Dastan M. Turdybekov, Assel A. Kelmyalene and Ivan V. Kulakov. Molecules 2022, 27(21), 7598; https://doi.org/10.3390/molecules27217598 - 05 Nov 2022

- Synthesis, Biosynthesis, and Biological Activity of Diels–Alder Adducts from Morus Genus: An Update. by Carola Tortora, Luca Pisano, Valeria Vergine, Francesca Ghirga, Antonia Iazzetti, Andrea Calcaterra, Violeta Marković, Bruno Botta and Deborah Quaglio. Molecules 2022, 27(21), 7580; https://doi.org/10.3390/molecules27217580 - 04 Nov 2022
- Synthesis and Evaluation of 11-Butyl Matrine Derivatives as Potential Anti-Virus Agents. by Wanjun Ni, Lizhong Wang, Hongjian Song, Yuxiu Liu and Qingmin Wang. Molecules 2022, 27(21), 7563; https://doi.org/10.3390/molecules27217563 04 Nov 2022
- Identification of Bcl2 as a Stably Expressed qPCR Reference Gene for Human Colon Cancer Cells Treated with Cottonseed-Derived Gossypol and Bioactive Extracts and Bacteria-Derived Lipopolysaccharides. by Heping Cao and Kandan Sethumadhavan. Molecules 2022, 27(21), 7560; https://doi.org/10.3390/molecules27217560 - 04 Nov 2022
- Eight Indole Alkaloids from the Roots of Maerua siamensis and Their Nitric Oxide Inhibitory Effects. by Sasiwimon Nukulkit, Angkana Jantimaporn, Preeyaporn Poldorn, Mattaka Khongkow, Thanyada Rungrotmongkol, Hsun-Shuo Chang, Rutt Suttisri and Chaisak Chansriniyom. Molecules 2022, 27(21), 7558; https://doi.org/10.3390/molecules27217558 04 Nov 2022
- Extraction with Acidified Methanol—An Easy and Effective Method of Methyl Chlorogenate Formation, as Studied by ESI-MS. by Karolina Szymborska, Rafał Frański and Monika Beszterda-Buszczak. Molecules 2022, 27(21), 7543; https://doi.org/10.3390/molecules27217543 - 03 Nov 2022
- Pyridine Carboxamides Based on Sulfobetaines: Design, Reactivity, and Biological Activity. by Eugene P. Kramarova, Sophia S. Borisevich, Edward M. Khamitov, Alexander A. Korlyukov, Pavel V. Dorovatovskii, Anastasia D. Shagina, Konstantin S. Mineev, Dmitri V. Tarasenko, Roman A. Novikov, Alexey A. Lagunin, Ivan Boldyrev, Aiarpi A. Ezdoglian, Natalia Yu. Karpechenko, Tatiana A. Shmigol, Yuri I. Baukov and Vadim V. Negrebetsky. Molecules 2022, 27(21), 7542; https://doi.org/10.3390/molecules27217542 - 03 Nov 2022
- Antioxidants of Amaranth, Quinoa and Buckwheat Wholemeals and Heat-Damage Development in Pseudocereal-Enriched Einkorn Water Biscuits. by Lorenzo Estivi, Luisa Pellegrino, Johannes A. Hogenboom, Andrea Brandolini and Alyssa Hidalgo. Molecules 2022, 27(21), 7541; https://doi.org/10.3390/molecules27217541 - 03 Nov 2022
- Phytochemical Analysis and Anti-Cancer Properties of Extracts of Centaurea castriferrei Borbás & Waisb Genus of Centaurea L. by Joanna Kubik, Łukasz Waszak, Grzegorz Adamczuk, Ewelina Humeniuk, Magdalena Iwan, Kamila Adamczuk, Mariola Michalczuk, Agnieszka Korga-Plewko and Aleksandra Józefczyk. Molecules 2022, 27(21), 7537; https://doi.org/10.3390/molecules27217537 - 03 Nov 2022
- Phytochemistry and Pharmacology of Medicinal Plants Used by the Tenggerese Society in Java Island of Indonesia. by Ari Satia Nugraha, Riza Putri Agustina, Syafi Mirza, Dinar Mutia Rani, Naura Bathari Winarto, Bawon Triatmoko, Antonius Nugraha Widhi Pratama, Paul A. Keller and Phurpa Wangchuk Molecules 2022, 27(21), 7532; https://doi.org/10.3390/molecules27217532 - 03 Nov 2022
- Propolis Efficacy: The Quest for Eco-Friendly Solvents. by Ana Sofia Freitas, Ana Cunha, Pier Parpot, Susana M. Cardoso, Rui Oliveira and Cristina Almeida-Aguiar. Molecules 2022, 27(21), 7531; https://doi.org/10.3390/molecules27217531 - 03 Nov 2022
- Combination of Spray-Chilling and Spray-Drying Techniques to Protect Carotenoid-Rich Extracts from Pumpkin (Cucurbita moschata) Byproducts, Aiming at the Production of a Powdered Natural Food Dye. by Priscilla Magalhães de Lima, Gustavo César Dacanal, Lorena Silva Pinho, Samuel Henrique Gomes de Sá, Marcelo Thomazini and Carmen Sílvia Favaro-Trindade. Molecules 2022, 27(21), 7530; https://doi.org/10.3390/molecules27217530 - 03 Nov 2022
- Iphiona mucronata (Forssk.) Asch. & Schweinf. A Comprehensive Phytochemical Study via UPLC-Q-TOF-MS in the Context of the Embryo- and Cytotoxicity Profiles. by Łukasz Pecio,

Asmaa M. Otify, Fatema R. Saber, Yasser A. El-Amier, Moataz Essam Shalaby, Solomiia Kozachok, Amira K. Elmotayam, Łukasz Świątek, Adrianna Skiba andKrystyna Skalicka-Woźniak. Molecules 2022, 27(21), 7529; https://doi.org/10.3390/molecules27217529 - 03 Nov 2022

- Influence of Altitudes and Development Stages on the Chemical Composition, Antioxidant, and Antimicrobial Capacity of the Wild Andean Blueberry (Vaccinium floribundum Kunth). by Mabel Guevara-Terán, Katherine Padilla-Arias, Andrea Beltrán-Novoa, Ana M. González-Paramás, Francesca Giampieri, Maurizio Battino, Wilson Vásquez-Castillo, Paulina Fernandez-Soto, Eduardo Tejera and José M. Alvarez-Suarez. Molecules 2022, 27(21), 7525; https://doi.org/10.3390/molecules27217525 - 03 Nov 2022
- Progress in the Preclinical and Clinical Study of Resveratrol for Vascular Metabolic Disease. by Dongxiao Fan, Chenshu Liu, Zhongyu Zhang, Kan Huang, Tengyao Wang, Sifan Chen and Zilun Li. Molecules 2022, 27(21), 7524; https://doi.org/10.3390/molecules27217524 - 03 Nov 2022
- Recent Advances in Herbal-Derived Products with Skin Anti-Aging Properties and Cosmetic Applications. by Erika F. Costa, Wagner V. Magalhães and Luiz C. Di Stasi. Molecules 2022, 27(21), 7518; https://doi.org/10.3390/molecules27217518 03 Nov 2022
- Phenolic Compounds in Active Packaging and Edible Films/Coatings: Natural Bioactive Molecules and Novel Packaging Ingredients. by Ajit Kumar Singh, Jae Young Kim and Youn Suk Lee. Molecules 2022, 27(21), 7513; https://doi.org/10.3390/molecules27217513 - 03 Nov 2022
- Multivariate Analysis Revealed Ultrasonic-Assisted Extraction Improves Anti-Melanoma Activity of Non-Flavonoid Compounds in Indonesian Brown Algae Ethanol Extract. by Anggit Listyacahyani Sunarwidhi, Ari Hernawan, Andri Frediansyah, Sri Widyastuti, Ni Wayan Riyani Martyasari, Angga Susmana Abidin, Hasriaton Padmi, Ervina Handayani, Ni Wayan Putri Utami, Farreh Alan Maulana, Muhammad Syach Maulad Ichfa and Eka Sunarwidhi Prasedya. Molecules 2022, 27(21), 7509; https://doi.org/10.3390/molecules27217509 - 03 Nov 2022
- Parkia platycephala Lectin (PPL) Inhibits Orofacial Nociception Responses via TRPV1 Modulation. by Gerlânia de Oliveira Leite, Sacha Aubrey Alves Rodrigues Santos, Romério Rodrigues dos Santos Silva, Claudener Souza Teixeira and Adriana Rolim Campos. Molecules 2022, 27(21), 7506; https://doi.org/10.3390/molecules27217506 - 03 Nov 2022
- Bioactive Compounds and Antioxidant Activity from Spent Coffee Grounds as a Powerful Approach for Its Valorization. by Carolina Andrade, Rosa Perestrelo and José S. Câmara. Molecules 2022, 27(21), 7504; https://doi.org/10.3390/molecules27217504 03 Nov 2022
- Effects of Tebuconazole Application on Soil Microbiota and Enzymes. by Małgorzata Baćmaga, Jadwiga Wyszkowska, Agata Borowik and Jan Kucharski. Molecules 2022, 27(21), 7501; https://doi.org/10.3390/molecules27217501 03 Nov 2022
- Phenolic Profile, EPR Determination, and Antiproliferative Activity against Human Cancer Cell Lines of Anthyllis vulneraria Extracts. by Manel Ouerfelli, Isidoro Metón, Idoia Codina-Torrella and María Pilar Almajano. Molecules 2022, 27(21), 7495; https://doi.org/10.3390/molecules27217495 - 03 Nov 2022
- Inhibitory Effect of Isoliquiritigenin in Niemann-Pick C1-Like 1-Mediated Cholesterol Uptake. by Jun Zeng, Wenjing Liu, Bing Liang, Lingyu Shi, Shanbo Yang, Jingsen Meng, Jing Chang, Xiaokun Hu, Renshuai Zhang and Dongming Xing. Molecules 2022, 27(21), 7494; https://doi.org/10.3390/molecules27217494 - 03 Nov 2022
- Potential Anti-Tumor Activity of Nardoguaianone L Isolated from Nardostachys jatamansi DC. in SW1990 Cells. by Chun-Yan Sang, Yi-Dan Zheng, Li-Mei Ma, Kai Wang, Cheng-Bo Wang, Tian Chai, Komila A. Eshbakova and Jun-Li Yang. Molecules 2022, 27(21), 7490; https://doi.org/10.3390/molecules27217490 - 03 Nov 2022
- Antihypothyroid Effect of Salidroside. by Nazym K. Korbozova, Nataliya O. Kudrina, Nataliya A. Zhukova, Alexander E. Grazhdannikov, Irina V. Blavachinskaya, Gulnaz A. Seitimova, Timur

E. Kulmanov, Tatyana G. Tolstikova and Nina V. Terletskaya. Molecules 2022, 27(21), 7487; https://doi.org/10.3390/molecules27217487 - 02 Nov 2022

- Medicinal Values and Potential Risks Evaluation of Ginkgo biloba Leaf Extract (GBE) Drinks Made from the Leaves in Autumn as Dietary Supplements. by Xiaojia Su, Ruirui Shi, Haiyan Hu, Linfeng Hu, Qichao Wei, Yuanyuan Guan, Jingling Chang and Chengwei Li. Molecules 2022, 27(21), 7479; https://doi.org/10.3390/molecules27217479 - 02 Nov 2022
- The Chemical Composition and Antimitotic, Antioxidant, Antibacterial and Cytotoxic Properties of the Defensive Gland Extract of the Beetle, Luprops tristis Fabricius. by Ovungal Sabira, Attuvalappil Ramdas Vignesh, Anthyalam Parambil Ajaykumar, Sudhir Rama Varma, Kodangattil Narayanan Jayaraj, Merin Sebastin, Kalleringal Nikhila, Annet Babu, Vazhanthodi Abdul Rasheed, Valiyaparambil Sivadasan Binitha, Zeena koldath Vasu and Madathilpadi Subrahmanian Sujith. Molecules 2022, 27(21), 7476; https://doi.org/10.3390/molecules27217476 - 02 Nov 2022
- Blueberry Anthocyanins from Commercial Products: Structure Identification and Potential for Diabetic Retinopathy Amelioration. by Rui Li, Zhan Ye, Wei Yang, Yong-Jiang Xu, Chin-Ping Tan and Yuanfa Liu. Molecules 2022, 27(21), 7475; https://doi.org/10.3390/molecules27217475 02 Nov 2022
- First Evidence of a Combination of Terpinen-4-ol and α-Terpineol as a Promising Tool against ESKAPE Pathogens. by Bianca Johansen, Raphaël E. Duval and Jean-Christophe Sergere. Molecules 2022, 27(21), 7472; https://doi.org/10.3390/molecules27217472 - 02 Nov 2022
- Advances in Research on Bioactivity, Toxicity, Metabolism, and Pharmacokinetics of Usnic Acid In Vitro and In Vivo. by Hanxue Wang, Min Xuan, Cheng Huang and Changhong Wang. Molecules 2022, 27(21), 7469; https://doi.org/10.3390/molecules27217469 02 Nov 2022
- Ginger Constituent 6-Shogaol Attenuates Vincristine-Induced Activation of Mouse Gastroesophageal Vagal Afferent C-Fibers. by Mayur J. Patil, Yongming Huang, Mingwei Yu, Xinzhong Dong, Bradley J. Undem and Shaoyong Yu. Molecules 2022, 27(21), 7465; https://doi.org/10.3390/molecules27217465 - 02 Nov 2022
- Antineoplastic and Antitrypanosomal Properties of Propolis from Tetragonula biroi Friese. by Samyah Alanazi. Molecules 2022, 27(21), 7463; https://doi.org/10.3390/molecules27217463
 - 02 Nov 2022
- Janus-Type AIE Fluorophores: Synthesis and Properties of π-Extended Coumarin-Bearing Triskelions. by Masafumi Ueda, Mirai Kokubun, Nao Yanagi, Norifumi Yamamoto and Yasuhiro Mazaki. Molecules 2022, 27(21), 7450; https://doi.org/10.3390/molecules27217450 - 02 Nov 2022
- Development of a GC/Q-ToF-MS Method Coupled with Headspace Solid-Phase Microextraction to Evaluate the In Vitro Metabolism of β-Caryophyllene. by Joseph Lee, Mei Wang, Goutam Mondal, Ikhlas A. Khan and Charles R. Yates. Molecules 2022, 27(21), 7441; https://doi.org/10.3390/molecules27217441 - 02 Nov 2022
- Production of High-Quality Red Fruit Juices by Athermal Membrane Processes. by Rosanna Morelli, Carmela Conidi, Rosa Tundis, Monica R. Loizzo, Massimo D'Avella, Rosario Timpone and Alfredo Cassano. Molecules 2022, 27(21), 7435; https://doi.org/10.3390/molecules27217435 - 01 Nov 2022
- Chemical Composition and Biological Activities of Prangos ferulacea Essential Oils. by Natale Badalamenti, Viviana Maresca, Michela Di Napoli, Maurizio Bruno, Adriana Basile and Anna Zanfardino. Molecules 2022, 27(21), 7430; https://doi.org/10.3390/molecules27217430 - 01 Nov 2022
- Phenolic Profiles, Antioxidant and Anti-Inflammatory Activities of Hydrodistillation Wastewaters from Five Lamiaceae Species. by Edoardo Napoli, Giuseppe Ruberto, Alessandra Carrubba, Mauro Sarno, Claudia Muscarà, Antonio Speciale, Mariateresa Cristani, Francesco Cimino and Antonella Saija. Molecules 2022, 27(21), 7427; https://doi.org/10.3390/molecules27217427 - 01 Nov 2022

- Synthesis of Novel Bromophenol with Diaryl Methanes—Determination of Their Inhibition Effects on Carbonic Anhydrase and Acetylcholinesterase. by Necla Oztaskin, Suleyman Goksu, Yeliz Demir, Ahmet Maras and İlhami Gulcin. Molecules 2022, 27(21), 7426; https://doi.org/10.3390/molecules27217426 01 Nov 2022
- Characterization of Conyza bonariensis Allelochemicals against Broomrape Weeds. by Antonio Cala Peralta, Gabriele Soriano, Jesús G. Zorrilla, Marco Masi, Alessio Cimmino and Mónica Fernández-Aparicio. Molecules 2022, 27(21), 7421; https://doi.org/10.3390/molecules27217421 - 01 Nov 2022
- Targeted and Non-Targeted HPLC Analysis of Coffee-Based Products as Effective Tools for Evaluating the Coffee Authenticity. by Jitka Klikarová and Lenka Česlová. Molecules 2022, 27(21), 7419; https://doi.org/10.3390/molecules27217419 01 Nov 2022
- Impact of TMPRSS2 Expression, Mutation Prognostics, and Small Molecule (CD, AD, TQ, and TQFL12) Inhibition on Pan-Cancer Tumors and Susceptibility to SARS-CoV-2. by Jiewen Fu, Shuguang Liu, Qi Tan, Zhiying Liu, Jie Qian, Ting Li, Jiaman Du, Binghui Song, Dabing Li, Lianmei Zhang, Jiayue He, Kan Guo, Baixu Zhou, Hanchun Chen, Shangyi Fu, Xiaoyan Liu, Jingliang Cheng, Tao He and Junjiang Fu. Molecules 2022, 27(21), 7413; https://doi.org/10.3390/molecules27217413 01 Nov 2022
- Curcumin and Its Derivatives Induce Apoptosis in Human Cancer Cells by Mobilizing and Redox Cycling Genomic Copper Ions. by Mohammed Ahmed Ismail Alhasawi, Mohammad Aatif, Ghazala Muteeb, Mir Waqas Alam, Mohamed El Oirdi and Mohd Farhan. Molecules 2022, 27(21), 7410; https://doi.org/10.3390/molecules27217410 - 01 Nov 2022
- Effect of Grinding Process Parameters and Storage Time on Extraction of Antioxidants from Ginger and Nutmeg. by Szymon Poliński, Patrycja Topka, Małgorzata Tańska, Sylwia Kowalska, Sylwester Czaplicki and Aleksandra Szydłowska-Czerniak. Molecules 2022, 27(21), 7395; https://doi.org/10.3390/molecules27217395 31 Oct 2022
- Phenylethanoid and Phenylmethanoid Glycosides from the Leaves of Ligustrum robustum and Their Bioactivities. by Shi-Hui Lu, Hao-Jiang Zuo, Jing Huang, Ran Chen, Jia-Ping Pan and Xiu-Xia Li. Molecules 2022, 27(21), 7390; https://doi.org/10.3390/molecules27217390 31 Oct 2022
- Simultaneous Extraction and Determination of Characteristic Steroidal Saponins and Homoisoflavonoids in Zhejiang Ophiopogon japonicus. by Yaoyao Zhu, Liling Wang, Meixu Chen, Yifeng Zhou and Jun Huang. Molecules 2022, 27(21), 7380; https://doi.org/10.3390/molecules27217380 - 30 Oct 2022
- Interactions between Gut Microbiota and Polyphenols: New Insights into the Treatment of Fatigue. by Chuanhong Luo, Xichuan Wei, Jiao Song, Xiaorong Xu, Haozhou Huang, Sanhu Fan, Dingkun Zhang, Li Han and Junzhi Lin. Molecules 2022, 27(21), 7377; https://doi.org/10.3390/molecules27217377 - 30 Oct 2022
- The Design of Rapid Self-Healing Alginate Hydrogel with Dendritic Crosslinking Network. by Dingxuan Wang, Yuhan Li, Haobo Zhang, Zhaorong Ren, Kefan Fan, Jue Cheng, Junying Zhang and Feng Gao. Molecules 2022, 27(21), 7367; https://doi.org/10.3390/molecules27217367 29 Oct 2022
- Characterization of Polyhydroxybutyrate, PHB, Synthesized by Newly Isolated Haloarchaea Halolamina spp. by Nashwa Hagagy, Amna A. Saddiq, Hend M. Tag, Samy Selim, Hamada AbdElgawad and Rosa María Martínez-Espinosa. Molecules 2022, 27(21), 7366; https://doi.org/10.3390/molecules27217366 - 29 Oct 2022
- Application of Metabolomics in Fungal Research. by Guangyao Li, Tongtong Jian, Xiaojin Liu, Qingtao Lv, Guoying Zhang and Jianya Ling. Molecules 2022, 27(21), 7365; https://doi.org/10.3390/molecules27217365 - 29 Oct 2022
- Isolation and Identification of Secondary Metabolites Produced by Phytopathogenic Fungus Corynespora cassiicola from Hevea brasiliensis. by Xiaoyan Yang, Zhikai Guo, Yang Yang,

Ailiman Abulaizi, Zijun Xiong, Shiqing Zhang, Boxun Li and Guixiu Huang. Molecules 2022, 27(21), 7360; https://doi.org/10.3390/molecules27217360 - 29 Oct 2022

- Functionalization of 7-Hydroxy-pyranoflavylium: Synthesis of New Dyes with Extended Chromatic Stability. by Ana Rita Pereira, Victor de Freitas, Nuno Mateus and Joana Oliveira. Molecules 2022, 27(21), 7351; https://doi.org/10.3390/molecules27217351 29 Oct 2022
- Potential Application of Luteolin as an Active Antibacterial Composition in the Development of Hand Sanitizer Products. by Meihua Xi, Yujie Hou, Ruolin Wang, Minhui Ji, Yingying Cai, Jingfang Ao, Heyu Shen, Mei Li, Jun Wang and Anwei Luo. Molecules 2022, 27(21), 7342; https://doi.org/10.3390/molecules27217342 28 Oct 2022
- Bioactive Compounds, Pharmacological Actions, and Pharmacokinetics of Genus Acacia. by Gaber El-Saber Batiha, Nosheen Akhtar, Abdulrahman A. Alsayegh, Wafaa Fouzi Abusudah, Najlaa Hamed Almohmadi, Hazem M. Shaheen, Thakur Gurjeet Singh and Michel De Waard. Molecules 2022, 27(21), 7340; https://doi.org/10.3390/molecules27217340 - 28 Oct 2022
- Essential Oil Chemotypes and Genetic Variability of Cinnamomum verum Leaf Samples Commercialized and Cultivated in the Amazon. by Júlia Karla A. M. Xavier, Talissa Gabriele C. Baia,Oscar Victor C. Alegria,Pablo Luis B. Figueiredo,Adriana R. Carneiro,Edith Cibelle de O. Moreira,José Guilherme S. Maia,William N. Setzer andJoyce Kelly R. da Silva. Molecules 2022, 27(21), 7337; https://doi.org/10.3390/molecules27217337 - 28 Oct 2022
- Chemical Profiling and Biological Activity of Extracts from Nine Norwegian Medicinal and Aromatic Plants. by Rune Slimestad, Amritha Johny, Mette Goul Thomsen, Christian Renè Karlsen and Jan Thomas Rosnes. Molecules 2022, 27(21), 7335; https://doi.org/10.3390/molecules27217335 - 28 Oct 2022
- Recent Updates on the Functional Impact of Kahweol and Cafestol on Cancer. by Salma Eldesouki, Rama Qadri, Rashid Abu Helwa, Hiba Barqawi, Yasser Bustanji, Eman Abu-Gharbieh and Waseem El-Huneidi. Molecules 2022, 27(21), 7332; https://doi.org/10.3390/molecules27217332 28 Oct 2022
- Chemical Composition and Antioxidant, Antimicrobial, and Anti-Inflammatory Properties of Origanum compactum Benth Essential Oils from Two Regions: In Vitro and In Vivo Evidence and In Silico Molecular Investigations. by Samiah Hamad Al-Mijalli, Nidal Naceiri Mrabti, Hayat Ouassou, Ryan A. Sheikh, Hamza Assaggaf, Saad Bakrim, Emad M. Abdallah, Mohammed Merae Alshahrani, Ahmed Abdullah Al Awadh, Learn-Han Lee, Yusra AlDhaheri,Amirhossein Sahebkar,Gokhan Zengin,Ammar A. Attar,Abdelhakim Bouyahya and Hanae Naceiri Mrabti. Molecules 2022, 27(21), 7329; https://doi.org/10.3390/molecules27217329 - 28 Oct 2022
- 7-Hydroxycoumarin Induces Vasorelaxation in Animals with Essential Hypertension: Focus on Potassium Channels and Intracellular Ca2+ Mobilization. by Rafael L. C. Jesus, Isnar L. P. Silva, Fênix A. Araújo, Raiana A. Moraes, Liliane B. Silva, Daniele S. Brito, Gabriela B. C. Lima, Quiara L. Alves and Darizy F. Silva. Molecules 2022, 27(21), 7324; https://doi.org/10.3390/molecules27217324 - 28 Oct 2022
- First Evidence of Gastroprotection by Schinus molle: Roles of Nitric Oxide, Prostaglandins, and Sulfhydryls Groups in Its Mechanism of Action. by María Elena Sánchez-Mendoza, Yaraset López-Lorenzo, Leticia Cruz-Antonio, Daniel Arrieta-Baez, Miranda Carolina Pérez-González and Jesús Arrieta. Molecules 2022, 27(21), 7321; https://doi.org/10.3390/molecules27217321 - 28 Oct 2022
- Variation of the Antioxidative Defense in Elaeis guineensis Jacq. Facing Bud Rot Disease in the Coastal Area of Ecuador. by Raluca A. Mihai, Galo M. Canchignia Guacollantes, Sebastián A. Villacrés Mesias, Larisa I. Florescu and Rodica D. Catana. Molecules 2022, 27(21), 7314; https://doi.org/10.3390/molecules27217314 - 27 Oct 2022
- Rutin Improves Anxiety and Reserpine-Induced Depression in Rats. by Ahmed I Foudah, Mohammed H Alqarni, Aftab Alam, Sushma Devi, Mohammad A Salkini and Prawez Alam. Molecules 2022, 27(21), 7313; https://doi.org/10.3390/molecules27217313 - 27 Oct 2022

- Brazilian Red Propolis Presents Promising Anti-H. pylori Activity in In Vitro and In Vivo Assays with the Ability to Modulate the Immune Response. by Mariana B. Santiago, Luis Fernando Leandro, Rafael B. Rosa, Murilo V. Silva ,Samuel C. Teixeira, João Paulo S. Servato, Sérgio Ricardo Ambrósio, Rodrigo Cassio S. Veneziani, Jennyfer A. Aldana-Mejía, Jairo K. Bastos andCarlos Henrique G. Martins. Molecules 2022, 27(21), 7310; https://doi.org/10.3390/molecules27217310 - 27 Oct 2022
- Native Mass Spectrometry Coupled to Spectroscopic Methods to Investigate the Effect of Soybean Isoflavones on Structural Stability and Aggregation of Zinc Deficient and Metal-Free Superoxide Dismutase. by Xinyu Bian, Xiaoyu Zhuang, Junpeng Xing, Shu Liu, Zhiqiang Liu and Fengrui Song. Molecules 2022, 27(21), 7303; https://doi.org/10.3390/molecules27217303 -27 Oct 2022
- A Composition Analysis and an Antibacterial Activity Mechanism Exploration of Essential Oil Obtained from Artemisia giraldii Pamp. by Guiguo Huo, Xu Li, Mohamed Aamer Abubaker, Tingyu Liang, Ji Zhang and Xuelin Chen. Molecules 2022, 27(21), 7300; https://doi.org/10.3390/molecules27217300 - 27 Oct 2022
- Inhibitory Effect of Sodium Alginate Nanoemulsion Coating Containing Myrtle Essential Oil (Myrtus communis L.) on Listeria monocytogenes in Kasar Cheese. by Gökçe Polat Yemiş, Elif Sezer and Hatice Sıçramaz. Molecules 2022, 27(21), 7298; https://doi.org/10.3390/molecules27217298 - 27 Oct 2022
- Guaianolide Derivatives from the Invasive Xanthium spinosum L.: Evaluation of Their Allelopathic Potential. by Sylvain Baldi, Pascale Bradesi and Alain Muselli. Molecules 2022, 27(21), 7297; https://doi.org/10.3390/molecules27217297 27 Oct 2022
- Beyond the Bark: An Overview of the Chemistry and Biological Activities of Selected Bark Essential Oils. by Melanie Graf and Iris Stappen. Molecules 2022, 27(21), 7295; https://doi.org/10.3390/molecules27217295 27 Oct 2022
- Anticancer, Cardio-Protective and Anti-Inflammatory Potential of Natural-Sources-Derived Phenolic Acids. by Afnan, Ammara Saleem, Muhammad Furqan Akhtar, Ali Sharif, Bushra Akhtar, Rida Siddique, Ghulam Md Ashraf, Badrah S. Alghamdi and Saif A. Alharthy. Molecules 2022, 27(21), 7286; https://doi.org/10.3390/molecules27217286 26 Oct 2022
- Phytochemistry Meets Geochemistry—Blumenol C Sulfate: A New Megastigmane Sulfate from Palicourea luxurians (Rubiaceae: Palicoureeae). by Christoph Kornpointner, Nadine J. Hochenegger, Bao-Bao Shi, Andreas Berger, Johannes Theiner, Lothar Brecker and Johann Schinnerl. Molecules 2022, 27(21), 7284; https://doi.org/10.3390/molecules27217284 - 26 Oct 2022
- Atomic-Resolution Experimental Structural Biology and Molecular Dynamics Simulations of Hyaluronan and Its Complexes. by Olgun Guvench. Molecules 2022, 27(21), 7276; https://doi.org/10.3390/molecules27217276 26 Oct 2022
- Untargeted Metabolomics Reveals New Markers of Food Processing for Strawberry and Apple Purees. by Gabriela Salazar-Orbea, Rocío García-Villalba, Luis M. Sánchez-Siles, Francisco A. Tomás-Barberán and Carlos J. García. Molecules 2022, 27(21), 7275; https://doi.org/10.3390/molecules27217275 - 26 Oct 2022
- Immunosuppressive Sesquiterpene Pyridine Alkaloids from Tripterygium wilfordii Hook. f.by Yadan Wang, Jiangong Yan, Zhongmou Zhang, Minghui Chen, Xianfu Wu and Shuangcheng Ma. Molecules 2022, 27(21), 7274; https://doi.org/10.3390/molecules27217274 - 26 Oct 2022
- Anthocyanins Formulated with Carboxymethyl Starch for Gastric and Intestinal Delivery. by Luiz Bruno De Sousa Sabino, Francesco Copes, Solène Saulais, Edy Sousa De Brito, Ivanildo José Da Silva Júnior, Tien Canh Le, Mircea Alexandru Mateescu and Diego Mantovani. Molecules 2022, 27(21), 7271; https://doi.org/10.3390/molecules27217271 - 26 Oct 2022
- The Antiviral Activity of Caprylic Monoglyceride against Porcine Reproductive and Respiratory Syndrome Virus In Vitro and In Vivo. by Luyu Yang, Jianhua Wen, Yang Zhang,

Zheyan Liu, Zhipeng Luo, Lei Xu, Siyuan Lai, Huaqiao Tang, Xiangang Sun, Youjun Hu, Ling Zhu and Zhiwen Xu. Molecules 2022, 27(21), 7263; https://doi.org/10.3390/molecules27217263 - 26 Oct 2022

- An Overview on the Synthesis of Fused Pyridocoumarins with Biological Interest. by Matina D. Douka and Konstantinos E. Litinas. Molecules 2022, 27(21), 7256; https://doi.org/10.3390/molecules27217256 26 Oct 2022
- Studies on Chemical Composition of Pueraria lobata and Its Anti-Tumor Mechanism. by Xiaoxue Fang, Yegang Zhang, Yiming Cao, Mengyao Shan, Dimeng Song, Chao Ye and Difu Zhu. Molecules 2022, 27(21), 7253; https://doi.org/10.3390/molecules27217253 - 26 Oct 2022
- Dark Sweet Cherry (Prunus avium) Anthocyanins Suppressed ERK1/2-Akt/mTOR Cell Signaling and Oxidative Stress: Implications for TNBC Growth and Invasion. by Ana Carolina Silveira Rabelo, Susanne U. Mertens-Talcott, Boon P. Chew and Giuliana Noratto. Molecules 2022, 27(21), 7245; https://doi.org/10.3390/molecules27217245 - 25 Oct 2022
- Chemical Composition of Tagetes patula Flowers Essential Oil and Hepato-Therapeutic Effect against Carbon Tetrachloride-Induced Toxicity (In-Vivo). by Hanan Y. Aati, Mahmoud Emam, Jawaher Al-Qahtani, Sultan Aati, Abdulrahman Aati, Juergen Wanner and Mohamed M. Seif. Molecules 2022, 27(21), 7242; https://doi.org/10.3390/molecules27217242 25 Oct 2022
- Challenges in the Heterologous Production of Furanocoumarins in Escherichia coli. by Joana L. Rodrigues, Daniela Gomes and Lígia R. Rodrigues. Molecules 2022, 27(21), 7230; https://doi.org/10.3390/molecules27217230 25 Oct 2022
- Experimental and Theoretical Study on the Homodimerization Mechanism of 3-Acetylcoumarin. by Kristina B. Simeonova, Ana I. Koleva, Anna-Mariya R. Zlatanova, Nevena I. Petkova-Yankova, Hristiyan A. Aleksandrov, Petko St. Petkov andRositca D. Nikolova. Molecules 2022, 27(21), 7228; https://doi.org/10.3390/molecules27217228 - 25 Oct 2022
- Therapeutic Potential of Select Dietary Compounds in the Management of Hypertension and its Cardiovascular Complications. by Aleena Francis Valookaran, Jenny Bouchard, Basma Milad Aloud, Sijo Joseph Thandapilly and Thomas Netticadan. Molecules 2022, 27(21), 7222; https://doi.org/10.3390/molecules27217222 25 Oct 2022
- The Antibacterial and Larvicidal Potential of Bis-(2-Ethylhexyl) Phthalate from Lactiplantibacillus plantarum. by Muhammad Rizwan Javed, Mahwish Salman, Anam Tariq, Abdul Tawab, Muhammad Kashif Zahoor, Shazia Naheed, Misbah Shahid, Anam Ijaz andHazrat Ali. Molecules 2022, 27(21), 7220; https://doi.org/10.3390/molecules27217220 25 Oct 2022
- Fatty-Acid-Rich Agave angustifolia Fraction Shows Antiarthritic and Immunomodulatory Effect. by Enrique Jiménez-Ferrer, Gabriela Vargas-Villa, Gabriela Belen Martínez-Hernández, Manases González-Cortazar, Alejandro Zamilpa, Maribel Patricia García-Aguilar, Martha Lucía Arenas-Ocampo and Maribel Herrera-Ruiz. Molecules 2022, 27(21), 7204; https://doi.org/10.3390/molecules27217204 - 24 Oct 2022
- Variations in the Chemical Composition of Essential Oils in Native Populations of Korean Thyme, Thymus quinquecostatus Celak. by Minju Kim, Kandhasamy Sowndhararajan, Ponnuvel Deepa and Songmun Kim. Molecules 2022, 27(21), 7203; https://doi.org/10.3390/molecules27217203 - 24 Oct 2022
- Pharmacodynamic Interactions between Puerarin and Metformin in Type-2 Diabetic Rats. by Zhen Li, Shengguang Wang, Xinyu Wang, Peng Gao, Shiming Zhang, Yingning Mo, Dongsheng Zhao and Long Dai. Molecules 2022, 27(21), 7197; https://doi.org/10.3390/molecules27217197 - 24 Oct 2022
- Analysis of Changes in the Amount of Phytosterols after the Bleaching Process of Hemp Oils. by Andrzej Kwaśnica, Mirosława Teleszko, Damian Marcinkowski, Dominik Kmiecik, Anna Grygier and Wojciech Golimowski. Molecules 2022, 27(21), 7196; https://doi.org/10.3390/molecules27217196 - 24 Oct 2022

- Valorization of Phenolic and Carotenoid Compounds of Sechium edule (Jacq. Swartz) Leaves: Comparison between Conventional, Ultrasound- and Microwave-Assisted Extraction Approaches. by Elsa F. Vieira, Suene Souza, Manuela M. Moreira, Rebeca Cruz, Aline Boatto da Silva, Susana Casal and Cristina Delerue-Matos. Molecules 2022, 27(21), 7193; https://doi.org/10.3390/molecules27217193 - 24 Oct 2022
- Sources, Transformations, Syntheses, and Bioactivities of Monoterpene Pyridine Alkaloids and Cyclopenta[c]pyridine Derivatives. by Xuejian Zhang, Feiyan Tao, Tao Cui, Cheng Luo, Zhigang Zhou, Yuchuan Huang, Lanlan Tan, Wei Peng and Chunjie Wu. Molecules 2022, 27(21), 7187; https://doi.org/10.3390/molecules27217187 - 24 Oct 2022
- Review on Compounds Isolated from Eriocaulaceae Family and Evaluation of Biological Activities by Machine Learning. by Laysa Lanes Pereira Ferreira Moreira, Renata Priscila Barros de Menezes, Luciana Scotti, Marcus Tullius Scotti, Valdemar Lacerda Júnior and Warley de Souza Borges. Molecules 2022, 27(21), 7186; https://doi.org/10.3390/molecules27217186 - 24 Oct 2022
- Mutactimycin AP, a New Mutactimycin Isolated from an Actinobacteria from the Atacama Desert. by Rishi Vachaspathy Astakala, Gagan Preet,Bruce F. Milne, Julius Tibyangye, Valeria Razmilic, Jean Franco Castro, Juan A. Asenjo, Barbara Andrews, Rainer Ebel and Marcel Jaspars. Molecules 2022, 27(21), 7185; https://doi.org/10.3390/molecules27217185 - 24 Oct 2022
- Astounding Health Benefits of Jamun (Syzygium cumini) toward Metabolic Syndrome. by Maryam Khalid Rizvi, Roshina Rabail, Seemal Munir, Muhammad Inam-Ur-Raheem, Mir Muhammad Nasir Qayyum, Marek Kieliszek, Abdo Hassoun and Rana Muhammad Aadil. Molecules 2022, 27(21), 7184; https://doi.org/10.3390/molecules27217184 - 24 Oct 2022
- The Synthesis, Fungicidal Activity, and in Silico Study of Alkoxy Analogues of Natural Precocenes I, II, and III. by Khaled M. A. Ramadan, Hossam S. El-Beltagi, Zafar Iqbal and Eslam S. A. Bendary. Molecules 2022, 27(21), 7177; https://doi.org/10.3390/molecules27217177 - 24 Oct 2022
- The Role of Astaxanthin as a Nutraceutical in Health and Age-Related Conditions. by Geir Bjørklund, Amin Gasmi, Larysa Lenchyk, Mariia Shanaida, Saba Zafar, Pavan Kumar Mujawdiya, Roman Lysiuk, Halyna Antonyak, Sadaf Noor, Muhammad Akram, Kateryna Smetanina, Salva Piscopo, Taras Upyr and Massimiliano Peana. Molecules 2022, 27(21), 7167; https://doi.org/10.3390/molecules27217167 - 23 Oct 2022
- Valorisation, Green Extraction Development, and Metabolomic Analysis of Wild Artichoke By-Product Using Pressurised Liquid Extraction UPLC–HRMS and Multivariate Data Analysis. by Stefania Pagliari, Ciro Cannavacciuolo, Rita Celano, Sonia Carabetta, Mariateresa Russo, Massimo Labra and Luca Campone. Molecules 2022, 27(21), 7157; https://doi.org/10.3390/molecules27217157 - 22 Oct 2022
- Anxiolytic-like Effect of Quercetin Possibly through GABA Receptor Interaction Pathway: In Vivo and In Silico Studies. by Md. Shahazul Islam, Rajib Hossain, Taukir Ahmed, Md. Mizanur Rahaman, Khattab Al-Khafaji, Rasel Ahmed Khan, Chandan Sarkar, Mehedi Hasan Bappi, Edlane Martins de Andrade, Isaac Moura Araújo, Henrique Douglas Melo Coutinho, Grażyna Kowalska, Radosław Kowalski, Muhammad Asif Hanif and Muhammad Torequl Islam. Molecules 2022, 27(21), 7149; https://doi.org/10.3390/molecules27217149 - 22 Oct 2022





Phytochemistry and Pharmacology of Medicinal Plants Used by the Tenggerese Society in Java Island of Indonesia

Ari Satia Nugraha ^{1,2,*}, Riza Putri Agustina ¹, Syafi Mirza ¹, Dinar Mutia Rani ¹, Naura Bathari Winarto ¹, Bawon Triatmoko ¹, Antonius Nugraha Widhi Pratama ¹, Paul A. Keller ² and Phurpa Wangchuk ^{3,*}

- ¹ Drug Utilisation and Discovery Research Group, Faculty of Pharmacy, Universitas Jember, Jember 68121, Indonesia
- ² School of Chemistry, Faculty of Science Medicine and Health, University of Wollongong, Wollongong, NSW 2522, Australia
- ³ Centre for Molecular Therapeutics, Australian Institute of Tropical Health and Medicine, James Cook University, Cairns, QLD 4878, Australia
- * Correspondence: arisatia@unej.ac.id (A.S.N.); phurpa.wangchuk@jcu.edu.au (P.W.); Tel.: +62-331-324-736 (A.S.N.); +61-742321249 (P.W.)

Abstract: The archipelagic country of Indonesia is inhabited by 300 ethnic groups, including the indigenous people of Tengger. Based on the reported list of medicinal plants used by the Tengger community, we have reviewed each of them for their phytochemical constituents and pharmacological activities. Out of a total of 41 medicinal plants used by the Tengerrese people, 33 species were studied for their phytochemical and pharmacological properties. More than 554 phytochemicals with diverse molecular structures belonging to different chemical classes including flavonoids, terpenoids, saponins and volatiles were identified from these studied 34 medicinal plants. Many of these medicinal plants and their compounds have been tested for various pharmacological activities including anti-inflammatory, antimicrobial, wound healing, headache, antimalarial and hypertension. Five popularly used medicinal plants by the healers were *Garcinia mangostana, Apium graveolens, Cayratia clematidea, Drymocallis arguta* and *Elaeocarpus longifolius*. Only *A. graviolens* were previously studied, with the outcomes supporting the pharmacological claims to treat hypertension. Few unexplored medicinal plants are *Physalis lagascae, Piper amplum, Rosa tomentosa* and *Tagetes tenuifolia,* and they present great potential for biodiscovery and drug lead identification.

Keywords: Tengger; phytochemistry; pharmacology; *Cayratia clematidea*; *Drymocallis arguta*; *Elaeocarpus longifolius*; *Physalis lagascae*; *Piper amplum*; *Rosa tomentosa*; *Tagetes tenuifolia*

1. Introduction

Since ancient human civilizations, mankind has used biotic resources including plants for clothing, cosmetics, food and medication. The World Health Organisation (WHO) estimated that more than 80 % of the world's population rely on traditional medicines (TM) for their primary health needs [1]. Plants are the bulk ingredients used in these medicaments [1] with an estimated 50,000 plant species used worldwide, the majority of them contained within Asian medicines [2]. Most of the Asian medicinal plant knowledge is passed down uninterrupted from father to son using oral communication, or from master to apprentices using written scholarly traditions. The most popular scholarly medical traditions are Chinese traditional medicine [3], Indian Ayurvedic medicine [4] and Sowa Rigpa medicine (also practiced in Bhutan) [5]. The former oral traditions, which are predominantly practiced by remote tribes, are prone to disappearance or extinction [6,7].

In Indonesia there are more than 17,000 islands which are rich in biodiversity, especially the terrestrial plants. Indonesia has one of the highest numbers of higher plant species, with 22,500 species recorded so far. However, only a miniscule 4.4 % (1000 species) of these higher plant species are used as medicinal plants [8]. Since there are 300 ethnic



Citation: Nugraha, A.S.; Agustina, R.P.; Mirza, S.; Rani, D.M.; Winarto, N.B.; Triatmoko, B.; Pratama, A.N.W.; Keller, P.A.; Wangchuk, P. Phytochemistry and Pharmacology of Medicinal Plants Used by the Tenggerese Society in Java Island of Indonesia. *Molecules* **2022**, *27*, 7532. https://doi.org/10.3390/ molecules27217532

Academic Editor: Federica Pellati

Received: 11 September 2022 Accepted: 1 November 2022 Published: 3 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). groups/tribes in Indonesia, one would expect to find a rich medicinal plants diversity. One of these 300 thriving communities in Indonesia is the Tengger tribal community, residing in the Bromo mountain range (1600–2000 m above sea level, masl) of East Java, with the region known for its breathtaking views (Figure 1). The people still practice Hinduism from the old Majapahit Hindu Kingdom (1300–1500 A.D.) [9], which arrived in Indonesia in the first century through Indian traders, with Brahmin passengers as direct agents in transmitting Hinduism. For this reason, it is likely that Tenggerese ethnobotanical practices would resemble Indian Ayurvedic medicines. It is also expected that the mainstream Islamic traditional medical culture of Indonesia may have influenced the way Tenggerese medicines have evolved over centuries.




Figure 1. Map of Indonesia. (**A**) Location of Tengger community within a Javanese island. (**B**) Bromo mountain range where Tenggerese community live.

There is no historical document to substantiate their influences, and there is a need for such studies in Indonesia. There are a few reports on the ethnobotanical studies of the Tenggerese community, including medicinal plant surveys in the Tenggeresse village of Wonokitri Village, Tosari subdistrict, Pasuruan Regency [10]. Another ethnobotanical survey was also previously reported in a different village, Ngadisari village, Sukapura district, Probolinggo Region [11]. Nevertheless, there is no comprehensive review on the phytochemical and pharmacological constituents of Tenggerese medicinal plants. In this review, we have collected the medicinal plants used by the Tenggerese community residing in the Wonokerso village, the oldest village in the Tenggerese community where the "Karo (blessing)" ceremony originated. The information regarding medicinal plants and their medicinal uses were collected through discussion and interviews with local physicians known as "dukuns". The ethnopharmacological information was initiated in May 2015, and a list of dukuns medicinal plants was generated and are listed in Section 2 Table 1. Based on the list of plants in Table 1, we conducted a thorough literature search for each plant for their phytochemical and pharmacological activities. Figure 2 shows the schematic approach of this literature review. We have also consulted and compared the ethnobotanical information of our survey with the published information described previously from the surrounding villages. For example, *Foeniculum vulgare* and *Acorus calamus*, which were previously described as fever from the neighbouring villages, were also found in the current surveys described by dukuns [10–12]. In order to retrieve the phytochemical information of medicinal plants from Google Scholar (https://scholar.google.com/, accessed on 9 August 2022), PubMed (https://pubmed.ncbi.nlm.nih.gov/, accessed on 9 August 2022) and Scifinder (https://scifinder.cas.org, accessed on 9 August 2022), we used keywords such as plant name, chemical constituent, phytochemical composition, and isolated compounds. Chemical names and molecular structures were authenticated using PubChem (https: //pubchem.ncbi.nlm.nih.gov/, accessed on 11 August 2022) and Chemspider (http://www. chemspider.com/, accessed on 11 August 2022). To collect pharmacological information, data were searched using the same databases, with the bioactivity of each species collected based on Tenggerese traditional uses as part of the keywords. To maintain the quality of information, we included only Scopus and PubMed indexed articles. We retrieved the

literature from 1975 to 2022, conducted meta-analysis and presented it in a bar graph, as shown in Figure 3.




Figure 2. Flow chart of our approach to scoping literature is presented here.




Figure 3. Retrieved articles related to phytochemistry and pharmacological studies of medicinal plants used by the Tenggeresse people. Papers were collected from a previous report (1975–2022) of the same plants studied across the globe for similar pharmacological claims (Google Scholar, PubMed, and SciFinder Scholar).

2. Phytochemistry of Tenggerese Medicinal Plants

The analysis of the reported ethnobotanical studies of Tenggerese medicinal plants revealed 41 species of medicinal plants (Table 1). Of these, 33 were studied for their phyto-

chemical composition, and seven species remain unstudied. More than 404 phytochemicals with diverse molecular structures were identified from the 33 medicinal plants studied (see Table 1). These phytochemicals belong to different chemical classes including flavonoids, terpenoids, alkaloids, saponins and volatiles. While a few plants were reported to contain two or three phytochemicals, other plants have been extensively studied, and as many as 30 phytochemicals were either detected or isolated from a single plant. For example, ellagic acid was the only phytochemical reported from *Rubus rosa*, and therefore further in-depth analysis of this plant is required. On the other hand, 48 phytochemicals have been identified from *Acorus calamus* (Table 1). Seven species that were not studied for their phytochemicals are *Cayratia clemaidea*, *Drymocallis arguta*, *Elaeocarpus longifolius*, *Physalis lagascae*, *Piper amplum*, *Rosa tomentosa* and *Tagetes tenuifolia* (Figure 4).




Figure 4. Pictures of understudied medicinal plants used by the Indigenous people of Tengger. (**A**): *Cayratia clemaidea*; (**B**): *Drymocallis arguta*: (**C**): *Elaeocarpus longifolius*; (**D**): *Physalis lagascae*; (**E**): *Piper amplum*, (**F**): *Rosa tomentosa*; (**G**): *Tagetes tenuifolia*.

It is interesting to note that although the medicinal plants listed in Table 1 have been used for many generations by the people of Tengger in Indonesia, this review found that most of the phytochemical and pharmacological studies on these plants were reported from other countries, including North and South Americas, Europe, Middle East and East Asia, and South East Asian countries. There are only limited phytochemical and pharmacological studies reported on medicinal plants that grow in the Tengger region, or even Indonesia as a whole. The few medicinal plants that were extensively studied in Indonesia for their phytochemicals are *C. burmanii*, *C. nucifera* and *S. grandiflora*.

Species	Family	Tenggeresse Ethnopharmacological Uses of Plants	Parts Used for Chemical Isolation	Countries (Chemical Studies Reported)	Isolated Compounds
<i>Acorus calamus</i> Linnaeus	Acoraceae	Fever	Leaves, rhizome, stem	India	 β-Asarone, Camphene, Cymene, Calarene, α-Selinene, s-Cadinol, Isoshyobunone, β-Sesquiphellandrene, Preiso-calamendiol, Acorone [13]; (-)-4-Terpineol, Epieudesmin, Lysidine, (-)-Spathulenol, Borneol, Furyl ethyl ketone, Nonanoic acid, Bornyl acetate, Galgravin, Retusin, Butyl butanoate, Geranylacetate, Sakuranin, Acetic acid, Camphor, Isoelemicin, α-Ursolic acid, Acetophenone, Dehydroabietic acid, Isoeugenol Methylether, Apigenin, Dehydrodiisoeugenol, Linalool, Elemicin, Linolenic acid [14]; 2-Deca-4,7-dienol, Acoradin, Acoragermacrone, Acrenone, Aterpineol, β-Cadinene, Calacorene, Calamendiol, Galangin, Shyobunones, Sitosterol [15]; Calamusins A-I [16].
Allium sativum Linnaeus	Alliaceae	Wound or cut	Rhizome	Iraq	<i>E</i> -Ajoene, <i>Z</i> -Ajoene, Alliin, Allicin, 2-Vinyl-4 <i>H</i> -1,3-dithiin, Diallyl sulfide (DAS), Diallyl disulfide (DADS), Diallyl trisulfide (DATS), Allyl methyl sulfide (AMS) [17].
<i>Alyxia reinwardtii</i> Blume	Apocynaceae	Fever, Rheumatism	Stem	Thailand	Coumarin, 3-Hydroxycoumarin, 6-Hydroxycoumarin, 8-Hydroxycoumarin, Scopoletin, (+)-Pinoresinol, Zhebeiresinol and <i>p</i> -Hydroxybenzoic acid [18].
Anredera cordifolia (Ten.) Steenis	Basellaceae	Itchiness, Wound	Leaves	Brazil	Phytol, α-pinene, Larreagenin A, Vitexin, Isovitexin, Myricetin, Morin, Lupeol, β -Sitosterol, Ursolic acid [19].
<i>Apium graveolens</i> Linnaeus	Apiaceae	Hypertension	Leaves	China	Apigenin, Luteolin, Chlorogenic acid [20]; Linalool, D-Limonene, 3-N-Butylphthalide (NBP) [21].
<i>Borreria laevis</i> (Lam.) Griseb	Rubiaceae	Rheumatism	Aerial parts	Thailand	Borreline, Asperulosidic acid, 6-O-Acetylscandoside, 6α -Hydroxyadoxoside, Kaempferol 3-O- β -D-glucopyranoside, Kaempferol 3-O-rutinoside, Quercetin 3-O- β -D-galactopyranoside, Rutin [22,23].
<i>Brassica rapa</i> Linnaeus	Brassicaceae	Fever, Hypertension, Nutrition	Leaves, stem, flower buds, roots	Portugal	Kaempferol 3-O-sophoroside-7-O-glucoside, Kaempferol 3-O (feruloy1/caffeoy1)-sophoroside7-O-glucoside, Isorhamnetin 3,7-O-diglucoside, Isorhamnetin 3-O-glucoside [24].
<i>Capsicum pubescens</i> Dun.	Solanaceae	Tonic after hard labour	Fruit	Mexico	Carotenoids (Violaxanthin, <i>cis</i> -Violaxanthin, Luteoxanthin, Antheraxanthin, Lutein, Zeaxanthin, β -Carotene), Ascorbic acid and Capsaicinoids (Capsaicin, Dihydrocapsaicin) [25].
<i>Cayratia clematidea</i> (F. Müll.) Domin	Vitaceae	Stomach disorder	NA	NA	NA
Cinnamomum burmannii (Nees & T. Nees) Bl.	Laruaceae	Fever		China, Indonesia	<i>Trans</i> -Cinnamaldehyde, Coumarin, and <i>Trans</i> -Cinnamic acid [26]. Styrene, Benzaldehyde, Camphene, β -Pinene, Borneol, α -Terpineol, Procyanidin B1, Procyanidin B2, Procyanidin trimer, Catechin, Procyanidin dimer, Epicatechin, Coumarin, (<i>E</i>)-Cinnamic acid, (<i>E</i>)-Cinnamaldehyde, (<i>Z</i>)-Cinnamaldehyde, Cinnamyl alcohol, (<i>E</i>)-cinnamaldehyde, eugenol, and coumarin, procyanidin trimer, (<i>E</i>)-cinnamaldehyde, and (<i>Z</i>)-cinnamaldehyde [27]. catechin, epicatechin, procyanidin B2, quercitrin, 3,4-dihydroxybenzaldehyde, protocatechuic acid, and cinnamic acid [28]. (<i>E</i>)-Cinnamaldehyde, Cinnamyl alcohol, Coumarin, 3,4-Dihydrocoumarin, Kaempferol, Procyanidin dimer, Procyanidin trimer, Linalool [29]

Table 1. Phytochemistry of Tenggerese medicinal plants collated from literature studies of similar species studied across the globe.

Species

Cocos nucifera

Linnaeus

Linnaeus

Table 1. Cont. Tenggeresse Parts Used for Chemical **Countries** (Chemical Family Ethnopharmacological **Isolated Compounds** Isolation **Studies Reported)** Uses of Plants 2-Furaldehyde diethyl acetal and Palmitic acid [30]; Jezonofol, Cirrhusin A, Cassigarol G, Maackin A, Treoguiacyl glycerol-8'-vanil ether acid, Foetus health India, Indonesia, Brazil, UK Erythroguiacyl glycerol-8'-vanillic acid ether, Apigenin-7-O- β -D-glucoside, Aracaceae Fruit Piceatannol, p-Hydroxybenzoic acid, Protocatechuic acid, and Vanillic acid [31]; Two phenol compounds-catechin and Chlorogenic acid [32]. Cuminaldehyde, α -Pinene, β -Pinene, γ -Cymene, γ -Terpinene, Cuminum cyminum Apiaceae Fever Seed USA, Iraq α -Terpinen-7-al and β -Terpinen-7-al [33]; Bergapten, Methoxsalen [34]; Luteolin, Apigenin-7-O-glucoside [35]. Curcuminoids, Demethoxycurcumin, Bisdemethoxycurcumin [36]; Calebin-A [37]; *α*-Turmerone [38]; Epicatechins [39]; Cucurbitacin B, Fever, Thailand China Belgium

<i>Curcuma longa</i> Linnaeus	Zingiberaceae	Fever, Headache, Wound	Rhizome	Thailand, China, Belgium, Vietnam, Germany	Calebin-A [37]; α -Turmerone [38]; Epicatechins [39]; Cucurbitacin B, Curcumin [40]; Bisacurone B [41]; α -Curcumene, Zingiberene, Bisabolene, Sesquiphellandrene [42]; Turmeronol B, Turmeronol A, (<i>E</i>)- α -Atlantone [43]; Curlone [44].
Datura metel Linnaeus	Solanaceae	Fever	Leaves, Flower	China	Daturafolisides, Daturametelin [45]; Dmetelisproside A, Citroside A, Staphylionoside D [46]; Baimantuoluolines, Baimantuoluoside [47]; Cyclosieversioside F, Astragaloside II, Ginsenoside Rg1, Astrojanoside A, Celerioside E [48]; Isofraxidin, Scopatone, Daturadiol (3),1,4-Benzenediol, Arenarine D, Vanillin, <i>N-trans</i> -Feruloyl-tyramine, Scopoletin, G-Sitosterol and Hyoscyamilactol [49]. β -carotene, carotenoids [50]; β -Phellandrene, γ -Terpinene [51];
Daucus carota	Apiaceae	Eyesight	Roots, Stems, Flower	Italy, Korea	6-methoxymellein [52]; Camphorene, Carotol, β -Bisabolene, Isoelemicin [53].
<i>Drymocallis arguta</i> subsp. arguta	Rosaceae	Diarrhoea Anaemia	NA	NA	NA
Elaeocarpus longifolius Bl.	Elaeocarpaceae		NA	NA	NA

Species	Family	Tenggeresse Ethnopharmacological Uses of Plants	Parts Used for Chemical Isolation	Countries (Chemical Studies Reported)	Isolated Compounds
Erythrina variegata Linnaeus	Leguminoseae	Uses of Plants Diarrhoea	Whole plant Uteaves, Stem	Reported) China Serbia, Italy, Tunisia Turkey, Romania, China, India, Italy, Turkey, Algeria, Italy, Spain, Turkey, and Egypt	Xanthoxyletin [54]; eryvarinols A and B [55]; Protocatechuic acid, Chlorogenic acid, and Caffeic acid [56]; Erythrinin B [57]. Quercetin 3-glucuronide, Isoquercitrin, Rutin, Quercetin 3-arabinoside, Isorhammetin glycosides [58]; Dillapiol, Bergapten, Imperatorin, Psolaren [59]; Anethole, Limonene [60]; Gallic acid, Diosmin, Hesperidin, Kaempferol [61]; Carvacrol, Thymol, Anethol, <i>p</i> -Cymene and γ -Terpinene [62]; (E)-Anethole and <i>p</i> -Acetonylanisole [63]. α -Thujene, 1,&-Cineol, β -Ocimene, Linalool, Germacrene D, Anisketone, Apiol, <i>n</i> -Hexadecanoic acid, Cubebene, Benzene-1-methyl-4-(1-methylethyl)- <i>p</i> -cymene, 1,3,6-Octatriene, 3,7-dimethyl-, (E)-3-carene, 2-Heptene, 3-Methyl-butanal, β -Pinene, Camphene, Hexanal, α -Pinene, β -Phellandrene, α -Phellanrene, β -Myrcene, 4-Carene, 2-Heptanohe, Limonene, 4-Methyl-bicyclo[3.10]hex-2-ene, Eucalyptol, α -Pinene, γ -Terpinene, 7-Dimethyl-1,3,7-octriene, 2,4-Dimethyl-benzenamine, 3-Carene, Cathine, 2-Heptanol, 2-Propyn-1-ol, 2,6-Dimethyl-2,4,6-octatriene, Fenchone, 1-Methyl-4-(1-methylethyl)-benzene, cis-Limonene oxide, fo-Methyl-2-heptanol, Tetradecyl-oxirane, Estragole, Trans- <i>p</i> -2,8-menthadien-1-ol, β -Terpinol, <i>cis-p</i> -2,8-Menthadien, 4-Methyl-1-(methylethyl)-3-cyclohexen, 2-Methyl-3-heptanol, Tetradecyl-oxirane, Estragole, Trans- <i>p</i> -2,8-menthadien-1-ol, β -Terpinol, <i>cis-p</i> -2,8-Menthadien, 4-Methyl-1-(methylethyl)-3-cyclohexen, 2.,3-Cyclohexen-1-methanol, Epi-bicyclosesquiphellardene, <i>cis-p</i> -Menth-2,8-dienol, 1,4-Dimethoxy-benzene, 1-Methoy-4-(1-propenyl)-benzene, 1,2,4a,5,8,8a-Hexadehyde-naphthalene, 4-Methyl-15-(1-methylethyl)-phenol, Mannoheptulose, 2-Methyl-5-(1-methylethyl)-2-cyclohexen-1-oid, 1-Undecanol, Benzothiazole, <i>E</i> -Pinane, 2-Cyclohexen-1-oi, 2-Methyl-5-(1-methylethyl)-phenol, Mannoheptulose, 2-Methyl-5-(1-methylethyl)-2-cyclohexen-1-oid, 1-Undecanol, Benzothiazole, <i>E</i> -Pinane, 2-Cyclohexen-1-oid, 2-Methyl-5-(1-methylethyl)-phenol, Mannoheptulose, 2-Methyl-5-(1-methylethyl)-2-cyclohexen-1-oid, 1-Undecanol, Benzothiazole, <i>E</i> -Pinane, 2
					3-O-β-glucoside [66], quercetin, rutin, isoquercitrin [67], 3-O-caffeoylquinic acid,

Table 1. Cont.

Table 1. Cont.

Species	Family	Tenggeresse Ethnopharmacological Uses of Plants	Parts Used for Chemical Isolation	Countries (Chemical Studies Reported)	Isolated Compounds
<i>Garcinia mangostana</i> Linnaeus	Clusiaceae	Stomach disorder	Fruit	India	 α-Mangostin, β-Mangostin, γ-Mangostin, Garcinone-E, Methoxy-β-mangostin, Xanthone [69]; Mangostin, BR-Xanthone, Gartanin, 8-Desoxygartanin, Garcinone-D, Euxanthone, Xanthione [70]; Epichatechin, and Tannin [71].
Jatropha gossypiifolia Linnaeus	Euphorbiaceae	Rheumatism	Whole plant, Stem, Leaves	India, Nigeria, Thailand	Gossypifan, Gossypilin, Gossypidien [72]; Gadain, Jatroiden [73]; Jatrodien [74]; Arylnaphthalene, Galic, Vanilic, Syringic, 2,5-Dihydroxy benzoic, Caffeic, Rosmarinic, and <i>p</i> -Coumaric [75].
<i>Kaempferia galanga</i> Linnaeus	Zingiberaceae	Rheumatism	Rhizome	Thailand	(–)-Sandaracopimaradiene, Boesenberol, Sandaracopimaradien- 1α , 9α -diol, Kaempulchraol C, Kaempulchraol D [76].
<i>Malus prunifolia</i> (Willd.) Borkh.	Rosaceae	Diarrhoea	Fruit	China	Citric acid, <i>p</i> -Coumaric acid, Hyperoside, Myricetin, Naringenin, Quercetin, Kaempferol, Gentiopicroside, Ursolic acid, and 8-Epiloganic acid [77].
<i>Manihot esculenta</i> Crantz	Euphorbiaceae	Hypertension	Stem	Switzerland, China	Sporoge, Thecacorin, Longifoamide-B (Zeng Y, 2015); Yucalexin P-23, Yucalexin P-15, Protocatechuic acid, and Catalpinic acid [78]; Coniferaldehyde, Isovanillin, 6-Deoxyjacareubin, Scopoletin, Syringaldehyde, Pinoresinol, <i>p</i> -Coumaric acid, Ficusol, Balanophonin and Ethamiyan [79].
Musa paradisiaca Linnaeus	Musaceae	Diarrhoea Stomach disorder	Fruit	Brazil, India	Cycloeucalenone, 31-Norcyclolaudenone, 24-Methylene-cicloartanol [80]; α-Thujene, γ-Terpinene, α- and β-Pinene, Sabinene, β-Myrcene, Limonene, α-Capaene, Caryophyllene and (Z,E)-α Farnesene, Aceteugenol, Palmitic acid, Stearic acid, Palmitin, and Stearin [81]. Momilactones A and B [82]; Momilactone D, Momilactone E,
Oryza satioa Linnaeus	Poaceae	Vitaliser	Seed, Roots	Japan, Korea	Momilactories A and B [62], Momilactorie D, Momilactorie E, Momilactorie A, Sandaracopimaradien-3-one, Oryzalexin A [83]; Oryzativol C [84]; Oryzativol A [85]; ferulic acid, γ -Oryzanol, and Phytic acid [86]; Vanillin, Methyl trans-ferulate, <i>Trans-p</i> -Coumaric acid Methyl ester, <i>N</i> -Benzoyltryptamine, and <i>N</i> -(<i>Trans</i> -cinnamoyl)tryptamine [87].
Persea americana Mill.	Lauraceae	Hypertension	Seed	Brazil	Quercetin and Epicatechin [88]; Avocadene, Avocadyne, Avocadenol-A [89]; γ -Lactone Perseanolide [90].
<i>Physalis lagascae</i> Roem. & Schult.	Solanaceae	Diarrhoea Stomach disorder	NA	NA	NA
Piper amplum Kunth	Piperaceae	Rheumatism	NA	NA	NA

	Table 1	L. Cont.				
Species	Family	Tenggeresse Ethnopharmacological Uses of Plants	Parts Used for Chemical Isolation	Countries (Chemical Studies Reported)	Isolated Compounds	
Piper betle Linnaeus	Piperaceae	Bleeding	Leaves	India, Myanmar, China	Estragole, Linalool, <i>α</i> -Copaene, Anethole, Caryophyllene, <i>α</i> -Terpinene, <i>p</i> -Cymene, 1,8-Cineole, <i>β</i> -Caryophyllene, <i>α</i> -Humulene, Allyl pyrocatechol, Allylcatechol, Methyl eugenol, Estragol (methyl chavicol), Chavibetol, Chavibetol acetate, Safrol, 4-Allyl-2-methoxy-phenolacetate, and 3-Allyl-6-methoxyphenol [91]; Pipeneolignan A, Piperneolignan B, Hydroxychavicol, <i>p</i> -Hydroxycinnamaldehyde, Diallylcatechol [92]; Pipercerebrosides A and B [93]; Piperolactam A [94].	
Rosa tomentosa Sm.	Rosaceae	Fever	NA	NA	NA	
<i>Rubus rosa</i> L. H. Bailey	Rosaceae	Diarrhoea	Whole plant	USA	Elagic acid [95].	
Saccharum officinarum Linnaeus	Poaceae	Rheumatism	Stem	Brazil	Phenolic acid: <i>p</i> -Hydroxybenzoic, <i>p</i> -Hydroxycinnamic, Vanillic and Ferulic acid, Terpenoids: α -Tocopherol and β -Carotene, Flavonoid aglycone Tricin (5,7,4-trihydroxy-3,5-dimethoxyflavone) [96].	
Sechium edule	Cucurbitaceae	Fever (kindern)	Whole plant, Fruit	Mexico	Cinnamic acid, Linoleic, Palmitic, and Myristic acids [97].	
Sesbania grandiflora (L.)Pers.	Fabaceae	Fever	Leaves, Bark, Flowers	Indonesia	Gallic acid [98], 2-Arylbenzofuran [99]; Sesbagrandiflorains A and B [100]; Sesbagrandiflorain D and E, Spinosan A and Spinosan B [101]. Monoterpenes, Glycoalkaloids, and Acyl sugars [102];	
<i>Solanum lycopersicum</i> Linnaeus	Solanaceae	Nutrition	Fruit	USA, Japan, Korea, Chile	13-Oxo-9(Z),11(E),15(Z)-octadecatrienoic acid (13-oxo-OTA), a linolenic acid derivative [103]; Steroidal saponins, Alkaloids, Cerebroside, Phenolic compounds, Sterols, and Nucleosides [104]; Guanosine [105].	
Solanum nigrum Linnaeus	Solanaceae	Hypertension, Tonic drink after hard labour	Whole plant, Fruit, Seed	China, Korea	Lignanamides [106]; Solanine Α, 7α-OH Khasianine, 7α-OH Solamargine and 7α-OH Solasonine [107]; Saponins, Solanigroside A and Solanigroside B [108]; Steroidal glycosides (β2-Solamargine, Solamargine, and Degalactotigonin), Saponin (degalactotigonin) [108 Lunasin [109].	
<i>Tagetes tenuifolia</i> Cavanille	Asteraceae	Nasal bleeding	NA	NA	NA	
<i>Tamarindus indica</i> Linnaeus	Fabaceae	Nausea	Fruit	India	 9,12-Octadecadienoic acid (Z,Z)-, Cis-vaccenic acid, n-Hexadecanoic acid, Beta-Sitosterol, and Octadecanoic acid [110]; Proanthcyanidins (+)-Catechin, Procyanidin B2, (-)-Epicatechin, Procyanidin trimer, Procyanidin tetramer, Procyanidin pentamer, Procyanidin hexamer, Taxifolin, Apigenin, Eriodictyol, Luteolin and Naringenin [111]. Myristicin, Plumbagin, Methyl piperate, 6-Shogaol, 6-Gingerol and Piperine [112]; Geranyl 6-O-α-L-arabinopyranosyl-β-D-glucopyranoside, and Geranyl 6-O-β-D-ayiofuranosyl-β-D-glucopyranoside [113]. 	
Zingiber officinale Roscoe	Zingiberaceae	Headache	Rhizome	Japan, Thailand		

NA: Not available.

3. Biological Activities of Tenggerese Medicinal Plants

To provide a scientific basis to the traditionally claimed therapeutic indications of medicinal plants, it is critical to test the plants for their chemical and biological activities. This is often a challenging task for the traditional practitioners and researchers in Indonesia due to lack of expertise, technology and financial resources. However, since there are many overlapping medicinal plants between different cultures and countries, it is likely that some medicinal plants may have been studied previously. For example, a capsicum species which are used by Tengger people in alleviating post labour complication caused by inflammation [114] has been shown to possess antioxidant and anti-inflammatory activities. In these cases, in order to understand the scientific status of medicinal plants used by Tengger healers, a literature review on each of the plants listed in Table 1 was undertaken for their biological activities. Several medicinal plants have been studied for their biological activities including diarrhoea, wound healing, headache, rheumatism, hypertension, fever, and other disorders. We have discussed them separately.

3.1. Diarrhoea

Many medicinal plants were traditionally used for treating diarrhoea, e.g., the sap of *Musa paradisiaca* L. This plant was reported to possess anti-diarrheal activity in an animal model study [115]. The soluble plantain fibre of banana was also reported to prevent diarrhoea by blocking epithelial adhesion and M-cell translocation of intestinal pathogens [116]. A clinical study on children with acute watery diarrhoea who received green cooked banana supplement indicated a significant recovery of their health [117]. The dietary management of persistent diarrhea in hospitalized children showed that the green banana diet significantly shortened the duration of diarrhea by 18 h compared to the non-banana-supplemented group [118].

3.2. Wound Healing

Of the many plants used by the Tenggerese healers for treating wounds, an in vivo preclinical experiment of leaf extract of Anredera cordifolia in skin burn recovery using albino rats showed a better healing process [119]. This might be related to the antioxidant, anti-inflammatory, and antibacterial properties of the plant. Similarly, a rhizome of Curcuma longa is prepared traditionally in wound healing by the Tenggeresse healers. Its rhizome is rich in curcumin 1, which has been reported for its wound healing, anti-inflammatory, anti-infectious, antibacterial and antioxidant activities [120]. In addition, curcumin 1 advances cutaneous wound healing through tissue remodelling, granulation, tissue formation, collagen deposition and epithelial regeneration, and increases fibroblast proliferation and vascular density [120]. The bulb of Allium sativum (garlic) poultice was applied in wound healing which is rich in allin 2, cycloalliin 3, S-allyl-L-cysteine 4, S-methyl-L-cysteine 5, S-ethylcysteine 6, S-1-proponyl-L-cysteine 7, S-allylmercapto-L-cysteine 8, fructosyl-arginine 9, and β -chlorogenin 10. It also consists of L-arginine 11, L-cysteine 12, and L-methionine 13 (Figure 5) [121]. These compounds were tested to have wound healing activity. Dermatologic application of garlic is correlated with its antioxidant components (S-allyl-L-cysteine 4 and S-allylmercapto-L-cysteine 8), which are organosulfur compounds. In addition, a randomized placebo-controlled double-blinded study on garlic powder revealed the powder increases capillary skin perfusions after 5 h administrations. The pre-clinical trial of aged garlic extract on chicken skin wounds indicated an increase in the re-epithelialization and profuse dose-dependent neovascularization [122].




Figure 5. Wound healing compounds of medicinal plants used by Tenggerese (isolated from the same species found in other countries).

3.3. Headache

Zingiber officinale and Curcuma longa, which are frequently used in Tengger for the treatment of headache, possess several important pharmacological properties including analgesic and neuroprotective properties. A case report of a 42 year old patient with migraine/headache showed that they experienced a reduction in migraine attacks with much lower intensity after consuming ginger powder and using raw fresh ginger in their diet [123]. However, a double-blind placebo-controlled randomized clinical trial of Z. officinale revealed that the consumption of ginger did not have a substantial effect on migraine treatment. Nevertheless, the trial indicated significant activity in attenuating pain intensity [124]. This pain alleviating was associated with the modulatory effect of the trigeminal nociceptor in neurogenic inflammation, and also had neuroprotective effects by inhibition of the production of interleukin 6 (IL-6) and tumor necrosis factor alpha (TNF- α). 6-Gingerol 14 and 6-shogaol 15 (Figure 6) are the main chemical constituents of ginger [125]. Furthermore, a multimodal care for headache, which include C. longa as a management therapy, appeared to improve the patient's symptoms. The tension score of the headache was 3 out of 10 (0 being no pain and 10 being highest pain) in the first week of treatment, with no migraine experienced after that [126]. Curcumin 1 could significantly reduce the neurochemical changes and nerve fibre degeneration [127]. In addition, curcumin 1 (isolated from C. longa) and capsaicin analog [6]-gingerol 14 (isolated from Z. officinale) were reported to possess significant analgesic activities [128,129]. These two plants are commonly used as cooking spices in many parts of the world.






Figure 6. Analgesic compounds of medicinal plants used by Tenggerese (isolated from same species found in other countries).

3.4. Rheumatism and Anti-Inflammatory Agents

Seven medicinal plants prescribed by the Tenggerese healers to treat rheumatism are Alyxia reinwardtii, Borreria laevis, Foeniculum vulgare, Jatropha gossypiifolia, Kaempferia galanga, Saccharum officinarum, and Piper amplumi. Most are prepared as an ointment, juiced, or boiled to drink. Previous studies related to the pharmacological activities of *F. vulgare*, J. gossypiifolia, K. galanga, and S. officinarum supported the traditional claims of these plants as anti-rheumatoid. Locally known as "adas" in Tengger, the leaf of F. vulgare consists of monoterpene hydrocarbon and sesquiterpenes as the main components of their essential oils. The methanol extract of F. vulgare Mill. showed inhibitory effects against acute and subacute inflammatory diseases and possessed a central analgesic effect, validating its traditional use for arthritis [130]. An in vivo preclinical study of *F. vulgare* essential oils against the mouse ear edema model induced by TPA were reported to reduce the level of anti-inflammatory cytokines TNF- α , cyclooxygenase-2 (COX-2), IL-6, and p65 [131]. Additionally, a randomized double-blind trial of women with knee osteoarthritis showed that the extract capsule of *F. vulgare* significantly lowered the scores for pain, disability, total of WOMAC score and VAS variables [132]. In addition, the insignificant toxicity of *F. vulgare* infusion was reported based on an in vivo experiment using rats [133].

The plant *K. galanga* is used traditionally in rheumatism, and has been reported to possess significant anti-inflammatory activity in carrageenan-induced rats by limiting lipoxygenase (LOX), thereby suppressing the leukotriene B4 (LTB4) production [134]. Ethyl-*trans-p*-methoxycinnamate (EPMC) **16** is a dominant phytoconstituent in *K. galanga*. It showed significant anti-inflammatory activity with a minimum inhibitory concentration (MIC) of 100 mg/kg in a carrageenan-induced edema, and also showed non-selective inhibition activities of cyclooxygenases 1 and 2, with IC₅₀ values of 1.12 μ M and 0.83 μ M, respectively [135]. EPMC rich extract suppress acute and chronic inflammation progression in animal models through neutrophil infiltration inhibition [136]. In another recent study, EPMC was also reported to have potential activity to inhibit granuloma tissue formation and suppress cytokine production including IL-1 and TNF- α . The significant analgesic effect of EPMC was also shown in a tail flick experiment of rodents [137].

The herbal gel containing an aqueous extract of J. gossypiifolia was reported to have topical anti-inflammatory activity, either in acute or chronic models of inflammation. It also reduced the production of nitric oxide, leukocyte migration and inhibited edema formation. The flavonoids constituents may be hypothesized as the main active compounds in J. gossypiifolia [138]. In zymosan-induced arthritis mice, a mixture of fatty acids from S. officinarum wax oil (FAM) was reported to decrease the level of β-glucuronidase activity in the synovial fluid of treated mice. FAM also reduces bone erosion [139]. S. officinarum, K. galanga, and F. vulgare, which are used for treating rheumatism pain, have been reported to possess significant anti-inflammatory and analgesic properties when evaluated using a carrageenin-induced test, a hot plate and acetic acid-induced writhing tests [140]. Eight phenolic compounds that were isolated from A.reinwardtii Bl (coumarin 17, 3-hydroxycoumarin 18, 6-hydroxycoumarin 19, 8-hydroxycoumarin 20, scopoletin 21, (+)-pinoresinol 22, zhebeiresinol 23, and p-hydroxybenzoic acid 24) showed anti-inflammatory activities [18] (Figure 7). Brassica rapa [141] and A. reinwardtii [142] were reported to have anti-inflammatory, gastroprotective and antiulcer properties against indomethacin and aspirin-induced rats. In addition, Brassicaceae is composed of an antiinflammatory agent producing veggie species such as *B. oleracea*, which significantly inhibits oxidative/nitrosative stress and lipoperoxidation, based on an ex vivo experiment [143].




Figure 7. Compounds found in plants used by Tenggerese (isolated from same species found in other countries).

3.5. Hypertension

High blood pressure was habitually treated by Tenggerese using *Apium graveolens, Brassica rapa, Manihot esculenta, Persera americana,* and *Solanum nigrum.* The hexane, methanol, and aqueous ethanol extracts of *A. graveolens* seed was reported to reduce blood pressure in deoxycorticosterone acetate–induced hypertensive rats. Further studies revealed that *N*-butylphthalide **25** presented as the major constituent of the hexane extracts of *A. graveolens,* which might be responsible for lowering blood pressure activity. Apigenin **26** isolated from *A. graveolens* demonstrated anti-hypertensive effects in rats [144]. In addition, a randomized triple-blind, placebo-controlled, cross-over clinical trial of *A. graveolens* was reported to have beneficial effects in metabolic syndrome, including hypertension. Administration of *A. graveolens* extract could also alter the pharmacokinetic profile of oral anti-hypertensive drugs when given in combination, thereby enhancing their efficacy [145]. The oil of *P. americana,* commonly known as avocado, was also reported to decrease diastolic and systolic blood pressure by 21.2% and 15.5%, respectively. Besides its beneficial effect on hypertension, avocado oil was reported to suppress the reactive oxygen species (ROS) levels responsible for the pathogenesis of Angiotensin-II induced hypertension [146].

The aqueous extract of *P. americana* leaf showed a significant reduction in systolic blood pressure (SBP), diastolic blood pressure (DBP), and mean arterial pressure (MAP), but had no beneficial effect on heart rate [147]. The crude extracts of other hypotensive medicinal plants such as *B. rapa* and *M. esculenta* inhibited angiotensin I-converting enzyme (ACE) [148]. The dual inhibition of ACE and renin are known to be more effective in lowering blood pressure. A protein-derived glycinyl-histidinyl-serine (GHS) **27** (identified from *B. rapa*) has been known to exhibit dual anti-hypertensive effects [149]. *A. graveolens* extract from which junipediol A 8-O- β -D-glucoside (1- β -D-glucosyloxy-2-(3-methoxy-4-

hydroxyphenyl)propane-1,3-diol **28** was isolated also inhibited the angiotensin-converting enzyme (ACE) [150] (Figure 8).




Figure 8. Antihypertensive compounds of plants used by Tenggerese (isolated from same species reported from other countries).

3.6. Antimicrobial Activities

Some Tenggerese medicinal plants were reported to have potential antimicrobial properties against various strains. For example, Acorus calamus extract, used by Tengerrese traditional healers (dukuns) for treating fever, showed broad-spectrum bioactivities including inhibition of both gram negative and positive bacteria such as *Helicobacter pylori* [151], Propinibacterium acnes [152], Methycillin-resistant staphylococcus aureus (MRSA) [153], Enterobacter aaerogenes, Proteus mirabilis [154] multidrug-resistant enteric bacteria [155], and dengue virus (DENV) replication. In addition, A. calamus. was reported to contain antimicrobial compounds including β -asarone 29, eugenol 30, methyl isoeugenol 31, pinenes 32, myrcene 33, cymene 34 [156] and tatanan A [157]. (E)-Cinnamaldehyde 35, procyanidin B2 36 and (+)-catechin 37 isolated from Cinnamomum burmannii (Nees & T. Nees) Bl. showed antimicrobial activity [158]. Cuminaldehyde 38, β -pinene 39, and γ -terpinene 40 (Figure 9) isolated from Cuminum cyiminum seeds (locally known as 'jinten') showed antibacterial activities against Bacillus cereus, Staphylococcus aureus, and Escherichia coli. The oil from the seed of this plant increases membrane permeability leading to swelling, and the reduction of membrane function, thereby changing cell morphology and causing cell death [159]. Sechium edule was reported to possess antifungal activity against Candida spp and Aspergillus spp. [160]. D. metel showed antifungal activity against Aspergillus flavus, Microsporum canis, and Fusarium solani [161]. In addition, daturalone 41 isolated from Datura metel was reported to be effective against Klebsiella pneumoniae, Bacillus subtilis, Staphylococcus epidermis, and *Staphylococcus aureus* [161].

C. burmanii extract, which contains *E*-cinnamaldehyde **35** and several polyphenols as a predominant volatile oil component, showed antimicrobial activities against *B. cereus, Listeria monocytogenes, S. aureus Escherichia coli*, and *Salmonella anatum* [158]. Crude polar extracts (*n*-butane and ethanol) from *C. burmanii* were reported to be effective against *Listeria monocytogenes, Staphylococcus aureus, Escherichia coli* O157:H7, and *Salmonella anatum* with the inhibition zone ranging from 7.28–24.32 mm in which *n*-butane extract indicated higher activity [27,29]. *Curcuma longa*, which contains curcumin, demonstrated a wide-spectrum of antimicrobial properties against Vibrio harveyi, Vibrio alginolyticus, Vibrio vulnificus, Vibrio parahaemolyticus, Vibrio cholerae, Bacillus subtilis, Bacillus cereus, Aeromonas hydrophila, Streptococcus agalactiae, Staphylococcus aureus, Staphylococcus intermedius, Staphylococcus epidermidis, and Edwardsiella tarda.




Figure 9. Cont.





Figure 9. Antimicrobial compounds of plants used by Tenggerese (isolated from same species grown in other countries).

In addition, compounds isolated from these medicinal plants are known to exhibit various biological activities. For example, curcumin 1 elicited fungal inhibitory activities against four species of Rhizoctonia solani, Phytophthora infestans, Puccinia recondita, and Botrytis cinerea. The curcumin 1 also demonstrated antiviral activities against human immunodeficiency virus (HIV), hepatitis b virus (HBV), hepatitis c virus (HCV), and human papillomavirus (HPV) [162]. The isoflavonoid compounds such as bidwillon B 42, eryvarins V 43, and eryvarins W 44 [163], and orientanol E 45 [164], which were isolated from E. variegata, also demonstrated antibacterial activity against MRSA. Bidwillon B 42, in combination with mupirocin, was effective in eliminating MRSA infection of the nasal cavity and skin [165]. Gallic acid 46 and essential oils present in Sesbania grandiflora [98] and Foeniculum vulgare [166] were responsible for their antibacterial properties, respectively. The F. vulgare essential oil contains trans-anethole 47, fenchone 48, and limonene 49, which were reported to possess potent bioactivities against Mycobacterium tuberculosa, Shigella dysenteriae, Shigella flexneri, Vibrio cholerae, Staphylococcus aureus and Escherichia coli [166]. Essential oils in general are known for their antimicrobial properties and have great applications in making antimicrobial products, lotions, disinfectants and insect repellents (especially mosquitoes) [167]. Secondary metabolites such as dillapiole 50, psoralen 51, bergapten 52, scopoletin 53, imperatorin 54, and dillapional 55 from F. vulgare were reported to be responsible for antibacterial activity [12,59]. Indonesia is also gifted with a diverse array of lichens, which showed potent antibacterial properties-an area worth exploring for chemical and antimicrobial screening [168].

3.7. Antimalarial Activities

There are a number of medicinal plants used by Tenggerese healers for treating fever and malaria. The methanolic extract of the root of *Sesbania* species (used by dukuns for treating fever arising from malaria infection) was reported to have significant antiplasmodial activity, with a minimum inhibition concentration value of 62.5 µg/mL [169]. The dichloromethane extract of *Acorus calamus* was reported to have antiplasmodial activity with an IC₅₀ value of 5.07 µg/mL against the chloroquine-sensitive (CQS) strain of *Plasmodium falciparum* [170]. Further studies showed that curcumin 1 isolated from the roots of *Curcuma longa* inhibited *P. falciparum* growth with an IC₅₀ of ~5 µM. Additionally, in mice infected with *Plasmodium berghei*, oral administration of curcumin 1 was reported to have significant activity in reducing the blood parasitemia by 80–90% [171]. A previous study reported that curcumin 1 was responsible for the inhibition of glycogen synthase Kinase-3 β , which might be contributing to the antimalarial activity [172]. In addition, the moderate anti-malarial activities of *Datura metel* leaf methanol extract were reported with an IC₅₀ value of 22 ± 0.6 µg/mL against *P.falciparum* [173]. The aqueous extract of *Cuminum cyminum* seeds was also reported to have plasmodial growth inhibition by 9% against *P. falciparum* strain FCR3 [174]. From *Erythrina variegata*, Warangalone **56** (8(3,3-dimethyl-allyl)-4'-hydroxy-2''',2'''-dimethylpyran-[6,7,*b*] isoflavone) (Figure 10) had been isolated from stem bark, and possessed antimalarial activity with an IC⁵⁰ value of 4.8 and 3.7 μ g/mL against both the sensitive (3D7) and resistant (K1) strain of *P. falciparum* [175].



Warangalone 56

Figure 10. Antimalarial compound of plants used by Tenggeresse (isolated from same species grown in other countries).

4. Conclusions

This review evaluated 41 medicinal plants used by the indigenous people of the Tengger community, and revealed that 554 phytochemicals have been isolated from 33 plant species with flavonoids and terpenoids as the major chemical components. Most of the plants and their phytochemicals have been tested for various pharmacological activities including anti-inflammatory, antimicrobial, antimalarial, wound healing, headache, and hypertension. Although these medicinal plants grow plentifully in Indonesia, most of the studies were reported on the plants that grow in China, India and Thailand. Only three medicinal plants were phytochemically studied in Indonesia. The research cost and lack of modern laboratory equipment have limited Indonesian researchers in conducting extensive phytochemical and pharmacological studies. The few species that have not been evaluated scientifically presents great potential for biodiscovery. These medicinal plants are: Cayratia clemaidea, Drymocallis arguta, Elaeocarpus longifolius, Physalis lagascae, Piper amplum, Rosa tomentosa and Tagetes tenuifolia. The Cayratia clemaidea, Drymocallis arguta, Elaeocarpus longifolius and Physalis lagascae species has a potential application in treating diarrhea, as this is common among the Indonesian population, especially living in the rural areas where there is lack of food and water sanitation. In addition, Piper amplum, Rosa tomentosa and Tagetes tenuifolia are valuable for bioprospecting to discover new therapeutic agents to treat rheumatism, fever agents and nasal bleeding.

Author Contributions: Conceptualization, A.S.N., A.N.W.P., B.T., P.A.K. and P.W. methodology, A.S.N., A.N.W.P., B.T., P.A.K. and P.W.; software, A.S.N. and A.N.W.P.; validation, A.S.N., A.N.W.P., B.T., P.A.K. and P.W.; formal analysis, A.S.N., A.N.W.P., B.T. and P.W.; investigation, A.S.N., A.N.W.P., B.T., R.P.A., S.M., D.M.R., N.B.W. and P.W.; resources, A.S.N. and P.A.K.; data curation, A.S.N., A.N.W.P., B.T., R.P.A., S.M., D.M.R., N.B.W. and N.B.W.; writing—original draft preparation, A.S.N., A.N.W.P., B.T., R.P.A., S.M., D.M.R., N.B.W., P.A.K. and P.W.; writing—review and editing, A.S.N., D.M.R., N.B.W., P.A.K. and P.W.; writing—review and editing, A.S.N., D.M.R., N.B.W., P.A.K. and P.W.; writing—review and editing, A.S.N., D.M.R., N.B.W., P.A.K. and P.W.; supervision, A.S.N., P.A.K. and P.W.; project administration, A.S.N., B.T. and N.B.W.; funding acquisition, A.S.N., A.N.W.P. and B.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding. The APC was funded by MDPI as an invited paper.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors thank the people of Tengger, the Faculty of Pharmacy for funding, Mohammad Hasan (Vice Chancellor, University of Jember), Sapanyana (Chief of Wonokerso Village)

and Sutomo (Spiritual Leader of the People of Tengger). We thank Hildawati Ilham, Aries Syafitri, Syamsu Dhuha, Putri Sakinah, Virda Fitra Mandasari for technical support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Ekor, M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Front. Pharmacol.* **2014**, *4*, 177. [CrossRef] [PubMed]
- 2. Wangchuk, P.; Tobgay, T. Contributions of medicinal plants to the Gross National Happiness and Biodiscovery in Bhutan. *J. Ethnobiol. Ethnomed.* **2015**, *11*, 48. [CrossRef] [PubMed]
- Yuan, H.; Ma, Q.; Ye, L.; Piao, G. The Traditional Medicine and Modern Medicine from Natural Products. *Molecules* 2016, 21, 559. [CrossRef] [PubMed]
- Al Rashid, M.H.; Kundu, A.; Mandal, V.; Wangchuk, P.; Mandal, S.C. Preclinical and Clinical Trials of Indian Medicinal Plants in Disease Control. In *Herbal Medicine in India: Indigenous Knowledge, Practice, Innovation and Its Value*; Sen, S., Chakraborty, R., Eds.; Springer: Singapore, 2020; pp. 119–142. [CrossRef]
- Yeshi, K.; Gyal, Y.; Sabernig, K.; Phuntsho, J.; Tidwell, T.; Jamtsho, T.; Dhondup, R.; Tokar, E.; Wangchuk, P. An integrated medicine of Bhutan: Sowa Rigpa concepts, botanical identification, and the recorded phytochemical and pharmacological properties of the eastern Himalayan medicinal plants. *Eur. J. Integr. Med.* 2019, 29, 100927. [CrossRef]
- Nugraha, A.S.; Keller, P.a. Revealing indigenous Indonesian traditional medicine: Anti-infective agents. *Nat. Prod. Commun.* 2011, 6, 1953–1966. [CrossRef]
- Roosita, K.; Kusharto, C.M.; Sekiyama, M.; Fachrurozi, Y.; Ohtsuka, R. Medicinal plants used by the villagers of a Sundanese community in West Java, Indonesia. J. Ethnopharmacol. 2008, 115, 72–81. [CrossRef]
- Gewali, M.B.; Awale, S. Aspects of Traditional Medicine in Nepal; Institute of Natural Medicine University of Toyama: Toyama, Japan, 2008.
- 9. Smith-Hefner, N.J. A Social History of Language Change in Highland East Java. J. Asian Stud. 2011, 48, 257–271. [CrossRef]
- 10. Azrianingsih, R.; Kusumahati, A. Perception and appreciation of tenggerese of medicinal plants in Wonokitri Village, Tosari subdistrict, Pasuruan Regency. *AIP Conf. Proc.* **2018**, 2019, 020016. [CrossRef]
- 11. Jadid, N.; Kurniawan, E.; Himayani, C.E.S.; Andriyani; Prasetyowati, I.; Purwani, K.I.; Muslihatin, W.; Hidayati, D.; Tjahjaningrum, I.T.D. An ethnobotanical study of medicinal plants used by the Tengger tribe in Ngadisari village, Indonesia. *PLoS ONE* **2020**, *15*, e0235886. [CrossRef]
- 12. Badgujar, S.A.-O.; Patel, V.V.; Bandivdekar, A.H. Foeniculum vulgare Mill: A review of its botany, phytochemistry, pharmacology, contemporary application, and toxicology. *BioMed Res. Int.* 2014, 2014, 32. [CrossRef]
- 13. Paithankar, V.V.; Belsare, S.L.; Charde, R.M.; Vyas, J.V. Acorus Calamus: An Overview. Int. J. Biomed. Res. 2011, 2, 518–529. [CrossRef]
- 14. Balakumbahan, R.; Rajamani, K.; Kumanan, K. Acorus calamus: An overview. J. Med. Plants Res. 2010, 4, 2740–2745.
- 15. Imam, H.; Riaz, Z.; Azhar, M.; Sofi, G.; Hussain, A. Sweet flag (*Acorus calamus* Linn.): An incredible medicinal herb. *Int. J. Green Pharm.* **2013**, *7*, 288. [CrossRef]
- 16. Hao, Z.-Y.; Liang, D.; Luo, H.; Liu, Y.-F.; Ni, G.; Zhang, Q.-J.; Li, L.; Si, Y.-K.; Sun, H.; Chen, R.-Y.; et al. Bioactive Sesquiterpenoids from the Rhizomes of Acorus calamus. *J. Nat. Prod.* **2012**, *75*, 1083–1089. [CrossRef]
- El-Saber Batiha, G.; Magdy Beshbishy, A.; GWasef, L.; Elewa, Y.H.; AAl-Sagan, A.; Abd El-Hack, M.E.; Taha, A.E.; MAbd-Elhakim, Y.; Prasad Devkota, H. Chemical Constituents and Pharmacological Activities of Garlic (*Allium sativum* L.): A Review. *Nutrients* 2020, 12, 872. [CrossRef] [PubMed]
- 18. Rattanapan, J.; Sichaem, J.; Tip-Pyang, S. Chemical constituents and antioxidant activity from the stems of Alyxia reinwardtii. *Rec. Nat. Prod.* **2012**, *6*, 288–291.
- 19. Souza, L.F.; de Barros, I.B.I.; Mancini, E.; Martino, L.D.; Scandolera, E.; Feo, V.D. Chemical Composition and Biological Activities of the Essential Oil from Anredera cordifolia Grown in Brazil. *Nat. Prod. Commun.* **2014**, *9*, 1934578X1400900730. [CrossRef]
- Liu, G.; Zhuang, L.; Song, D.; Lu, C.; Xu, X. Isolation, purification, and identification of the main phenolic compounds from leaves of celery (*Apium graveolens* L. var. dulce Mill./Pers.). J. Sep. Sci. 2017, 40, 472–479. [CrossRef]
- Hedayati, N.; Bemani Naeini, M.; Mohammadinejad, A.; Mohajeri, S.A. Beneficial effects of celery (*Apium graveolens*) on metabolic syndrome: A review of the existing evidences. *Phytother. Res.* 2019, 33, 3040–3053. [CrossRef]
- 22. Conserva, L.M.; Jesu Costa Ferreira, J. Borreria and Spermacoce species (Rubiaceae): A review of their ethnomedicinal properties, chemical constituents, and biological activities. *Pharmacogn. Rev.* **2012**, *6*, 46. [CrossRef]
- 23. Noiarsa, P.; Yu, Q.; Matsunami, K.; Otsuka, H.; Ruchirawat, S.; Kanchanapoom, T. (Z)-3-Hexenyl diglycosides from Spermacoce laevis Roxb. *J. Nat. Med.* 2007, *61*, 406–409. [CrossRef]
- 24. Fernandes, F.; Valentão, P.; Sousa, C.; Pereira, J.A.; Seabra, R.M.; Andrade, P.B. Chemical and antioxidative assessment of dietary turnip (*Brassica rapa* var. *rapa* L.). *Food Chem.* **2007**, *105*, 1003–1010. [CrossRef]
- Pérez-Vázquez, M.A.K.; Pacheco-Hernández, Y.; Lozoya-Gloria, E.; Mosso-González, C.; Ramírez-García, S.A.; Romero-Arenas, O.; Villa-Ruano, N. Peppermint Essential Oil and Its Major Volatiles as Protective Agents against Soft Rot Caused by *Fusarium sambucinum* in Cera Pepper (*Capsicum pubescens*). *Chem. Biodivers.* 2022, 19, e202100835. [CrossRef] [PubMed]

- Ahmad, I.; Arifianti, A.E.; Sakti, A.S.; Saputri, F.C.; Abdul, M.i. Simultaneous natural deep eutectic solvent-based ultrasonicassisted extraction of bioactive compounds of cinnamon bark and sappan wood as a dipeptidyl peptidase IV inhibitor. *Molecules* 2020, 25, 3832. [CrossRef]
- Liang, Y.; Li, Y.; Sun, A.; Liu, X. Chemical compound identification and antibacterial activity evaluation of cinnamon extracts obtained by subcritical n-butane and ethanol extraction. *Food Sci. Nutr.* 2019, 7, 2186–2193. [CrossRef] [PubMed]
- Muhammad, D.R.A.; Tuenter, E.; Patria, G.D.; Foubert, K.; Pieters, L.; Dewettinck, K. Phytochemical composition and antioxidant activity of *Cinnamomum burmannii* Blume extracts and their potential application in white chocolate. *Food Chem.* 2021, 340, 127983. [CrossRef] [PubMed]
- 29. Rahayu, D.U.; Hakim, R.A.; Mawarni, S.A.; Satriani, A.R. Indonesian Cinnamon (*Cinnamomum burmannii*): Extraction, Flavonoid Content, Antioxidant Activity, and Stability in the Presence of Ascorbic Acid. *Cosmetics* **2022**, *9*, 57. [CrossRef]
- Sethupathy, S.; Nithya, C.; Pandian, S.K. 2-Furaldehyde diethyl acetal from tender coconut water (*Cocos nucifera*) attenuates biofilm formation and quorum sensing-mediated virulence of Chromobacterium violaceum and Pseudomonas aeruginosa. *Biofouling* 2015, 31, 721–733. [CrossRef]
- 31. Elsbaey, M.; Ibrahim, M.A.A.; Bar, F.A.; Elgazar, A.A. Chemical constituents from coconut waste and their in silico evaluation as potential antiviral agents against SARS-CoV-2. *South Afr. J. Bot.* **2021**, *141*, 278–289. [CrossRef]
- Lima, E.B.C.; de Sousa, C.N.S.; Vasconcelos, G.S.; Meneses, L.N.; Silva Pereira, Y.F.E.; Ximenes, N.C.; Santos Júnior, M.A.; Matos, N.C.B.; Brito, R.; Miron, D.; et al. Antidepressant, antioxidant and neurotrophic properties of the standardized extract of Cocos nucifera husk fiber in mice. J. Nat. Med. 2016, 70, 510–521. [CrossRef]
- Zheljazkov, V.D.; Gawde, A.; Cantrell, C.L.; Astatkie, T.; Schlegel, V. Distillation Time as Tool for Improved Antimalarial Activity and Differential Oil Composition of Cumin Seed Oil. *PLoS ONE* 2015, *10*, e0144120. [CrossRef] [PubMed]
- Aldulaimi, O. Screening of Fruits of Seven Plants Indicated for Medicinal Use in Iraq. *Pharmacogn. Mag.* 2017, 13, S189–S195. [CrossRef] [PubMed]
- Neethu, S.; Veena, S.K.; Indulekha, V.C.; Eapen, J.; Radhakrishnan, K.V. Phytoconstituents assessment and development of standardization protocol for 'Nayopayam Kwatha', a polyherbal Ayurvedic formulation. J. Ayurveda Integr. Med. 2021, 12, 489–499. [CrossRef] [PubMed]
- 36. Wuthi-udomlert, M.; Grisanapan, W.; Luanratana, O.; Caichompoo, W. Antifungal activity of Curcuma longa grown in Thailand. *Southeast Asian J. Trop. Med. Public Health* **2000**, *31* (Suppl. 1), 178–182. [PubMed]
- Kim, D.S.; Kim, J.Y. Total synthesis of Calebin-A, preparation of its analogues, and their neuronal cell protectivity against beta-amyloid insult. *Bioorganic Med. Chem. Lett.* 2001, 11, 2541–2543. [CrossRef]
- Aratanechemuge, Y.; Komiya, T.; Moteki, H.; Katsuzaki, H.; Imai, K.; Hibasami, H. Selective induction of apoptosis by arturmerone isolated from turmeric (*Curcuma longa* L.) in two human leukemia cell lines, but not in human stomach cancer cell line. *Int. J. Mol. Med.* 2002, 9, 481–484. [CrossRef]
- 39. Saha, A.; Kuzuhara, T.; Echigo, N.; Suganuma, M.; Fujiki, H. New role of (-)-epicatechin in enhancing the induction of growth inhibition and apoptosis in human lung cancer cells by curcumin. *Cancer Prev. Res.* **2010**, *3*, 953–962. [CrossRef]
- Sun, Y.; Zhang, J.; Zhou, J.; Huang, Z.; Hu, H.; Qiao, M.; Zhao, X.; Chen, D. Synergistic effect of cucurbitacin B in combination with curcumin via enhancing apoptosis induction and reversing multidrug resistance in human hepatoma cells. *Eur. J. Pharmacol.* 2015, 768, 28–40. [CrossRef]
- Wang, Y.; Wang, L.; Zhu, X.; Wang, D.; Li, X. Choleretic Activity of Turmeric and its Active Ingredients. J. Food Sci. 2016, 81, H1800–H1806. [CrossRef]
- Lv, G.-P.; Hu, D.-J.; Zhou, Y.-Q.; Zhang, Q.-W.; Zhao, J.; Li, S.-P. Preparation and Application of Standardized Typical Volatile Components Fraction from Turmeric (*Curcuma longa* L.) by Supercritical Fluid Extraction and Step Molecular Distillation. *Molecules* 2018, 23, 1831. [CrossRef]
- Akter, J.; Islam, M.Z.; Takara, K.; Hossain, M.A.; Sano, A. Isolation and structural elucidation of antifungal compounds from Ryudai gold (*Curcuma longa*) against Fusarium solani sensu lato isolated from American manatee. *Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol.* 2019, 219, 87–94. [CrossRef] [PubMed]
- Le, T.; Beaufay, C.; Nghiem, D.; Pham, T.; Mingeot-Leclercq, M.-P.; Quetin-Leclercq, J. Evaluation of the Anti-Trypanosomal Activity of Vietnamese Essential Oils, with Emphasis on Curcuma longa L. and Its Components. *Molecules* 2019, 24, 1158. [CrossRef] [PubMed]
- Guo, R.; Liu, Y.; Xu, Z.-P.; Xia, Y.-G.; Yang, B.-Y.; Kuang, H.-X. Withanolides from the leaves of *Datura metel L. Phytochemistry* 2018, 155, 136–146. [CrossRef] [PubMed]
- 46. Guo, R.; Liu, Y.; Pan, J.; Guan, W.; Yang, B.-Y.; Kuang, H.-X. A new sesquiterpenoid with cytotoxic and anti-inflammatory activity from the leaves of *Datura metel* L. *Nat. Prod. Res.* **2021**, *35*, 607–613. [CrossRef]
- Liu, Y.; Pan, J.; Sun, Y.-P.; Wang, X.; Liu, Y.; Yang, B.-Y.; Kuang, H.-X. Immunosuppressive withanolides from the flower of *Datura* metel L. Fitoterapia 2020, 141, 104468. [CrossRef] [PubMed]
- 48. Liu, Y.; Wu, D.-D.; Zhou, Y.-Q.; Wu, J.-T.; Qi, Z.-T.; Algradi, A.M.; Pan, J.; Guan, W.; Yang, B.-Y.; Kuang, H.-X. A new ent-kaurane diterpenoid from the pericarps of Datura metel. *J. Asian Nat. Prod. Res.* **2021**, 1–7. [CrossRef]
- Han, X.-L.; Wang, H.; Zhang, Z.-H.; Tan, Y.; Wang, J.-H. Study on Chemical Constituents in Seeds of Datura metel from Xinjiang. Zhong Yao Cai Zhongyaocai J. Chin. Med. Mater. 2015, 38, 1646–1648.

- 50. Miękus, N.; Iqbal, A.; Marszałek, K.; Puchalski, C.; Świergiel, A. Green Chemistry Extractions of Carotenoids from Daucus carota L.-Supercritical Carbon Dioxide and Enzyme-Assisted Methods. *Molecules* **2019**, *24*, 4339. [CrossRef]
- Badalamenti, N.; Modica, A.; Ilardi, V.; Bruno, M.; Maresca, V.; Zanfardino, A.; Di Napoli, M.; Castagliuolo, G.; Varcamonti, M.; Basile, A. *Daucus carota* subsp. *maximus* (Desf.) Ball from Pantelleria, Sicily (Italy): Isolation of essential oils and evaluation of their bioactivity. *Nat. Prod. Res.* 2021, 1, 1–6. [CrossRef]
- 52. Liu, R.; Choi, H.S.; Kim, S.-L.; Kim, J.-H.; Yun, B.-S.; Lee, D.-S. 6-Methoxymellein Isolated from Carrot (Daucus carota L.) Targets Breast Cancer Stem Cells by Regulating NF-κB Signaling. *Molecules* **2020**, *25*, 4374. [CrossRef]
- 53. Gaglio, R.; Barbera, M.; Aleo, A.; Lommatzsch, I.; La Mantia, T.; Settanni, L. Inhibitory Activity and Chemical Characterization of *Daucus carota* subsp. *maximus Essential Oils. Chem. Biodivers.* **2017**, *14*, e1600477. [CrossRef] [PubMed]
- 54. Rasul, A.; Khan, M.; Yu, B.; Ma, T.; Yang, H. Xanthoxyletin, a coumarin induces S phase arrest and apoptosis in human gastric adenocarcinoma SGC-7901 cells. *Asian Pac. J. Cancer Prev. APJCP* **2011**, *12*, 1219–1223. [PubMed]
- 55. Tanaka, H.; Hirata, M.; Etoh, H.; Watanabe, N.; Shimizu, H.; Ahmad, M.; Terada, Y.; Fukai, T. Two diphenylpropan-1,2-diol syringates from the roots of Erythrina variegata. *J. Nat. Prod.* **2002**, *65*, 1933–1935. [CrossRef] [PubMed]
- 56. Liu, Q.; Yu, J.; Liao, X.; Zhang, P.; Chen, X. One-step separation of antioxidant compounds from Erythrina variegata by high speed counter-current chromatography. *J. Chromatogr. Sci.* 2015, *53*, 730–735. [CrossRef]
- Kobayashi, M.; Mahmud, T.; Yoshioka, N.; Shibuya, H.; Kitagawa, I. Indonesian medicinal plants. XXI. Inhibitors of Na+/H+ exchanger from the bark of Erythrina variegata and the roots of Maclura cochinchinensis. *Chem. Pharm. Bull.* 1997, 45, 1615–1619. [CrossRef]
- Kunzemann, J.; Herrmann, K. [Isolation and identification of flavon(ol)-O-glycosides in caraway (*Carum carvi L.*), fennel (*Foeniculum vulgare Mill.*), anise (*Pimpinella anisum L.*), and coriander (*Coriandrum sativum L.*), and of flavon-C-glycosides in anise. I. Phenolics of spices (author's transl)]. Z. Fur Lebensm. Unters. Forsch. 1977, 164, 194–200. [CrossRef]
- 59. Kwon, Y.S.; Choi, W.G.; Kim, W.J.; cKim, W.K.; Kim, M.J.; Kang, W.H.; Kim, C.M. Antimicrobial constituents of foeniculum vulgare. *Arch. Pharmacal. Res.* 2002, 25, 154–157. [CrossRef]
- 60. Senatore, F.; Oliviero, F.; Scandolera, E.; Taglialatela-Scafati, O.; Roscigno, G.; Zaccardelli, M.; De Falco, E. Chemical composition, antimicrobial and antioxidant activities of anethole-rich oil from leaves of selected varieties of fennel [*Foeniculum vulgare* Mill. ssp. *vulgare var. azoricum (Mill.) Thell*]. *Fitoterapia* **2013**, *90*, 214–219. [CrossRef]
- Váradyová, Z.; Pisarčíková, J.; Babják, M.; Hodges, A.; Mravčáková, D.; Kišidayová, S.; Königová, A.; Vadlejch, J.; Várady, M. Ovicidal and larvicidal activity of extracts from medicinal-plants against Haemonchus contortus. *Exp. Parasitol.* 2018, 195, 71–77. [CrossRef]
- Štrbac, F.; Bosco, A.; Maurelli, M.P.; Ratajac, R.; Stojanović, D.; Simin, N.; Orčić, D.; Pušić, I.; Krnjajić, S.; Sotiraki, S.; et al. Anthelmintic Properties of Essential Oils to Control Gastrointestinal Nematodes in Sheep-In Vitro and In Vivo Studies. *Vet. Sci.* 2022, 9, 93. [CrossRef]
- Chiboub, W.; Sassi, A.B.; Amina, C.M.H.; Souilem, F.; El Ayeb, A.; Djlassi, B.; Ascrizzi, R.; Flamini, G.; Harzallah-Skhiri, F. Valorization of the Green Waste from Two Varieties of Fennel and Carrot Cultivated in Tunisia by Identification of the Phytochemical Profile and Evaluation of the Antimicrobial Activities of Their Essentials Oils. *Chem. Biodivers.* 2019, *16*, e1800546. [CrossRef]
- 64. Faudale, M.; Viladomat, F.; Bastida, J.; Poli, F.; Codina, C. Antioxidant Activity and Phenolic Composition of Wild, Edible, and Medicinal Fennel from Different Mediterranean Countries. *J. Agric. Food Chem.* **2008**, *56*, 1912–1920. [CrossRef] [PubMed]
- Parejo, I.; Jauregui, O.; Sánchez-Rabaneda, F.; Viladomat, F.; Bastida, J.; Codina, C. Separation and Characterization of Phenolic Compounds in Fennel (*Foeniculum vulgare*) Using Liquid Chromatography–Negative Electrospray Ionization Tandem Mass Spectrometry. J. Agric. Food Chem. 2004, 52, 3679–3687. [CrossRef] [PubMed]
- 66. Nassar, M.I.; Aboutabl, E.S.A.; Makled, Y.A.; El-Khrisy, E.D.A.; Osman, A.F. Secondary metabolites and pharmacology of *Foeniculum vulgare* Mill. Subsp. *Piperitum. Rev. Latinoam. Química* **2010**, *38*, 10.
- 67. Cherng, J.-M.; Chiang, W.; Chiang, L.-C. Immunomodulatory activities of common vegetables and spices of Umbelliferae and its related coumarins and flavonoids. *Food Chem.* **2008**, *106*, 944–950. [CrossRef]
- Ghanem, M.T.M.; Radwan, H.M.A.; Mahdy, E.S.M.; Elkholy, Y.M.; Hassanein, H.D.; Shahat, A.A. Phenolic Compounds from *Foeniculum vulgare* (Subsp. *Piperitum*) (Apiaceae) Herb and Evaluation of Hepatoprotective Antioxidant Activity. *Pharmacogn. Res.* 2012, 4, 5.
- 69. Dharmaratne, H.R.W.; Piyasena, K.G.N.P.; Tennakoon, S.B. A geranylated biphenyl derivative from *Garcinia malvgostana*. *Nat. Prod. Res.* **2005**, *19*, 239–243. [CrossRef]
- 70. Gopalakrishnan, G.; Banumathi, B.; Suresh, G. Evaluation of the antifungal activity of natural xanthones from *Garcinia mangostana* and their synthetic derivatives. J. Nat. Prod. **1997**, 60, 519–524. [CrossRef]
- Ngawhirunpat, T.; Opanasopi, P.; Sukma, M.; Sittisombut, C.; Kat, A.; Adachi, I. Antioxidant, free radical-scavenging activity and cytotoxicity of different solvent extracts and their phenolic constituents from the fruit hull of mangosteen (*Garcinia mangostana*). *Pharm. Biol.* 2010, 48, 55–62. [CrossRef]
- 72. Das, B.; Anjani, G. ChemInform Abstract: Phytochemicals. Part 29. Gossypidien, a Lignan from Stems of Jatropha gossypifollia. *Cheminform* **2010**, *30*, 1. [CrossRef]
- 73. Oduola, T.; Adeosun, G.; Oduola, T.; Avwioro, G.; Oyeniyi, M. Mechanism of Action of Jatropha Gossypifolia Stem Latex As a Haemostatic Agent. *Eur. J. Gen. Med.* **2004**, *2*, 140–143. [CrossRef]

- 74. Saishri, R.; Ravichandran, N.; Vadivel, V.; Brindha, P. Pharmacognostic studies on leaf of *Jatropha Gossypifolia* L. *Int. J. Pharm. Sci. Res.* **2016**, *7*, 163–173. [CrossRef]
- 75. Povichit, N.; Phrutivorapongkul, A.; Suttajit, M.; Chaiyasut, C.; Leelapornpisid, P. Phenolic content and in vitro inhibitory effects on oxidation and protein glycation of some Thai medicinal plants. *Pak. J. Pharm. Sci.* **2010**, *23*, 403–408. [PubMed]
- 76. Tungcharoen, P.; Chatchai, W.; Tansakul, P.; Nakamura, S.; Matsuda, H.; Tewtrakul, S. Anti-inflammatory effect of isopimarane diterpenoids from Kaempferia galanga. *Phytother. Res.* 2019, 34, 612–623. [CrossRef]
- 77. Wen, C.; Wang, D.; Li, X.; Huang, T.; Huang, C.; Hu, K. Targeted isolation and identification of bioactive compounds lowering cholesterol in the crude extracts of crabapples using UPLC-DAD-MS-SPE/NMR based on pharmacology-guided PLS-DA. *J. Pharm. Biomed. Anal.* **2018**, 150, 144–151. [CrossRef]
- 78. Li, S.-S.; Hu, L.-F.; Zhao, Y.-X.; Zuo, W.-J.; Zeng, Y.-B.; Li, X.-N.; Mei, W.-L.; Dai, H.-F. A new diterpene from the stems of *Manihot* esculenta. J. Asian Nat. Prod. Res. 2011, 13, 961–964. [CrossRef]
- Yi, B.; Hu, L.; Mei, W.; Zhou, K.; Wang, H.; Luo, Y.; Wei, X.; Dai, H. Antioxidant phenolic compounds of cassava (*Manihot esculenta*) from Hainan. *Molecules* 2011, 16, 10157–10167. [CrossRef]
- Silva, A.A.S.; Morais, S.M.; Falcão, M.J.C.; Vieira, I.G.P.; Ribeiro, L.M.; Viana, S.M.; Teixeira, M.J.; Barreto, F.S.; Carvalho, C.A.; Cardoso, R.P.A.; et al. Activity of cycloartane-type triterpenes and sterols isolated from *Musa paradisiaca* fruit peel against *Leishmania infantum* chagasi. *Phytomedicine* 2014, 21, 1419–1423. [CrossRef]
- Fahim, M.; Ibrahim, M.; Zahiruddin, S.; Parveen, R.; Khan, W.; Ahmad, S.; Shrivastava, B.; Shrivastava, A.K. TLC-bioautography identification and GC-MS analysis of antimicrobial and antioxidant active compounds in *Musa* × *paradisiaca* L. fruit pulp essential oil. *Phytochem. Anal. PCA* 2019, 30, 332–345. [CrossRef]
- 82. Quan, N.V.; Tran, H.-D.; Xuan, T.D.; Ahmad, A.; Dat, T.D.; Khanh, T.D.; Teschke, R. Momilactones A and B Are α-Amylase and α-Glucosidase Inhibitors. *Molecules* **2019**, *24*, 482. [CrossRef]
- Cho, J.-G.; Cha, B.-J.; Min Lee, S.; Shrestha, S.; Jeong, R.-H.; Sung Lee, D.; Kim, Y.-C.; Lee, D.-G.; Kang, H.-C.; Kim, J.; et al. Diterpenes from the roots of *Oryza sativa* L. and their inhibition activity on NO production in LPS-stimulated RAW264.7 macrophages. *Chem. Biodivers.* 2015, 12, 1356–1364. [CrossRef] [PubMed]
- 84. Lee, T.K.; Lee, D.; Yu, J.S.; Jo, M.S.; Baek, S.C.; Shin, M.-S.; Ko, Y.-J.; Kang, K.S.; Kim, K.H. Biological Evaluation of a New Lignan from the Roots of Rice (*Oryza sativa*). *Chem. Biodivers*. **2018**, *15*, e1800333. [CrossRef] [PubMed]
- Kang, H.R.; Yun, H.S.; Lee, T.K.; Lee, S.; Kim, S.-H.; Moon, E.; Park, K.-M.; Kim, K.H. Chemical Characterization of Novel Natural Products from the Roots of Asian Rice (*Oryza sativa*) that Control Adipocyte and Osteoblast Differentiation. *J. Agric. Food Chem.* 2018, 66, 2677–2684. [CrossRef]
- 86. Manosroi, A.; Chutoprapat, R.; Abe, M.; Manosroi, W.; Manosroi, J. Anti-aging efficacy of topical formulations containing niosomes entrapped with rice bran bioactive compounds. *Pharm. Biol.* **2012**, *50*, 208–224. [CrossRef]
- Cho, J.-G.; Huh, J.; Jeong, R.-H.; Cha, B.-J.; Shrestha, S.; Lee, D.-G.; Kang, H.-C.; Kim, J.-Y.; Baek, N.-I. Inhibition effect of phenyl compounds from the *Oryza sativa* roots on melanin production in murine B16-F10 melanoma cells. *Nat. Prod. Res.* 2015, 29, 1052–1054. [CrossRef] [PubMed]
- Soldera-Silva, A.; Seyfried, M.; Campestrini, L.H.; Zawadzki-Baggio, S.F.; Minho, A.P.; Molento, M.B.; Maurer, J.B.B. Assessment of anthelmintic activity and bio-guided chemical analysis of *Persea americana* seed extracts. *Vet. Parasitol.* 2018, 251, 34–43. [CrossRef]
- Louis, M.R.L.M.; Rani, V.P.; Krishnan, P.; Reegan, A.D.; Balakrishna, K.; Ignacimuthu, S.; Packiam, S.M.; Maheswaran, R.; Shirota, O. Mosquito Larvicidal Activity of Compounds from Unripe Fruit Peel of Avocado (*Persea americana* Mill.). *Appl. Biochem. Biotechnol.* 2022, 1, 1–12. [CrossRef]
- 90. Reis, I.M.A.; Umehara, E.; Conceição, R.S.; de M. Oliveira, L.; Coelho Dos, S.M., Jr.; Costa-Silva, T.A.; Amaral, M.; Tempone, A.G.; Branco, A.; Lago, J.H.G. γ-Lactones from Persea americana and Persea fulva—In Vitro and in Silico Evaluation of Trypanosoma cruzi Activity. *Chem. Biodivers.* 2021, 18, e2100362. [CrossRef]
- Nayaka, N.M.D.M.W.; Sasadara, M.M.V.; Sanjaya, D.A.; Yuda, P.E.S.K.; Dewi, N.L.K.A.A.; Cahyaningsih, E.; Hartati, R. Piper betle (L): Recent Review of Antibacterial and Antifungal Properties, Safety Profiles, and Commercial Applications. *Molecules* 2021, 26, 2321. [CrossRef]
- 92. San, T.T.; Wang, Y.-H.; Hu, D.-B.; Yang, J.; Zhang, D.-D.; Xia, M.-Y.; Yang, X.-F.; Yang, Y.-P. A new sesquineolignan and four new neolignans isolated from the leaves of *Piper betle*, a traditional medicinal plant in Myanmar. *Bioorganic Med. Chem. Lett.* **2021**, *31*, 127682. [CrossRef]
- Chen, D.-Z.; Xiong, H.-B.; Tian, K.; Guo, J.-M.; Huang, X.-Z.; Jiang, Z.-Y. Two new sphingolipids from the leaves of *Piper betle* L. *Molecules* 2013, 18, 11241–11249. [CrossRef] [PubMed]
- 94. Amin, S.A.; Bhattacharya, P.; Basak, S.; Gayen, S.; Nandy, A.; Saha, A. Pharmacoinformatics study of Piperolactam A from *Piper betle* root as new lead for non steroidal anti fertility drug development. *Comput. Biol. Chem.* **2017**, *67*, 213–224. [CrossRef]
- Miyasaki, Y.; Rabenstein, J.D.; Rhea, J.; Crouch, M.-L.; Mocek, U.M.; Kittell, P.E.; Morgan, M.A.; Nichols, W.S.; Van Benschoten, M.M.; Hardy, W.D.; et al. Isolation and characterization of antimicrobial compounds in plant extracts against multidrug-resistant *Acinetobacter baumannii*. *PLoS ONE* 2013, *8*, e61594. [CrossRef] [PubMed]
- 96. Alves, V.G.; Souza, A.G.; Chiavelli, L.U.R.; Ruiz, A.L.T.G.; Carvalho, J.E.; Pomini, A.M.; Silva, C.C. Phenolic compounds and anticancer activity of commercial sugarcane cultivated in Brazil. *An. Acad. Bras. Cienc.* **2016**, *88*, 1201–1209. [CrossRef] [PubMed]

- Trejo-Moreno, C.; Castro-Martínez, G.; Méndez-Martínez, M.; Jiménez-Ferrer, J.E.; Pedraza-Chaverri, J.; Arrellín, G.; Zamilpa, A.; Medina-Campos, O.N.; Lombardo-Earl, G.; Barrita-Cruz, G.J.; et al. Acetone fraction from *Sechium edule* (Jacq.) S.w. edible roots exhibits anti-endothelial dysfunction activity. *J. Ethnopharmacol.* 2018, 220, 75–86. [CrossRef]
- Anantaworasakul, P.; Hamamoto, H.; Sekimizu, K.; Okonogi, S. Biological activities and antibacterial biomarker of *Sesbania* grandiflora bark extract. Drug Discov. Ther. 2017, 11, 70–77. [CrossRef]
- 99. Noviany, N.; Nurhidayat, A.; Hadi, S.; Suhartati, T.; Aziz, M.; Purwitasari, N.; Subasman, I. Sesbagrandiflorain A and B: Isolation of two new 2-arylbenzofurans from the stem bark of *Sesbania grandiflora*. *Nat. Prod. Res.* **2018**, *32*, 2558–2564. [CrossRef]
- 100. Noviany, N.; Samadi, A.; Carpenter, E.L.; Abugrain, M.E.; Hadi, S.; Purwitasari, N.; Indra, G.; Indra, A.; Mahmud, T. Structural revision of sesbagrandiflorains A and B, and synthesis and biological evaluation of 6-methoxy-2-arylbenzofuran derivatives. *J. Nat. Med.* 2020, 75, 66–75. [CrossRef]
- 101. Tjahjandarie, T.S.; Tanjung, M.; Saputri, R.D.; Rahayu, D.O.; Gunawan Alfiah Nur, I.; Aldin, M.F. Two new 2-arylbenzofurans from *Sesbania grandiflora* L. and their cytotoxicity towards cancer cell. *Nat. Prod. Res.* **2020**, *35*, 5637–5642. [CrossRef]
- 102. Kang, J.-H.; Shi, F.; Jones, A.D.; Marks, M.D.; Howe, G.A. Distortion of trichome morphology by the hairless mutation of tomato affects leaf surface chemistry. *J. Exp. Bot.* 2009, *61*, 1053–1064. [CrossRef]
- 103. Takahashi, H.; Hara, H.; Goto, T.; Kamakari, K.; Wataru, N.; Mohri, S.; Suzuki, H.; Shibata, D.; Kawada, T. 13-Oxo-9(Z),11(E),15(Z)octadecatrienoic Acid Activates Peroxisome Proliferator-Activated Receptor γ in Adipocytes. *Lipids* 2014, 50, 3–12. [CrossRef] [PubMed]
- 104. Seong, S.H.; Jung, H.A.; Choi, J.S. Discovery of Flazin, an Alkaloid Isolated from Cherry Tomato Juice, As a Novel Non-Enzymatic Protein Glycation Inhibitor via in Vitro and in Silico Studies. J. Agric. Food Chem. 2021, 69, 3647–3657. [CrossRef] [PubMed]
- 105. Fuentes, E.; Alarcón, M.; Astudillo, L.; Valenzuela, C.; Gutiérrez, M.; Palomo, I. Protective mechanisms of guanosine from *Solanum lycopersicum* on agonist-induced platelet activation: Role of sCD40L. *Molecules* 2013, 18, 8120–8135. [CrossRef] [PubMed]
- 106. Li, C.-X.; Song, X.-Y.; Zhao, W.-Y.; Yao, G.-D.; Lin, B.; Huang, X.-X.; Li, L.-Z.; Song, S.-J. Characterization of enantiomeric lignanamides from *Solanum nigrum* L. and their neuroprotective effects against MPP(+)-induced SH-SY5Y cells injury. *Phytochemistry* 2019, 161, 163–171. [CrossRef]
- Gu, X.-Y.; Shen, X.-F.; Wang, L.; Wu, Z.-W.; Li, F.; Chen, B.; Zhang, G.-L.; Wang, M.-K. Bioactive steroidal alkaloids from the fruits of *Solanum nigrum*. *Phytochemistry* 2018, 147, 125–131. [CrossRef]
- 108. Zhou, X.-L.; He, X.-J.; Zhou, G.-X.; Ye, W.-C.; Yao, X.-S. Pregnane glycosides from Solanum nigrum. J. Asian Nat. Prod. Res. 2007, 9, 517–523. [CrossRef]
- 109. Jeong, J.B.; De Lumen, B.O.; Jeong, H.J. Lunasin peptide purified from *Solanum nigrum* L. protects DNA from oxidative damage by suppressing the generation of hydroxyl radical via blocking fenton reaction. *Cancer Lett.* **2010**, *293*, 58–64. [CrossRef]
- Fagbemi, K.O.; Aina, D.A.; Olajuyigbe, O.O. Soxhlet Extraction versus Hydrodistillation Using the Clevenger Apparatus: A Comparative Study on the Extraction of a Volatile Compound from Tamarindus indica Seeds. *Sci. World J.* 2021, 2021, 5961586. [CrossRef]
- 111. Sudjaroen, Y.; Haubner, R.; Würtele, G.; Hull, W.E.; Erben, G.; Spiegelhalder, B.; Changbumrung, S.; Bartsch, H.; Owen, R.W. Isolation and structure elucidation of phenolic antioxidants from Tamarind (*Tamarindus indica* L.) seeds and pericarp. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* 2005, 43, 1673–1682. [CrossRef]
- 112. Rattarom, R.; Sakpakdeejaroen, I.; Hansakul, P.; Itharat, A. Cytotoxic activity against small cell lung cancer cell line and chromatographic fingerprinting of six isolated compounds from the ethanolic extract of Benjakul. *J. Med. Assoc. Thail. Chotmaihet Thangphaet* **2014**, *97* (Suppl. 8), S70-5.
- 113. Sekiwa, Y.; Kobayashi, A.; Kubota, K.; Takenaka, M. First isolation of geranyl disaccharides from ginger and their relations to aroma formation. *Nat. Prod. Lett.* **2001**, *15*, 267–274. [CrossRef] [PubMed]
- 114. Della Valle, A.; Dimmito, M.P.; Zengin, G.; Pieretti, S.; Mollica, A.; Locatelli, M.; Cichelli, A.; Novellino, E.; Ak, G.; Yerlikaya, S.; et al. Exploring the Nutraceutical Potential of Dried Pepper Capsicum annuum L. on Market from Altino in Abruzzo Region. *Antioxidants* 2020, 9, 400. [CrossRef] [PubMed]
- 115. Yakubu, M.T.; Nurudeen, Q.O.; Salimon, S.S.; Yakubu, M.O.; Jimoh, R.O.; Nafiu, M.O.; Akanji, M.A.; Oladiji, A.T.; Williams, F.E. Antidiarrhoeal Activity of *Musa paradisiaca* Sap in Wistar Rats. *Evid.-Based Complement. Altern. Med. Ecam* 2015, *4*, 1–9. [CrossRef]
- 116. Roberts, C.L.; Keita, Å.V.; Parsons, B.N.; Prorok-Hamon, M.; Knight, P.; Kennedy, N.; Söderholm, J.D.; Rhodes, J.M.; Campbell, B.J. Soluble plantain fibre blocks epithelial adhesion and M-cell translocation of intestinal pathogens. *Gut* **2011**, *60*, A96. [CrossRef]
- 117. Gunasekaran, D.; Chandramohan, A.; Karthikeyan, K.; Balasubramaniam, B.; Jagadeesan, P.; Soundararajan, P. Effect of Green Banana (*Musa paradisiaca*) on Recovery in Children With Acute Watery Diarrhea With No Dehydration: A Randomized Controlled Trial. *Indian Pediatr.* 2020, *57*, 1114–1118. [CrossRef] [PubMed]
- 118. Alvarez-Acosta, T.; León, C.; Acosta-González, S.; Parra-Soto, H.; Cluet-Rodriguez, I.; Rossell, M.R.; Colina-Chourio, J.A. Beneficial Role of Green Plantain [*Musa paradisiaca*] in the Management of Persistent Diarrhea: A Prospective Randomized Trial. *J. Am. Coll. Nutr.* 2009, 28, 169–176. [CrossRef]
- 119. Yuniarti, W.M.; Lukiswanto, B.S. Effects of herbal ointment containing the leaf extracts of Madeira vine (*Anredera cordifolia* (Ten.) Steenis) for burn wound healing process on albino rats. *Vet. World* 2017, *10*, 808–813. [CrossRef]
- 120. Akbik, D.; Ghadiri, M.; Chrzanowski, W.; Rohanizadeh, R. Curcumin as a wound healing agent. Life Sci. 2014, 116, 1–7. [CrossRef]
- 121. Pazyar, N.; Feily, A. Garlic in dermatology. Dermatol. Rep. 2011, 3, e4. [CrossRef]

- 122. Ejaz, S.; Chekarova, I.; Cho, J.; Lee, S.; Ashraf, S.; Lim, C. Effect of aged garlic extract on wound healing: A new frontier in wound management. *Drug Chem. Toxicol.* **2009**, *32*, 191–203. [CrossRef]
- 123. Mustafa, T.; Srivastava, K.C. Ginger (Zingiber officinale) in migraine headache. J. Ethnopharmacol. 1990, 29, 267–273. [CrossRef]
- 124. Martins, L.B.; Rodrigues, A.M.d.S.; Monteze, N.M.; Tibaes, J.R.B.; Amaral, M.H.A.; Gomez, R.S.; Teixeira, A.L.; Ferreira, A.V.M. Double-blind placebo-controlled randomized clinical trial of ginger (*Zingiber officinale Rosc.*) in the prophylactic treatment of migraine. *Cephalalgia* 2019, 40, 88–95. [CrossRef] [PubMed]
- 125. Kim, Y.J.; Lim, H.-S.; Kim, B.-Y.; Seo, C.-S.; Jeong, S.-J. Quantitative Analysis and Biological Efficacies regarding the Neuroprotective and Antineuroinflammatory Actions of the Herbal Formula Jodeungsan in HT22 Hippocampal Cells and BV-2 Microglia. *Evid.-Based Complement. Altern. Med.* 2017, 2017, 6360836. [CrossRef] [PubMed]
- 126. Martin, B.R. Multimodal Care for Headaches, Lumbopelvic Pain, and Dysmenorrhea in a Woman With Endometriosis: A Case Report. J. Chiropr. Med. 2021, 20, 148–157. [CrossRef]
- 127. Liu, Z.; Huang, P.; Law, S.; Tian, H.; Leung, W.; Xu, C. Preventive Effect of Curcumin Against Chemotherapy-Induced Side-Effects. *Front. Pharmacol.* **2018**, *9*, 1374. [CrossRef]
- 128. Mittal, N.; Joshi, R.; Hota, D.; Chakrabarti, A. Evaluation of antihyperalgesic effect of curcumin on formalin-induced orofacial pain in rat. *Phytother. Res.* 2009, 23, 507–512. [CrossRef]
- 129. Young, H.-Y.; Luo, Y.-L.; Cheng, H.-Y.; Hsieh, W.-C.; Liao, J.-C.; Peng, W.-H. Analgesic and anti-inflammatory activities of [6]-gingerol. J. Ethnopharmacol. 2005, 96, 207–210. [CrossRef]
- 130. Choi, E.M.; Hwang, J.K. Antiinflammatory, analgesic and antioxidant activities of the fruit of Foeniculum vulgare. *Fitoterapia* **2004**, *75*, 557–565. [CrossRef]
- Li, C.; Cai, Q.; Wu, X.; Tan, Z.; Yao, L.; Huang, S.; Zhang, W.; Hong, Z.; Chen, Z.; Zhang, L.A.-O. Anti-inflammatory Study on the Constituents of *Angelica sinensis* (Oliv.) Diels, *Angelica dahurica* (Hoffm.) Benth. & Hook.f. ex Franch. & Sav., *Angelica pubescence* Maxim and *Foeniculum vulgare* Mill. J. Essent. Oils 2022, 71, 1207–1219.
- 132. Alazadeh, M.; Azadbakht, M.; Niksolat, F.; Asgarirad, H.; Moosazadeh, M.; Ahmadi, A.; Yousefi, S.S. Effect of sweet fennel seed extract capsule on knee pain in women with knee osteoarthritis. *Complement. Ther. Clin. Pract.* **2020**, *40*, 101219. [CrossRef]
- 133. Rifqiyati, N.; Wahyuni, A. Fennel (*Foeniculum vulgare*) leaf infusion effect on mammary gland activity and kidney function of lactating rats. *Nusant. Biosci.* 2019, *11*, 5. [CrossRef]
- 134. Dwita, L.P.; Hikmawanti, N.P.E.; Yeni; Supandi. Extract, fractions, and ethyl-p-methoxycinnamate isolate from Kaempferia galanga Elicit anti-inflammatory activity by limiting leukotriene B4 (LTB4) production. J. Tradit. Complement. Med. 2021, 11, 563–569. [CrossRef] [PubMed]
- Umar, M.I.; Asmawi, M.Z.; Sadikun, A.; Atangwho, I.J.; Yam, M.F.; Altaf, R.; Ahmed, A. Bioactivity-guided isolation of ethylp-methoxycinnamate, an anti-inflammatory constituent, from *Kaempferia galanga* L. extracts. *Molecules* 2012, 17, 8720–8734. [CrossRef] [PubMed]
- 136. Jagadish, P.C.; Latha, K.P.; Mudgal, J.; Nampurath, G.K. Extraction, characterization and evaluation of Kaempferia galanga L. (Zingiberaceae) rhizome extracts against acute and chronic inflammation in rats. J. Ethnopharmacol. 2016, 194, 434–439. [CrossRef] [PubMed]
- 137. Muhammad Ihtisham, U.; Mohd Zaini, A.; Amirin, S.; Amin Malik Shah Abdul, M.; Fouad Saleih, R.A.-S.; Loiy Elsir Ahmed, H.; Rabia, A.; Mohamed, B.K.A. Ethyl-p-methoxycinnamate isolated from kaempferia galanga inhibits inflammation by suppressing interleukin-1, tumor necrosis factor-α, and angiogenesis by blocking endothelial functions. *Clinics* 2014, 69, 134–144. [CrossRef]
- 138. Xavier-Santos, J.B.; Félix-Silva, J.; Passos, J.G.R.; Gomes, J.A.S.; Fernandes, J.M.; Garcia, V.B.; de Araujo-Junior, R.F.; Zucolotto, S.M.; Silva-Junior, A.A.; Fernandes-Pedrosa, M.F. Development of an effective and safe topical anti-inflammatory gel containing Jatropha gossypiifolia leaf extract: Results from a pre-clinical trial in mice. *J. Ethnopharmacol.* 2018, 227, 268–278. [CrossRef]
- 139. Ledon, N.; Casaco, A.; Remirez, D.; González, A.; Cruz, J.; Gonzalez, R.; Capote, A.; Tolón, Z.; Rojas, E.; Rodríguez, V.J.; et al. Effects of a mixture of fatty acids from sugar cane (*Saccharum officinarum* L.) wax oil in two models of inflammation: Zymosaninduced arthritis and mice tail test of psoriasis. *Phytomedicine Int. J. Phytother. Phytopharm.* 2007, 14, 690–695. [CrossRef]
- 140. Ledon, N. Anti-inflammatory and analgesic effects of a mixture of fatty acids isolated and purified from sugar cane wax oil. *Planta Med.* **2003**, 2069, 367–369. [CrossRef]
- 141. Alotaibi, B.; Mokhtar, F.A.-O.; El-Masry, T.A.; Elekhnawy, E.; Mostafa, S.A.; Abdelkader, D.H.; Elharty, M.E.; Saleh, A.; Negm, W.A.-O. Antimicrobial Activity of *Brassica rapa* L. Flowers Extract on Gastrointestinal Tract Infections and Antiulcer Potential Against Indomethacin-Induced Gastric Ulcer in Rats Supported by Metabolomics Profiling. *J. Inflamm. Res.* 2021, 14, 7411. [CrossRef]
- 142. Nugroho, A.E.; Wijayanti, A.D.; Mutmainah, M.; Susilowati, R.; Rahmawati, N. Gastroprotective Effect of Combination of Hot Water Extracts of Licorice (*Glycyrrhiza glabra*), Pulasari Stem Bark (*Alyxia reinwardtii*), and Sembung Leaf (*Blumea balsamifera*) Against Aspirin-Induced Gastric Ulcer Model Rats. J. Evid. Based Complement. Altern. Med. 2016, 21, NP77–NP84. [CrossRef]
- 143. Mollica, A.; Stefanucci, A.; Zengin, G.; Locatelli, M.; Macedonio, G.; Orlando, G.; Ferrante, C.; Menghini, L.; Recinella, L.; Leone, S.; et al. Polyphenolic composition, enzyme inhibitory effects ex-vivo and in-vivo studies on two Brassicaceae of north-central Italy. *Biomed. Pharmacother.* 2018, 107, 129–138. [CrossRef] [PubMed]
- 144. Chauhan, N. Plants Used as Antihypertensive. Nat. Prod. Bioprospect. 2020, 11, 155–184. [CrossRef]
- 145. Siska, S.; Mun'im, A.; Bahtiar, A.; Suyatna, F.D. Effect of Apium graveolens Extract Administration on the Pharmacokinetics of Captopril in the Plasma of Rats. Sci. Pharm. 2018, 86, 6. [CrossRef] [PubMed]

- 146. Márquez-Ramírez, C.; Paz, J.; Ortiz-Avila, O.; Raya-Farias, A.; González-Hernández, J.; Rodríguez-Orozco, A.; Salgado-Garciglia, R.; Saavedra-Molina, A.; Godínez-Hernández, D.; Cortés-Rojo, C. Comparative effects of avocado oil and losartan on blood pressure, renal vascular function and mitochondrial oxidative stress in hypertensive rats. *Nutrition* 2018, 54, 60–67. [CrossRef]
- 147. Sokpe, A.; Mensah, M.; Koffuor, G.; Thomford, K.P.; Arthur, R.; Jibira, Y.; Baah, M.; Adedi, B.; Agbemenyah, H. Hypotensive and Antihypertensive Properties and Safety for Use of Annona muricata and Persea americana and Their Combination Products. *Evid. Based Complement. Altern. Med.* 2020, *6*, 1–13. [CrossRef] [PubMed]
- 148. Rong, H.; Alashi, A.; Malomo, S.; Girgih, A.; Chao, D.; Ju, X.; Aluko, R. Antihypertensive and free radical scavenging properties of enzymatic rapeseed protein hydrolysates. *Food Chem.* **2013**, *141*, 153–159. [CrossRef]
- 149. He, R.; Malomo, S.A.; Girgih, A.T.; Ju, X.; Aluko, R.E. Glycinyl-Histidinyl-Serine (GHS), a Novel Rapeseed Protein-Derived Peptide Has Blood Pressure-Lowering Effect in Spontaneously Hypertensive Rats. J. Agric. Food Chem. 2013, 61, 8396–8402. [CrossRef]
- 150. Simaratanamongkol, A.; Umehara, K.; Noguchi, H.; Panichayupakaranant, P. Identification of a new angiotensin-converting enzyme (ACE) inhibitor from Thai edible plants. *Food Chem.* **2014**, *165*, 92–97. [CrossRef]
- 151. Prasad, A.; Devi, A.T.; Prasad, M.N.N.; Zameer, F.; Shruthi, G.; Shivamallu, C. Phyto anti-biofilm elicitors as potential inhibitors of Helicobacter pylori. *3 Biotech* **2019**, *9*, 53. [CrossRef]
- 152. Kim, W.-J.; Hwang, K.-H.; Park, D.-G.; Kim, T.-J.; Kim, D.-W.; Choi, D.-K.; Moon, W.-K.; Lee, K.-H. Major constituents and antimicrobial activity of Korean herb Acorus calamus. *Nat. Prod. Res.* 2011, 25, 1278–1281. [CrossRef]
- Aqil, F.; Ahmad, I.; Owais, M. Evaluation of anti-methicillin-resistantStaphylococcus aureus (MRSA) activity and synergy of some bioactive plant extracts. *Biotechnol. J.* 2006, 1, 1093–1102. [CrossRef] [PubMed]
- Mickymaray, S.; Al Aboody, M.S. In Vitro Antioxidant and Bactericidal Efficacy of 15 Common Spices: Novel Therapeutics for Urinary Tract Infections? *Medicina* 2019, 55, 289. [CrossRef] [PubMed]
- 155. Ahmad, I.; Aqil, F. In vitro efficacy of bioactive extracts of 15 medicinal plants against ESβL-producing multidrug-resistant enteric bacteria. *Microbiol. Res.* 2007, 162, 264–275. [CrossRef]
- 156. Sharma, V.; Singh, I.; Chaudhary, P. Acorus calamus (The Healing Plant): A review on its medicinal potential, micropropagation and conservation. *Nat. Prod. Res.* 2014, 28, 1454–1466. [CrossRef]
- 157. Yao, X.; Ling, Y.; Guo, S.; He, S.; Wu, W.; Zhang, Q.; Zou, M.; Nandakumar, K.S.; Chen, X.; Liu, S. Tatanan A from the *Acorus calamus* L. root inhibited dengue virus proliferation and infections. *Phytomedicine* **2018**, *42*, 258–267. [CrossRef]
- Shan, B.; Cai Yz Fau-Brooks, J.D.; Brooks Jd Fau-Corke, H.; Corke, H. Antibacterial properties and major bioactive components of cinnamon stick (*Cinnamomum burmannii*): Activity against foodborne pathogenic bacteria. J. Agric. Food Chem. 2007, 55, 5484–5490. [CrossRef] [PubMed]
- Wongkattiya, N.; Sanguansermsri, P.; Fraser, I.H.; Sanguansermsri, D.A.-O. Antibacterial activity of cuminaldehyde on food-borne pathogens, the bioactive component of essential oil from *Cuminum cyminum* L. collected in Thailand. *J. Complement. Integr. Med.* 2019, *16*, 1. [CrossRef]
- Ordoñez, A.A.; Ordoñez Rm Fau-Zampini, I.C.; Zampini Ic Fau-Isla, M.I.; Isla, M.I. Design and quality control of a pharmaceutical formulation containing natural products with antibacterial, antifungal and antioxidant properties. *Int. J. Pharm.* 2009, 378, 51–58. [CrossRef]
- 161. Bawazeer, S.; Rauf, A. In Vitro Antibacterial and Antifungal Potential of Amyrin-Type Triterpenoid Isolated from Datura metel Linnaeus. *BioMed. Res. Int.* 2021, 2021, 1543574. [CrossRef]
- 162. Zorofchian, S.; Kadir, H.; Hassandarvish, P.; Tajik, H.; Abu Bakar, S.; Zandi, K. A Review on Antibacterial, Antiviral, and Antifungal Activity of Curcumin. *BioMed. Res. Int.* 2014, 2014, 186864. [CrossRef]
- 163. Tanaka, H.; Atsumi, I.; Shirota, O.; Sekita, S.; Sakai, E.; Sato, M.; Murata, J.; Murata, H.; Darnaedi, D.; Chen, I.-S. Three new constituents from the roots of *Erythrina variegata* and their antibacterial activity against methicillin-resistant *Staphylococcus aureus*. *Chem. Biodivers.* 2011, *8*, 476–482. [CrossRef] [PubMed]
- 164. Tanaka, H.; Atsumi, I.; Hasegawa, M.; Hirata, M.; Sakai, T.; Sato, M.; Yamaguchi, R.; Tateishi, Y.; Tanaka, T.; Fukai, T. Two New Isoflavanones from the Roots of *Erythrina variegata*. *Nat. Prod. Commun.* **2015**, *10*, 1934578X1501000330. [CrossRef]
- 165. Sato, M.; Tanaka, H.; Yamaguchi, R.; Kato, K.; Etoh, H. Synergistic effects of mupirocin and an isoflavanone isolated from Erythrina variegata on growth and recovery of methicillin-resistant *Staphylococcus aureus*. Int. J. Antimicrob. Agents 2004, 24, 241–246. [CrossRef] [PubMed]
- 166. Camacho-Corona, M.D.R.; Ramírez-Cabrera, M.A.; Santiago, O.G.; Garza-González, E.; Palacios, I.d.P.; Luna-Herrera, J. Activity against drug resistant-tuberculosis strains of plants used in Mexican traditional medicine to treat tuberculosis and other respiratory diseases. *Phytother. Res. PTR* 2008, 22, 82–85. [CrossRef]
- Yeshi, K.; Wangchuk, P. Chapter 11—Essential oils and their bioactive molecules in healthcare. In *Herbal Biomolecules in Healthcare Applications*; Mandal, S.C., Nayak, A.K., Dhara, A.K., Eds.; Academic Press: Cambridge, MA, USA, 2022; pp. 215–237. [CrossRef]
- 168. Nugraha, A.S.; Dayli, I.R.; Sukrisno Putri, C.P.Z.; Firli, L.N.; Widhi Pratama, A.N.; Triatmoko, B.; Untari, L.F.; Wongso, H.; Keller, P.A.; Wangchuk, P. Isolation of Antibacterial Depside Constituents from Indonesian Folious Lichen, *Candelaria fibrosa*. J. Biol. Act. Prod. Nat. 2022, 12, 24–32. [CrossRef]
- Maregesi, S.; Van Miert, S.; Pannecouque, C.; Feiz Haddad, M.H.; Hermans, N.; Wright, C.W.; Vlietinck, A.J.; Apers, S.; Pieters, L. Screening of Tanzanian Medicinal Plants against Plasmodium falciparum and Human Immunodeficiency Virus. *Planta Med.* 2010, 76, 195–201. [CrossRef]

- 170. Nethengwe, M. Antiplasmodial/Antipyretic activity of some Zulu medicinal plants. J. Med. Plants Res. 2012, 6, 1255–1262. [CrossRef]
- 171. Reddy, R.C.; Vatsala, P.G.; Keshamouni, V.G.; Padmanaban, G.; Rangarajan, P.N. Curcumin for malaria therapy. *Biochem. Biophys. Res. Commun.* **2005**, 326, 472–474. [CrossRef]
- 172. Ali, A.; Sudi, S.; Sidek, H.; Embi, N.; Basir, R. The Antimalarial Effect of Curcumin Is Mediated by the Inhibition of Glycogen Synthase Kinase-3β. J. Med. Food 2016, 20, 152–161. [CrossRef]
- 173. Kamaraj, C.; Kaushik, N.K.; Rahuman, A.A.; Mohanakrishnan, D.; Bagavan, A.; Elango, G.; Zahir, A.A.; Santhoshkumar, T.; Marimuthu, S.; Jayaseelan, C.; et al. Antimalarial activities of medicinal plants traditionally used in the villages of *Dharmapuri regions* of South India. *J. Ethnopharmacol.* **2012**, *141*, 796–802. [CrossRef]
- 174. Sathiyamoorthy, P.; Lugasi-Evgi, H.; Schlesinger, P.; Kedar, I.; Gopas, J.; Pollack, Y.; Golan-Goldhirsh, A. Screening for cytotoxic and antimalarial activities in desert plants of the negev and bedouin market plant products. *Pharm. Biol.* 1999, 37, 188–195. [CrossRef]
- 175. Herlina, T.; Supratman, U.; Soedjanaatmadja, U.; Subarnas, A.; Sutardjo, S.; Abdullah, N.; Hayashi, H. Anti-malarial compound from the stem bark of Erythrina variegata. *Indones. J. Chem.* **2010**, *9*, 308–311. [CrossRef]