



**KEANEKARAGAMAN JENIS POHON DI KAMPUS
UNIVERSITAS JEMBER**

Tree Species Diversity in Campus of Jember University

ARTIKEL

Oleh

**Nanda Septesa Mikorizawan
NIM. 041810401082**

**JURUSAN BIOLOGI
FAKULTAS MATEMATIKA DAN ILMU PENGETAHUAN ALAM
UNIVERSITAS JEMBER
2012**



**KEANEKARAGAMAN JENIS POHON DI KAMPUS
UNIVERSITAS JEMBER**

Tree Species Diversity in Campus of Jember University

ARTIKEL

diajukan guna melengkapi tugas akhir dan memenuhi salah satu syarat
untuk menyelesaikan Program Studi Biologi (S1)
dan mencapai gelar Sarjana Sains

Oleh

**Nanda Septesa Mikorizawan
NIM. 041810401082**

**JURUSAN BIOLOGI
FAKULTAS MATEMATIKA DAN ILMU PENGETAHUAN ALAM
UNIVERSITAS JEMBER
2012**

PENGESAHAN

Artikel berjudul “**Keanekaragaman Jenis Pohon di Kampus Universitas Jember**” telah diperiksa dan disetujui oleh Dosen Pembimbing Utama (DPU) dan Dosen Pembimbing Anggota (DPA) pada:

hari, tanggal :

tempat : Fakultas Matematika dan Ilmu Pengetahuan Alam
Unversitas Jember

Dosen Pembimbing

DPU,

DPA,

Prof. Sudarmadji, MA., Ph.D
NIP. 195005071982121001

Drs. Moh. Imron Rosyidi, MSc.
NIP. 196205051988021001

Tree Species Diversity in Campus of Jember University

Nanda Septesa Mikorizawan¹, Sudarmadji², Moh. Imron Rosyidi².

*Biology Department, Faculty of Mathematic and Natural Science
Jember University*

ABSTRACT

*Jember University campus with an area of 65.7 ha covered by many different types of plants that can form the flora. This study aims to determine the diversity of tree species in Jember University Campus. The study was conducted in May upto October 2011. Measurement of Jember University campus area and the extent of each Faculty and Program Studies were conducted using the Global Positioning System (GPS). Census method was used to identify and count the number of trees at the study site. Environmental parameters were measured at each study site, include: the degree of acidity (pH) of soil, soil texture, air temperature, and humidity. Species diversity index is determined using the Shannon-Weiner formula. Abundance type is determined by the criteria used in the abundance of individual species of trees. Importance value index of type is used to observe the dominance of plant species at each location. The results showed that in Jember University Campus, there were forty-three species of trees with a total of as many as 789 tree species and the species density was achieved by *Swietenia macrophylla* King. Jacq. (6.41), followed by *Samanea saman* Jacq. Merr. (3.28) and *Spathodea campanulata* Beauv. (2.41), respectively. Of all types of existing trees, three species dominates, namely *Swietenia macrophylla* King. Jacq., *Samanea saman* Jacq. Merr., And *Spathodea campanulata* Beauv. .. The index value the importance of trees in the University of Jember ranged from 0.85 at the tree *Tectona grandis* L. f. and *Bombax ceiba* L. until 46.39 on the tree *Swietenia macrophylla* King. Jacq. .. The most dominant tree species on the University Campus Jember is *Swietenia macrophylla* King. Jacq. followed by *Samanea saman* Jacq. Merr. and *Spathodea campanulata* Beauv., respectively. Tree species diversity index values in the range of 0.20 at the Nursing Science Program (PSIK) up to 2.66 at the Faculty of Agriculture, which means it has relatively low species diversity through the medium. The total area of study site is 27.75 ha with a mean value of 1.98 ha each faculty.*

Key word: Global Positioning System (GPS), tree species diversity, species diversity index, importance value index of type.

¹*Student of Biology Department, Faculty of Mathematic and Natural Science.*

²*Lecturer of Biology Department, Faculty of Mathematic and Natural Science.*