Digital Repository Universitas Jember

ISSN 2086-0382 E-ISSN 2477-3344

CAUCH

Jurnal Matematika Murni dan Aplikasi

Volume 4, Issue 3, November 2016













Volume 4, Issue 3, November 2016

PUBLICATION ETHICS

Journal Cauchy is a peer-reviewed electronic national journal. This statement clarifies ethical behaviour of all parties involved in the act of publishing an article in this journal, including the author, the chief editor, the Editorial Board, the peer-reviewer and the publisher (Mathematics Department of Maulana Malik Ibrahim State Islamic University of Malang). This statement is based on COPE's Best Practice Guidelines for Journal Editors.

Ethical Guideline for Journal Publication

The publication of an article in a peer-reviewed Cauchy is an essential building block in the development of a coherent and respected network of knowledge. It is a direct reflection of the quality of the work of the authors and the institutions that support them. Peer-reviewed articles support and embody the scientific method. It is therefore important to agree upon standards of expected ethical behavior for all parties involved in the act of publishing: the author, the journal editor, the peer reviewer, the publisher and the society.

As publisher of Pure and applied mathematics journal, we take our duties to back up over all stages of publishing seriously and we recognize our ethical and other responsibilities. We are committed to ensuring that advertising, reprint or other commercial revenue has no impact or influence on editorial decisions.

Publication decisions

The editor of Cauchy is responsible for deciding which of the articles submitted to the journal should be published. The validation of the work in question and its importance to researchers and readers must always drive such decisions. The editors may be guided by the policies of the journal's editorial board and constrained by such legal requirements as shall then be in force regarding libel, copyright infringement and plagiarism. The editors may confer with other editors or reviewers in making this decision.

Fair Play

An editor at any time evaluates manuscripts for their intellectual content without regard to race, gender, sexual orientation, religious belief, ethnic origin, citizenship, or political philosophy of the authors.

Confidentiality

The editor and any editorial staff must not disclose any information about a submitted manuscript to anyone other than the corresponding author, reviewers, potential reviewers, other editorial advisers, and the publisher, as appropriate.

Any manuscripts received for review must be treated as confidential documents. They must not be shown to or discussed with others except as authorized by the editor.

Disclosure and conflicts of interest

Unpublished materials disclosed in a submitted manuscript must not be used in an editor's own research without the express written consent of the author.



Volume 4, Issue 3, November 2016

PREFACE

Cauchy is a national journal published by Mathematics Department, Science and Technology Faculty, Maulana Malik Ibrahim State Islamic University of Malang. This is the second issue of this year. It contains 6 (six) article from all over the country, not only from local area of East Java. Those articles covered graph theory, numerical analysis, applied mathematics, statistics, economic mathematics, and algebra. This issue was authored by 13 authors and co-authors.

In the first article, the author discussed the application of pontryagin's minimum principle in optimum time of missile manoeuvring. Analysis of surface-to-surface missile which considered objective and dynamical system showed that trajectory tracking of this missile was split up in three sub-intervals, namely flying, climbing, and diving in which had different performances of speed.

The second article discussed about power of test path analysis and partial least square analysis. The result showed that the usage of path analysis provides better value than *PLS* at R^2 value in terms of the power of the test. Behavioral conditions of work of civil servants in Kediri City Government did not follow the rules of the science of human resources in terms of election officials echelon structural or civil servants.

The third article, entitled "Estimation Parameters and Modelling Zero Inflated Negative Binomial" elaborated that the estimation parameter of Zero inflated Negative Binomial (ZINB) model was conducted using Maximum Likelihood Estimation (MLE) and to maximize the likelihood function used the EM (Expectation Maximization) algorithm. For parameter test predictor variable that has significant effect on the number of cases of Tetanus Neonatorum are pregnant mothers visit K4 (X₁) and maternal mothers assisted by health workers (X₃) for the negative binomial state models, while zero inflation state model predictor variable that has significant effect on the number of cases of Tetanus Neonatorum include the percentage of neonates visit (X₄).

The fourth article emphasized the discussion to Leontief input-output method for the fresh milk distribution linkage analysis. Leontief Input-Output Method can illustrate the intersectoral linkages and identify the key sector in the fresh milk distribution. It is clearly shown that the merging of the collectors' and the fresh milk processing industries' activities offers high indices of both backward linkages and forward linkages so that it can be the key sector in the fresh milk distribution.



Volume 4, Issue 3, November 2016

PREFACE

The last article entitled "Some Properties from Construction of Finite Projective Planes of Order 2 and 3". It showed that a finite projective plane is a geometric construction that extends the concept of a plane. It consists of a set of points and lines together with an incidence relation between them and order 2 is the smallest order. In this paper some properties from construction of finite projective planes of order 2 and order 3 which is as a comparison are observed.





Volume 4, Issue 3, November 2016

PUBLICATION ETHICS

Journal Cauchy is a peer-reviewed electronic national journal. This statement clarifies ethical behaviour of all parties involved in the act of publishing an article in this journal, including the author, the chief editor, the Editorial Board, the peer-reviewer and the publisher (Mathematics Department of Maulana Malik Ibrahim State Islamic University of Malang). This statement is based on COPE's Best Practice Guidelines for Journal Editors.

Ethical Guideline for Journal Publication

The publication of an article in a peer-reviewed Cauchy is an essential building block in the development of a coherent and respected network of knowledge. It is a direct reflection of the quality of the work of the authors and the institutions that support them. Peer-reviewed articles support and embody the scientific method. It is therefore important to agree upon standards of expected ethical behavior for all parties involved in the act of publishing: the author, the journal editor, the peer reviewer, the publisher and the society.

As publisher of Pure and applied mathematics journal, we take our duties to back up over all stages of publishing seriously and we recognize our ethical and other responsibilities. We are committed to ensuring that advertising, reprint or other commercial revenue has no impact or influence on editorial decisions.

Publication decisions

The editor of Cauchy is responsible for deciding which of the articles submitted to the journal should be published. The validation of the work in question and its importance to researchers and readers must always drive such decisions. The editors may be guided by the policies of the journal's editorial board and constrained by such legal requirements as shall then be in force regarding libel, copyright infringement and plagiarism. The editors may confer with other editors or reviewers in making this decision.

Fair Play

An editor at any time evaluates manuscripts for their intellectual content without regard to race, gender, sexual orientation, religious belief, ethnic origin, citizenship, or political philosophy of the authors.

Confidentiality

The editor and any editorial staff must not disclose any information about a submitted manuscript to anyone other than the corresponding author, reviewers, potential reviewers, other editorial advisers, and the publisher, as appropriate.

Any manuscripts received for review must be treated as confidential documents. They must not be shown to or discussed with others except as authorized by the editor.

Disclosure and conflicts of interest

Unpublished materials disclosed in a submitted manuscript must not be used in an editor's own research without the express written consent of the author.



Volume 4, Issue 3, November 2016

PUBLICATION ETHICS

Contribution to Editorial Decisions

Peer review assists the editor in making editorial decisions and through the editorial communications with the author may also assist the author in improving the paper.

Promptness

Any selected referee who feels unqualified to review the research reported in a manuscript or knows that its prompt review will be impossible should notify the editor and excuse himself from the review process.

Standards of Objectivity

Reviews should be conducted objectively. Personal criticism of the author is inappropriate. Referees should express their views clearly with supporting arguments.

Acknowledgement of Sources

Reviewers should identify relevant published work that has not been cited by the authors. Any statement that an observation, derivation, or argument had been previously reported should be accompanied by the relevant citation. A reviewer should also call to the editor's attention any substantial similarity or overlap between the manuscript under consideration and any other published paper of which they have personal knowledge.

Disclosure and Conflict of Interest

Privileged information or ideas obtained through peer review must be kept confidential and not used for personal advantage. Reviewers should not consider manuscripts in which they have conflicts of interest resulting from competitive, collaborative, or other relationships or connections with any of the authors, companies, or institutions connected to the papers.

Reporting standards

Authors of reports of original research should present an accurate account of the work performed as well as an objective discussion of its significance. Underlying data should be represented accurately in the paper. A paper should contain sufficient detail and references to permit others to replicate the work. Fraudulent or knowingly inaccurate statements constitute unethical behaviour and are unacceptable.

Data Access and Retention

Authors are asked to provide the raw data in connection with a paper for editorial review, and should be prepared to provide public access to such data (consistent with the ALPSP-STM Statement on Data and Databases), if practicable, and should in any event be prepared to retain such data for a reasonable time after publication.



Volume 4, Issue 3, November 2016

PUBLICATION ETHICS

Originality and Plagiarism

The authors should ensure that they have written entirely original works, and if the authors have used the work and/or words of others that this has been appropriately cited or quoted.

Multiple, Redundant or Concurrent Publication

An author should not in general publish manuscripts describing essentially the same research in more than one journal or primary publication. Submitting the same manuscript to more than one journal concurrently constitutes unethical publishing behaviour and is unacceptable.

Acknowledgement of Sources

Proper acknowledgment of the work of others must always be given. Authors should cite publications that have been influential in determining the nature of the reported work.

Authorship of the Paper

Authorship should be limited to those who have made a significant contribution to the conception, design, execution, or interpretation of the reported study. All those who have made significant contributions should be listed as co-authors. Where there are others who have participated in certain substantive aspects of the research project, they should be acknowledged or listed as contributors. The corresponding author should ensure that all appropriate co-authors and no inappropriate co-authors are included on the paper, and that all co-authors have seen and approved the final version of the paper and have agreed to its submission for publication.

Hazards and Human or Animal Subjects

If the work involves chemicals, procedures or equipment that have any unusual hazards inherent in their use, the author must clearly identify these in the manuscript.

Disclosure and Conflicts of Interest

All authors should disclose in their manuscript any financial or other substantive conflict of interest that might be construed to influence the results or interpretation of their manuscript. All sources of financial support for the project should be disclosed.

Fundamental errors in published works

When an author discovers a significant error or inaccuracy in his/her own published work, it is the author's obligation to promptly notify the journal editor or publisher and cooperate with the editor to retract or correct the paper.



On The Local Metric Dimension of Line Graph of Special Graph

Marsidi¹, Dafik², Ika Hesti Agustin³, Ridho Alfarisi⁴

¹Mathematics Edu. Depart. IKIP Jember Indonesia ²Mathematics Edu. Depart. University of Jember Indonesia ³Mathematics Depart. University of Jember Indonesia ⁴Mathematics Depart. ITS Surabaya Indonesia

Email: marsidiarin@gmail.com, d.dafik@gmail.com, ikahestiagustin@gmail.com, alfarisi38@gmail.com.

ABSTRACT

Let G be a simple, nontrivial, and connected graph. $W = \{w_1, w_2, w_3, ..., w_k\}$ is a representation of an ordered set of k distinct vertices in a nontrivial connected graph G. The metric code of a vertex v, where $v \in G$, the ordered $r(v|W) = (d(v, w_1), d(v, w_2), ..., d(v, w_k))$ of k-vector is representations of v with respect to W, where $d(v, w_i)$ is the distance between the vertices v and w_i for $1 \le i \le k$. Furthermore, the set W is called a local resolving set of G if $r(u|W) \ne r(v|W)$ for every pair u, v of adjacent vertices of G. The local metric dimension ldim(G) is minimum cardinality of W. The local metric dimension exists for every nontrivial connected graph G. In this paper, we study the local metric dimension of line graph of special graphs, namely path, cycle, generalized star, and wheel. The line graph L(G) of a graph G has a vertex for each edge of G, and two vertices in L(G) are adjacent if and only if the corresponding edges in G have a vertex in common.

Keywords: metric dimension, local metric dimension number, line graph, resolving set.

INTRODUCTION

All graph in this paper are simple, nontrivial, and undirected, for more detail basic definition of graph, see [1]. In [2] define the distance d(u, v) between two vertices u and v in a connected graph G is the length of a shortest path between these two vertices. Suppose that $W = \{w_1, w_2, w_3, ..., w_k\}$ is an ordered set of vertices of a nontrivial connected graph G. The metric representation of v with respect to W is the k-vector $r(v|W) = (d(v, w_1), d(v, w_2), ..., d(v, w_k))$. Distance in graphs has also been used to distinguish all of the vertices of a graph. The set W is called a resolving set for G if distinct vertices of G have distinct representations with respect to W. The metric dimension dim(G) of G is the minimum cardinality of resolving set for G [3]. Furthermore, we consider those ordered sets W of vertices of G for which any two vertices of G having the same code with respect to W are not adjacent in G. If $r(u|W) \neq r(v|W)$ for every pair u, v of adjacent vertices of G, then W is called a local metric set of G. The minimum k for which G has a local metric k-set is the local metric dimension of G, which is denoted by ldim(G) [4].

In this paper, we study the local metric dimension number of line graph of special graphs, namely path, cycle, generalized star, and wheel graph. Line graphs are a special case of intersection graphs. The line graph L(G) of a graph G has a vertex for each edge of G, and two vertices in L(G) are adjacent if and only if the corresponding edges in G have a vertex in common. Thus, the line graph L(G) is the intersection graph corresponding to the endpoint sets of the edges

of G. As for an example, Figure 1 shows a graph G and its line graph L(G) [5]. The results show that distinct vertices of each L(G), with G is a special graph, namely path, cycle, generalized star, and wheel graph have distinct representations with respect to *W*, and their local metric dimension attain a minimum number. Furthermore, [6] showed that on the metric dimension, the upper dimension and the resolving number of graphs, [7] showed the metric dimension of amalgamation of cycles, [8] studied the constant metric dimension of regular graphs, [2] obtained resolvability in graphs and the metric dimension of a graph. The last, [9] showed the Fault-tolerant metric and partition dimension of graph.



Figure 1. A graph and its line graph

RESULTS AND DISCUSSIONS

We have some observation to show the result of line graph of path, cycle, generalized star, and wheel graph. Thus, we have four main theorem about local metric dimension to discuss as follows.

Observation 1. The order and the size of $L(P_n)$ are $|V(L(P_n))| = n - 1$ and $|E(L(P_n))| = n - 2$, respectively.

Proof. Line graph of Path $L(P_n)$ is connected, simple, and undirected graph with vertex set $V(L(P_n)) = \{v_i; 1 \le i \le n-1\}$ and edge set $E(L(P_n)) = \{v_iv_{i+1}; 1 \le i \le n-2\}$. Thus, $|V(L(P_n))| = n - 1$ and $|E(L(P_n))| = n - 2$. See Figure 2 (a) and 2 (b) for illustration.

Theorem 1. For $n \ge 2$, the local metric dimension of line graph of path is $ldim(L(P_n)) = 1$.

Proof. By observation 1, line graph of path $L(P_n)$ is isomorphic to P_{n-1} . Thus, the vertex set and the edge set of $L(P_n)$ are $V(L(P_n)) = \{v_i; 1 \le i \le n-1\}$ and $E(L(P_n)) = \{v_iv_{i+1}; 1 \le j \le n-2\}$, respectively. The number of vertices $|V(L(P_n))| = n-1$ and the size $|E(L(P_n))| = n-2$. By Theorem 3, $ldim(P_n) = 1$, thus $ldim(L(P_n)) = 1$. It concludes the proof. See Figure 2 (c) for illustration.



Figure 2. (a) P_8 , (b) $L\{P_8\}$, (c) Construction of local resolving set $W = \{v_1\}$

Observation 2. The order and the size of $L(C_n)$ are $|V(L(C_n))| = n$ and $|E(L(P_n))| = n$, respectively.

Proof. Line graph of cycle $L(C_n)$ is connected, simple, and undirected graph with vertex set $V(L(C_n)) = \{v_i; 1 \le i \le n\}$ and edge set $E(L(C_n)) = \{v_i v_{i+1}; 1 \le j \le n-1\} \cup \{v_n v_1; \}$. Thus, $|V(L(C_n))| = n$ and $|E(L(C_n))| = n$. See Figure 3 (a) and 3 (b) for illustration.

Theorem 2. For $n \ge 3$, the local metric dimension of line graph of cycle is

 $ldim(L(C_n)) = \begin{cases} 1, & for \ n \ even \\ 2, & for \ n \ odd \end{cases}$ **Proof**. By observation 2, line graph of cycle $L(C_n)$ is isomorphic to C_n . Thus, the vertex set and the edge set of $L(C_n)$ are $V(L(C_n)) = \{v_i; 1 \le i \le n\}$ and $E(L(C_n)) = \{e_j; 1 \le j \le n\}$, respectively. The order of vertices $|V(L(C_n))| = n$ and the size $|E(L(C_n))| = n$. By Theorem 3,

$$ldim(C_n) = \begin{cases} 1, & \text{for } n \text{ even} \\ 2, & \text{for } n \text{ odd} \end{cases}$$

thus

$$dim(L(C_n)) = \begin{cases} 1, & \text{for } n \text{ even} \\ 2, & \text{for } n \text{ odd} \end{cases}$$

It concludes the proof. See Figure 3 (c) for illustration



Figure 3. (a) C_8 , (b) $L\{C_8\}$, (c) Construction of local resolving set $W = \{v_1\}$

Observation 3. The order and the size of $L(S_{n,m})$ are $|V(L(S_{n,m}))| = mn$ and $|E(L(S_{n,m}))| = mn + mn$ $\frac{n(n-3)}{2}$, respectively.

Proof. Line graph of generalized star $L(S_{n,m})$ is connected, simple, and undirected graph with vertex set $V(L(S_{n,m})) = \{x_{i,j}; 1 \le i \le n, 1 \le j \le m\}$ and edge set $E(L(S_{n,m})) = \{x_{i,j}x_{i,j+1}; 1 \le i \le m\}$ $n, 1 \le j \le m - 1$ $\bigcup \{x_{i,1}x_{t,1}; i \ne t, 1 \le i, t \le n\}$. Thus, $|V(L(S_{n,m}))| = mn$ and $|E(L(S_{n,m}))| = mn$ $mn + \frac{n(n-3)}{2}$. See Figure 4 (a) and 4 (b) for illustration.

Theorem 3. For $n \ge 3$, the local metric dimension of line graph of generalized star is $dim(L(S_{n,m})) = n - 1.$

Proof. By observation 3, the order and the size of $L(S_{n,m})$ are $|V(L(S_{n,m}))| = mn$ and $|E(L(S_{n,m}))| = mn + \frac{n(n-3)}{2}$, respectively. Suppose the lower bound is $ldim(L(S_{n,m})) \ge n-2$ with the resolving set $W' = \{x_{i,1}; 1 \le i \le n-2\}$. Thus two vertices of $V(L(S_{n,m})) - W'$ with respect to W' are

$$r(x_{n-1,1}|W') = r(x_{n,1}|W') = \left(\underbrace{1,\dots,1}_{n-2 \text{ times}}\right)$$

It is easy to see that the line graph of generalized star has the same vertex representation respecting to W'. Thus, the lower bound is $ldim(L(S_{n,m})) \ge n-1$. Now, we will show that $ldim(L(S_{n,m})) \le n-1$ by determining the resolving set $W = \{x_{i,1}; 1 \le i \le n-1\}$ and the vertex representation of $V(L(S_{n,m})) - W$ respect to W, as follows

$$r(x_{n,j}|W) = \left(\underbrace{j, \dots, j}_{n-1 \text{ times}}\right); 1 \le j \le m$$
$$r(x_{i,j}|W) = \left(\underbrace{j, \dots, j}_{i-1 \text{ times}}, j-1, \underbrace{j, \dots, j}_{n-i-1 \text{ times}}\right); 1 \le i \le n-1, 2 \le j \le m$$

It easy to see that $\forall u, v \in V(L(S_{n,m})) - W$ have a different representation respect to W for every pair u, v of adjacent vertices in $L(S_{n,m})$. The cardinality of resolving set $L(S_{n,m})$ is n - 1, thus $ldim(L(S_n)) \leq n - 1$. It concludes the proof. See Figure 4 (c).



Figure 4. (a) $S_{6,3}$, (b) $L(S_{6,3})$, (c) Construction of local resolving set $W = \{x_{1,1}, x_{2,1}, x_{3,1}, x_{4,1}, x_{5,1}\}$

Observation 4. The order and the size of $L(W_n)$ are $|V(L(W_n))| = 2n$ and $|E(L(W_n))| = \frac{n(n+5)}{2}$, respectively.

Proof. Line graph of wheel $L\{W_n\}$ is connected, simple, and undirected graph with vertex set $V(L(W_n)) = \{x_{i,j}; 1 \le i \le n, 1 \le j \le 2\}$ and edge set $E(L(S_{n,m})) = \{x_{i,1}x_{i,2}; 1 \le i \le n\} \cup \{x_{i,1}x_{t,1}; i \ne t, 1 \le i, t \le n\} \cup \{x_{i,2}x_{i+1,2}; 1 \le i \le n-1\} \cup \{x_{i,2}x_{i+1,1}; 1 \le i \le n-1\} \cup \{x_{1,2}x_{n,2}\} \cup \{x_{1,1}x_{n,2}\}$. Thus, $|V(L(W_n))| = 2n$ and $|E(L(W_n))| = \frac{n(n+5)}{2}$. See Figure 5 (a) and 5 (b) for illustration.

Theorem 4. For $n \ge 3$, the local metric dimension of line graph of wheel is $LDim(L(W_n)) = n - 1$. **Proof.** By observation 4, the order and the size of $L(W_n)$ are $|V(L(W_n))| = 2n$ and $|E(L(W_n))| = \frac{n(n+5)}{2}$, respectively. Suppose the lower bound of $L(W_n)$ is $ldim(L(W_n)) \ge n - 2$ with the resolving set $W' = \{x_{i,1}; 1 \le i \le n - 2\}$. Thus two vertices of $V(L(W_n)) - W'$ with respect to W' are

$$r(x_{n-1,1}|W') = r(x_{n,1}|W') = \left(\underbrace{1, ..., 1}_{n-2 \text{ times}}\right)$$

It is easy to see that the line graph of wheel possess the same vertex representation respecting to W'. Thus the lower bound is $ldim(L\{W_n\}) \ge n - 1$. Now, we will show that $ldim(L\{W_n\}) \le n - 1$

by determining the resolving set $W = \{x_{i,1}; 1 \le i \le n-1\}$ and the vertex representation of $V(L\{W_n\}) - W$ respect to W, as follows.

$$W = \{x_{1,1}, x_{2,1}, x_{3,1}, x_{4,1}, x_{5,1}, x_{6,1}, x_{7,1}\}$$
$$r(x_{n,1}|W) = \begin{pmatrix} 1, \dots, 1\\ n-1 \text{ times} \end{pmatrix}$$
$$(x_{i,2}|W) = \begin{pmatrix} 2, \dots, 2\\ i-1 \text{ times} \end{pmatrix}, 1, \underbrace{2, \dots, 2}_{n-i-1 \text{ times}} \end{pmatrix}; 2 \le i \le n-1$$
$$r(x_{1,2}|W) = \begin{pmatrix} 1, 1, \underbrace{2, \dots, 2}_{n-3 \text{ times}} \end{pmatrix}$$

r

 $r(x_{n,2}|W) = \begin{pmatrix} 1, 2, ..., 2 \\ n-2 \text{ times} \end{pmatrix}$ It easy to see that $\forall u, v \in V(L(W_n)) - W$ have a different representation respect to W for every pair u, v of adjacent vertices in $L(W_n)$. The cardinality of resolving set $L(W_n)$ is n - 1, thus $ldim(L(W_n)) \leq n - 1$. It concludes the proof. See Figure 5 (c) for illustration.



Figure 5. (a) W_8 , (b) $L\{W_8\}$, (c) Construction of local resolving set

CONCLUSION

We have shown the local metric dimension number of line graph of special graphs, namely line graph of path, cycle, star, and wheel. The results show that the local metric dimension numbers attain the best lower bound. However we have not found the sharpest lower bound for any connected graph, therefore we proposed the following open problem.

Open Problem 1. *Let G be a connected graph, obtain the best lower bound of the local metric dimension of any graph G.*

REFERENCES

- [1] G. Chartrand, E. Salehi, and P. Zhang, "The partition dimension of a graph," *Aequationes Math.*, vol. 59, pp. 45–54, 2000.
- [2] G. Chartrand, L. Eroh, and M. A. Johnson, "Resolvability in graphs and the metric dimension of a graph," *Discrate Appl. Math.*, vol. 105, pp. 99–113, 2000.
- [3] M. Imran, A. Q. Baig, S. Ahtsham, U. Haq, and I. Javaid, "On the metric dimension of circulant graphs," *Appl. Math. Lett.*, vol. 25, no. 3, pp. 320–325, 2012.
- [4] G. A. B. Ramirez, C. G. Gomez, and J. A. R. Valazquez, "Closed formulae for the local metric dimension of corona product graphs," *Electron. notes Discret. Math.*, vol. 46, pp. 27–34, 2014.
- [5] LECTURE 1, "INTRODUCTION TO GRAPH MODELS," pp. 1–25.
- [6] D. Garijo, A. González, and A. Márquez, "On the metric dimension , the upper dimension and the resolving number of graphs," *Discret. Appl. Math.*, vol. 161, no. 10–11, pp. 1440–1447, 2013.
- [7] H. Iswadi, E. T. Baskoro, A. N. M. Salman, R. Simanjuntak, N. Science, and U. Surabaya, "THE METRIC DIMENSION OF AMALGAMATION," *Far East J. Math. Sci.*, vol. 41, no. 1, pp. 19–31, 2010.
- [8] I. Javaid, M. T. Rahim, and K. Ali, "Families of regular graphs with constant metric dimension," *Util. Math.*, no. October 2016, 2008.
- [9] M. A. Chaudhry, I. Javaid, and M. Salman, "Fault-Tolerant Metric and Partition Dimension of Graphs," *Util. Math.*, 2010.

