

21/1
Vol. 21 No. 1
JANUARY 2021

P-ISSN: 1411-8033
E-ISSN: 2614-0101



mutiara medika

Jurnal Kedokteran dan Kesehatan

RESEARCH ARTICLE

- 01 The Influence of Chewing Habits on the Degree of Impacted Cerumen
Asti Widuri
- 07 Mother's Behavior in Using Smartphone to Find Child's Health Information in Pontianak City
Fitriyah, Rini Andriani, Mitra Handini
- 15 The Duration of Reading the Qur'an with the Stress Levels in Medical Faculty Students in Palembang
Erina Ahmad Ghifari, Syahrul Muhammad, M. Avif Ababil
- 20 Relationship between Knowledge and Using Personal Protective Equipment in Tuberculosis Ward and Polyclinic of Hospital X
Ekorini Listiowati, Fitria Eka Rianti
- 26 The Correlation between Body Mass Index to Hypertension in a Rural Area in East Java
Alvin Hartanto Kurniawan, Beneditus Hanjaya Suwandi, Dicky Teguh Prakoso, Reynaldy Cliftianto Moksidy, Irma Talitha, Maharani Kartika Anggraeni, Danang Zulkifli
- 32 Correlation the Destiny of *Dermatophagoides* sp. in Households with the Score for Allergic Rhinitis in Jember, East Java
Erma Sulistyanyingsih, Elisa Fadia Lalli, Cholis Abrori
- 39 Comparison of Serum Interleukin 10 Levels between Leprosy and Non-Leprosy Population
Enda Silvia Putri, Marniati Marniati, Arafah Husna, Afriani Maifizar
- 45 Effect of Anti Mycobacterium Tuberculosis Nanoherbal Legundi Leaf Extract (*Vitex trifolia*)
Bella Atika Dwiyantri, Siti Mulyani, Oche Axena Yulhan, Humairah Medina Liza Lubiz, Ance Roslina
- 51 The Correlation between Osteoarthritis Knee Damage Based on Ultrasound with Kellgren-Lawrence
Kanti Wensri Sevni Kurniawati, Muhammad Ilyas, Mirna Muis, Andi Alfian Z, Faridin, Sri Asriyani
- 60 Corticosteroid Expression Level of Cyclin-D1 between Endometriomas and Ovarian Carcinomas
Alfun Dhiya An, Supriyatningsih
- 67 Correlation between Cup Ratio and Optic Nerve Disc with High Myopia
Nur Shani Meida, Akhmad Ikluddin, Yunani Setyandriani, Nadia Dina Ayu

Accredited SK No: 10/E/KPT/2019, April 4th, 2019

Published by Universitas Muhammadiyah Yogyakarta
in collaboration with Asosiasi Pendidikan Kedokteran dan Kesehatan Muhammadiyah

Mutiara Medika: Jurnal Kedokteran dan Kesehatan

Journal Title	Mutiara Medika: Jurnal Kedokteran dan Kesehatan
ISSN	2614-0101 (online) 1411-8033 (print)
DOI Prefix	Prefix 10.18196 by  Crossref
Editor in Chief	Tri Wulandari Kesetyaningsih
Publisher	Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta in collaboration with the Association of Medicine and Health Education of Muhammadiyah (Asosiasi Pendidikan Kedokteran dan Kesehatan Muhammadiyah-APKKM)
Frequency	2 issues per year
Citation Analysis	Google Scholar

Mutiara Medika: Jurnal Kedokteran dan Kesehatan or also known as **MMJJK** is peer reviewed journal published by Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta in collaboration with the Association of Medicine and Health Education of Muhammadiyah (Asosiasi Pendidikan Kedokteran dan Kesehatan Muhammadiyah-APKKM). MMJJK is accredited grade 3 by Ministry of Research, Technology and Higher Education of Republic of Indonesia Decree (SK) [No. 200/M/KPT/2020](#).

Since the first issued in January 2001, MMJJK is publishing scientific articles consistently, both research and case report. MMJJK publishes the new editions every January and July. By January 2016, MMJJK has published both printed (book) and electronic (PDF) versions. Electronic articles are accessible openly on the [web page](#). MMJJK is indexed by [SINTA \(Science and Technology Index\) Grade 3](#), [Google Scholar](#), [Base](#), [Garuda \(Garba Rujukan Digital\)](#), and [Crossref](#). MMJJK focuses on all aspects of human health (but not limited):

1. Biomedical Sciences
2. Clinical Medicine
3. Public Health Sciences
4. Islamic Medicine

IMPORTANT: Before you submit a manuscript, make sure that your paper is prepared using MMJJK template ([research article](#) and [case report](#)), has been proofread and polished carefully, and conformed to the [MMJJK AUTHOR GUIDELINES](#).

Online Submissions

- If you already have a Username/Password? **GO TO LOGIN**
- If you are new comers and need a Username/Password? **GO TO REGISTRATION**

Registration and login are required to submit items online and to check the status of current submissions.

PURCHASE ORDER

Subscription or order hardcopy Mutiara Medika: Jurnal Kedokteran dan Kesehatan [click here](#)



Editorial Office:
Journal Room, G1 (Biomedic) Building, Ground Floor, Faculty of Medicine and Health Science
Universitas Muhammadiyah Yogyakarta,
Jalan Lingkar Selatan (Brawijaya), Tamantirto, Kasihan, Bantul, Daerah Istimewa Yogyakarta,
Indonesia
Phone: +62 274 387 656 (ext: 231)
WA : +62 882 3323 4696
Website: <http://journal.umy.ac.id/index.php/mm>
E-mail: mmjkk@umy.university



Mutiara Medika: Jurnal Kedokteran dan Kesehatan is licensed under a [Creative Commons Attribution 4.0 International License](#). [View My Stats](#)

Editorial Team

Editor in Chief

1. [Tri Wulandari Kesetyaningsih](#), Scopus ID: 56607303300, Department of Parasitology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

Managing Editor

1. [Sri Nabawiyati Nurul Makiyah](#), Department of Histology and Immunology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

Editors

1. [Sri Sundari](#), Universitas Muhammadiyah Yogyakarta, Indonesia
2. [Ana Majdawati](#), Scopus ID: 57209331456, Department of Radiology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia
3. [EM Sutrisna](#), Scopus ID: 56725762600, Farmakologi dan Farmakogenomik, Universitas Muhammadiyah Surakarta, Indonesia
4. [Muhammad Sasmito Djati](#), Scopus ID: 56403414400, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Indonesia
5. [Sri Andarini](#), Scopus ID: 57188996572, Department of Public Health, Faculty of Medicine, Universitas Brawijaya, Indonesia
6. [Titiek Hidayati](#), Universitas Muhammadiyah Yogyakarta, Indonesia
7. [Eko Suhartono](#), Universitas Lambung Mangkurat, Indonesia
8. [Lilis Suryani](#), Scopus ID: 57188995983, Department of Microbiology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia
9. [Erna Rochmawati](#), Scopus ID: 36150260600, Department of Medical Surgical Nursing, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia
10. [Iman Permana](#), Scopus ID: 57201316848, Department of Public Health, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta

Editorial Board Member

1. [Anya Ahmed](#), Scopus ID: 55587710100, School of Health and Society, University of Salford, United Kingdom
2. [Muhammad Taher](#), Scopus ID: 57201790510, Department of Pharmacy, Faculty of Pharmacy, International Islamic University, Malaysia
3. [Paula Ormandy](#), Scopus ID: 13007313400, School of Nursing, University of Salford, United Kingdom

Peer Reviewers

1. [Yoshizo Matsuka](#), Scopus ID: 7003862097, Department of Stomatognathic Function and Occlusal Reconstruction Faculty of Dentistry, Tokushima University, Japan

[Sri Andarini](#), Scopus ID: 57188996572, Department of Public Health, Faculty of Medicine, Universitas Brawijaya, Indonesia

[Muhammad Sasmito Diati](#), Scopus ID: 56403414400, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Indonesia

[Muhaimin Rifa'i](#), Scopus ID: 8269391600, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Indonesia

[Ratna Indriawati](#), Scopus ID: 57204094606, Department of Physiology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Supriyatiningasih Supriyatiningasih](#), Scopus ID: 57194828856, Department of Obstetrics & Gynecology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Marsetyawan Heparis Nur Ekandaru Soesatyo](#), Department of Histology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada

[Sagiran Sagiran](#), Scopus ID: 56922140700, Department of Surgery, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Ahmad Hamim Sadewa](#), Scopus ID: 6506553494, Departement of Biochemistry, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada

[Lantip Ruijito](#), Scopus ID: 56656540400, Department of Anatomy and Clinical Genetics, Faculty of Medicine, Universitas Jenderal Soedirman

[Ikhlas Muhammad Jenie](#), Scopus ID: 57192904333, Department of Physiology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Farida Juliantina Rachmawaty](#), Scopus ID: 56505407000, Department of Microbiology, Faculty of Medicine, Universitas Islam Indonesia, Indonesia

[Nur Shani Meida](#), Universitas Muhammadiyah Yogyakarta, Indonesia

[Warih Andan Puspitosari](#), Scopus ID: 57205700481, Department of Pshychiatry, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Juliani Ibrahim](#), Scopus ID: 36164160500, Department of Public Health, Faculty of Medicine, Universitas Muhammadiyah Makassar, Indonesia

[Mustofa Mustofa](#), Scopus ID: 57203104599, Department of Pharmacology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Indonesia

[Joerg Haier](#), Scopus ID: 7004044592, Department of Geburtshilfe und Frauenheilkunde, Universitatklinikum Muenster/Westfaliche Wilhelms, University of Muenster, Germany

[Sabtanti Harimurti](#), Scopus ID: 36448103200, Department of Pharmaceutical Chemistry, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Yoni Astuti](#), Scopus ID: 56575015700, Department of Biochemistry, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

Digital Repository Universitas Jember

[Ardi Pramono](#), Scopus ID: 56515234600, Department of Anesthesiology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Indonesia

[Wiwik Kusumawati](#), Scopus ID: 57160948400, Department of Medical Education, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta

[Supanji Supanji](#), Scopus ID: 55710928300, Department of Ophthalmology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Indonesia

[Sri Rahayu Lestari](#), Scopus ID: 56403367700, Animal Physiology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Malang, Indonesia

[Restu Syamsul Hadi](#), Scopus ID: 57192266794, Biomedical Sciences, Faculty of Medicine, Universitas Yarsi, Indonesia

[Firdaus Hamid](#), Scopus ID: 35933184500, Department of Microbiology, Faculty of Medicine, Universitas Hasanuddin, Indonesia

[EM Sutrisna](#), Scopus ID: 56725762600, Farmakologi dan Farmakogenomik, Universitas Muhammadiyah Surakarta, Indonesia

[Neni Trilusiana Rahmawati](#), Scopus ID: 6504816137, Department of Anatomy, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Indonesia

[Irwan Supriyanto](#), Department of Psychiatry, Faculty of Medicine, Universitas Gadjah Mada, Indonesia

[Denny Agustiningasih](#), Scopus ID: 56946943700, Department of Physiology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Indonesia

[Lilik Zuhriyah](#), Scopus ID: 55753460700, Department of Public Health, Faculty of Medicine, Universitas Brawijaya

Vol 21, No 1: January 2021

Table of Contents

Research

The Influence of Chewing Habits on the Degree of Impacted Cerumen

10.18196/mmjkk.v21i1.9576

Abstract view : 121 times

Asti Widuri

Mother's Behaviour in Using Smartphone to Find Child's Health Information in Pontianak City

10.18196/mmjkk.v21i1.8624

Abstract view : 87 times

Fithriyyah Fithriyyah, Rini Andriani, Mitra Handini

The Duration of Reading the Qur'an with the Stress Levels in Medical Faculty Students in Palembang

10.18196/mmjkk.v21i1.8488

Abstract view : 113 times

Ahmad Ghiffari, Syahrul Muhammad, M. Avif Ababil

Relationship between Knowledge and Using Personal Protective Equipment in Tuberculosis Ward and Polyclinic of Hospital X

10.18196/mmjkk.v21i1.6157

Abstract view : 45 times

Ekorini Listiowati, Fitria Eka Rianti

Correlation between Body Mass Index to Hypertension in A Rural Area in East Java

10.18196/mmjkk.v21i1.7385

Abstract view : 132 times

Alvin Hartanto Kurniawan, Benedictus Hanjaya Suwandi, Dicky Teguh Prakoso, Reynaldy Cliftianto Moksidy, Irma Talitha, Maharani Kartika Anggraeni, Danang Zulkifli

The Density of Dermatophagoides sp. in Households and its Correlation with the Score for Allergic Rhinitis in Jember, East Java

10.18196/mmjkk.v21i1.7925

Abstract view : 51 times

Erma Sulistyaningsih, Elisa Fadia Laili, Cholis Abrori

Comparison of Serum Interleukin 10 Levels between Leprosy and Non-Leprosy Population

10.18196/mmjkk.v21i1.6514

Abstract view : 45 times

Adika Kodrati, Eddy Mart Salim, Zen Hafy

Effects of AntiMycobacterium Tuberculosis Nanoherbal Legundy Leaf Extract (Vitex trifolia)

10.18196/mmjkk.v21i1.6497

Abstract view : 84 times

Atika Dwiyaniti, Siti Mulyani, Oche Axena Yulhan, Humairah Medina Liza Lubis, Ance Roslina

Correlation between Osteoarthritis Knee Damage Based on Ultrasound with Kellgren-Lawrence Classification

10.18196/mmjkk.v21i1.7225

Abstract view : 64 times

Wensri sevni Kurniawati, Muhammad Ilyas, Mirna Muis, Andi Alfian Z, Faridin Faridin, Sri Asriyani

Expression Level of Cyclin-D1 between Endometriomas and Ovarian Carcinomas

[10.18196/mmjkk.v21i1.7212](#)

Abstract view : 41 times

Alfun Dhiya An, Supriyatiningsih Supriyatiningsih

[Correlation between Cup Ratio and Optic Nerve Disc with High Myopia](#)

[10.18196/mmjkk.v21i1.8615](#)

Abstract view : 54 times

Nur Shani Meida, Ahmad Ikliluddin, Yunani Setyandriana, Nadia Dina Ayu



Correlation the Density of *Dermatophagoides sp.* in Households with the Score for Allergic Rhinitis in Jember, East Java

Hubungan Kepadatan *Dermatophagoides sp.* di dalam Rumah dengan Skor Rinitis Alergi di Jember, Jawa Timur

Erma Sulistyaningsih¹, Elisa Fadia Laili², Cholis Abrori²

¹Department of Parasitology, Faculty of Medicine, University of Jember. Jalan Kalimantan 37 Kampus Bumi Tega Boto Jember 68121

² Faculty of Medicine, University of Jember, Jalan Kalimantan 37 Kampus Bumi Tega Boto Jember 68121

DATA OF ARTICLE:

Received: 3 Jan 2020

Reviewed: 25 Jun 2020

Revised: 19 Nov 2020

Accepted: 05 Dec 2020

*CORRESPONDENCE:

sulistyaningsih.fk@unej.ac.id

DOI:

10.18196/mmjkk.v21i1.7925

TYPE OF ARTICLE:

Research

Abstract: The prevalence of allergic rhinitis (AR) in Indonesia is an estimated 24.3% and increasing each year. The majority cause of AR is house dust mites (HDMs) allergens, especially *Dermatophagoides sp.* The screening tool to estimate the prevalence and diagnose AF in a population setting is the Score for Allergic Rhinitis (SFAR) questionnaire. This study aimed to investigate the correlation between the density of *Dermatophagoides sp.* in households and the SFAR in Jember, East Java. The study enrolled 30 housewives as respondents, which were selected by stratified random sampling. Respondents were interviewed using the SFAR questionnaire. The dust sample from each house of the respondent was collected, the HDMs either *Dermatophagoides sp.* or others were identified and the density of *Dermatophagoides sp.* was calculated. Data were analyzed using the Pearson Test. *Dermatophagoides sp.* were found in 21 out of 30 samples and 11 out of 30 respondents had an SFAR of ≥ 7 . The Pearson analysis showed the value of $p = 0.555$ and $r = 0.136$. *Dermatophagoides sp.* was the majority of species of HDMs, but there was no significant correlation between the density of *Dermatophagoides sp.* in households and the SFAR among housewives in Jember.

Keywords: allergic rhinitis; *Dermatophagoides sp.*; house dust mites; SFAR

Abstrak: Prevalensi rinitis alergi (AR) di Indonesia diperkirakan 24,3% dan cenderung meningkat setiap tahun. Penyebab AR mayoritas adalah alergen tungau debu rumah (HDM) terutama *Dermatophagoides sp.* Alat skrining untuk memperkirakan prevalensi dan mendiagnosis AF dalam populasi adalah kuesioner Score for Allergic Rhinitis (SFAR). Penelitian ini bertujuan untuk mengetahui hubungan antara kepadatan *Dermatophagoides sp.* di rumah tangga dan SFAR di Jember, Jawa Timur. Penelitian ini melibatkan 30 ibu rumah tangga sebagai responden yang dipilih secara stratified random sampling. Responden diwawancarai menggunakan kuesioner SFAR. Pengambilan sampel debu dari rumah masing-masing responden. HDMs baik *Dermatophagoides sp.* atau yang lainnya diidentifikasi dan selanjutnya kepadatan *Dermatophagoides sp.* dihitung. Data dianalisis menggunakan Uji Pearson. *Dermatophagoides sp.* didapatkan sebanyak 21 dari 30 sampel yang diambil dan 11 diantara 30 responden memiliki SFAR ≥ 7 . Analisis Pearson menunjukkan nilai $p = 0,555$ dan $r = 0,136$ yang menunjukkan bahwa tidak terdapat korelasi yang bermakna antara kepadatan *Dermatophagoides sp.* dengan SFAR ibu rumah tangga di Jember.

Kata kunci: rhinitis alergi, *Dermatophagoides sp.*, tungau debu, SFAR

INTRODUCTION

Allergic rhinitis (AR) is a symptomatic disease of the nose due to the inflammation of the nose caused by an allergen. It is very common condition in four seasons countries, estimated to affect around one in every five people in the United Kingdom. The prevalence of AR in Latin America was 5.5-45.1%, Europe was 23-30%, Africa was 7.2-54.1%, Australia was 12-41.3% and South-East Asia was 5.5-44.2%.¹ The prevalence of AR in Indonesia variable, it is estimated 24.3% and tends to increase each year.² Although AR is not commonly a severe complaint, it has a negative impact on patients social lives, the school performance of children, and productivity in the workplace.³

The cause of more than 50% of AR in Indonesia is house dust mites (HDM), and *Dermatophagoides pteronyssinus* (Der p1), and *D. farinae* (Der f1) is the most common HDM allergen.⁴ The external part of the body, part of the body and feces of *Dermatophagoides sp.* contains Der p1 and Der f1 allergens which can activate the A and D proteins on the lung surfactant that can inhibit the binding of allergen to IgE. The Der p1 also can induce the production of proinflammatory cytokines through proteinase-activated receptor 2 (PAR2) independent in the respiratory tract and stimulate the development of Th₂, which further activate B cell to produce IgE. IgE and allergen's binding causes degranulation of mast cells and basophil to release histamine, resulting in the symptoms of AR.⁵ The symptom of AR includes the runny or stuffy nose, sneezing, red, itchy and watery eyes, and swelling around the eyes. Exposure to 100-500 HDM/g of dust or 10 mg Der p1/g is a risk factor for an allergic reaction. Exposure to 500 HDM/g of dust in a long time results in IgE-antibody responses.^{6,7,8}

The prevalence of RA in developed and developing countries tends to increase. However, there is limited assessment available to diagnose, especially in population settings, where there are no objective allergy measurements and no health professionals. The Score for Allergic Rhinitis (SFAR) is a quantitative questionnaire developed by experts as a helping tool to diagnose AR. The questionnaire has been validated in three different ways, i.e., diagnosis validation by the medical specialist, internal validation using psychometric methods, and population acceptability methods.⁹ Study in France and six African countries on using the SFAR to diagnose AR reported that the SFAR gives the same or higher positive predictive value and specificity as the International Study of Asthma and Allergies in Childhood (ISAAC) questionnaire. The study also showed that nonspecialists could use the SFAR without performing a physical examination, and the AR diagnosis was nearly matched with the gold standard's diagnosis, confirmed by skin prick test. This questionnaire has eight components, including (1) nasal symptoms in the past year, i.e., sneezing, runny nose and blocked nose; (2) nasal symptoms accompanied with itchy-watery eyes; (3) month of the year that the nasal symptoms occur; (4) triggers of nasal symptoms (pollens and house dust); (5) allergic status; (6) previous diagnosis of allergy; (7) previous positive allergy test; and (8) familial history of allergy. A total score of SFAR is 16, and the cutoff value of ≥ 7 can differentiate individuals with and without AR.^{10,11}

There were some studies on the density of house dust mites, allergic rhinitis and its symptoms in association with the attitude and practice,^{12,13,14} but no study using the SFAR questionnaire to measure the allergic rhinitis, especially among housewives. This study aimed to determine the correlation between the density of *Dermatophagoides sp.* in households and the SFAR among housewives in Jember, East Java.

MATERIAL AND METHOD

Study design and ethical clearance

The study was a cross-sectional study and was conducted in Jember, Indonesia. The study's procedure has been approved by the Ethical Committee of Faculty of Medicine, University of Jember, with reference number 1281/H25.1.11/KE/2019. The sample was chosen by stratified random sampling. We enrolled 30 housewives, which match with inclusion criteria, i.e., 21-34 years old, do not have an outside job, and signed the informed consent after explanation. We excluded women who have a pet such as dogs, cats or birds, and living with a smoking family member. The inclusion criteria for collected house dust were dust from a 1-year-old or older mattress, dust from a mattress that has not been dried in the past week, and dust from a mattress with more than three days of bedsheet replacement. The exclusion criteria were dust from the mattress, which was still wrapped in plastic and made from cotton.

Data collection

We collected dust samples at each house by using a 350 W vacuum cleaner. Dust was collected from a mattress in an area of 1 m² for 2 minutes. The collected dust was placed into a ziplock and labeled with the

respondent's code and sampling date. This study interviewed a housewife at each house using the SFAR questionnaire.

Dust samples were transported to the Laboratory of Parasitology, Medical Faculty, University of Jember. Each dust sample was weighed and recorded. As much as 0.1 g of dust sample was diluted with distilled water and observed under a stereoscopic microscope. Each HDM found was taken using a mite needle and fixed using Hoyer's medium on object glass. The fixed HDM was identified using a light microscope based on Collof key of identification¹⁵ and consulted with the entomologist of the Biology Department, Faculty of Mathematics and Natural Sciences, University of Jember. The identification of HDMs in this study only reached the genus stage. The density of *Dermatophagoides sp.* were calculated using the following formula.¹⁶

$$\text{HDMs density} = \frac{\text{mass of dust (g)}}{0.1} \times \text{HDMs found in 0.1 g of dust}$$

Statistical Analysis

The data was analyzed for its normality using Shapiro-Wilk. Data was transformed once before it had a normal distribution. P-value was used to analyze the significant criteria. The correlation test was performed using Pearson Test

RESULT

There are 30 respondents involved in this study. The result of the interview showed that All job at the time. The majority of respondents were 25-29 years old women (46.67%), as shown in Table 1.

Table 1. The Distribution of Respondents' Age

Age (years)	Total (n)	Percentage (%)
20-24	5	16.67
25-29	14	46.67
30-34	11	36.67

Table 2. Identification of HDMs in Dust Samples, its Density and the SFAR's Score of Respondents

Code	Dust mass (g)	Number of HDM found (n/ 0.1 g of dust)	HDM density (n/ g of dust)	Number of <i>Dermatophagoides sp.</i> found (n/0.1 g of dust)	<i>Dermatophagoides sp.</i> density (n/ g of dust)	SFAR
01	1.3	13	169	1	13	6
02	0.5	8	40	1	5	0
03	0.8	0	0	0	0	0
04	0.8	5	40	2	16	12
05	3.7	5	185	1	37	6
06	0.9	5	45	2	18	7
07	0.3	7	21	0	0	5
08	0.9	14	126	1	9	11
09	0.9	16	144	3	27	11
10	0.4	4	16	0	0	6
11	0.6	15	90	2	12	12
12	0.8	7	56	1	8	10
13	1.0	3	30	1	10	0
14	0.6	3	18	1	6	2
15	1.1	12	132	3	33	6
16	1.2	4	48	1	12	5
17	0.6	6	36	2	12	12
18	1.0	0	0	0	0	2

19	0.7	4	28	1	7	0
20	0.6	0	0	0	0	5
21	0.8	3	24	1	8	0
22	1.0	7	70	3	30	2
23	1.2	3	36	1	12	6
24	0.5	0	0	0	0	5
25	1.1	7	77	0	0	10
26	0.3	6	18	1	3	4
27	0.8	0	0	0	0	7
28	2,1	4	84	1	21	2
29	0,7	0	0	0	0	8
30	0,6	7	42	1	6	15
Total	27.8	168	1575	31	305	
Average	0.93	5.6	52.5	1.03	10.17	5.9

The HDMs identification of dust samples and the SFAR of respondents were shown in Table 2. HDMs were found in 24 out of 30 dust samples, and *Dermatophagoides* sp. were found in 21 out of 24 dust samples with HDMs. The density of *Dermatophagoides* sp. were ranged between 0 - 37 HDM/g of dust, while the average density was 10.17 HDM/g of dust. The identified *Dermatophagoides* sp. were shown in Figure 1. The SFAR of respondents were in the range of 0-15, none of them had the maximum score. As many as 11 respondents had a score of ≥ 7 , meaning had an AR.




Figure 1. Morphology of *Dermatophagoides* sp. found in the sample (a) Female (b) Male

The data of the density of *Dermatophagoides* sp. were not normally distributed, so the data transformation was performed. Data were normally distributed after performing data transformation ($p=0.848$). The test for linearity showed that the correlation between the two variables was linear ($p=0.834$). Pearson Test to determine the correlation between the density of *Dermatophagoides* sp. and the SFAR showed $p = 0.555$ and $r = 0.136$. The result showed no significant correlation between the density of *Dermatophagoides* sp. in households and the SFAR among housewives in Jember (Table 3).

Table 3. The Result of the Statistical Analysis

		Density of <i>Dermatophagoides</i> sp	SFAR
Density of <i>Dermatophagoides</i> sp	Pearson Correlation	1	.136
	Sig. (2-tailed)		.555
	N	21	21
SFAR	Pearson Correlation	.136	1
	Sig. (2-tailed)	.555	
	N	21	30

DISCUSSION

In this study, women were chosen as respondents, as reported by previous studies that women have higher experience of AR than men. This might be due to a different perception of health disorders between women and men, so women had a higher intention to seek professional medical assistance than men.^{2,3,17} All respondents were housewives without an outside job at the time. Housewives tended to get more exposure to indoor allergens such as HDM allergens because of their daily activities such as house cleaning and households furnishings.^{18,19} The majority of respondents were 25-29 years old women (46.67%). The previous study reported that AR's patients were mostly in productive ages, 20-29 years old, and most AR cases were 18-34 years old. AR was rarely found in a person over 60 years old due to aging process which causes the degradation of the allergen-specific level.^{20,21,22}

Houset dust mites were identified in 24 out of 30 dust samples, and *Dermatophagoides sp.* were found in 21 out of 24 dust samples, while the 3 samples were having other HDMs but *Dermatophagoides sp.* Previous studies showed that *Pyroglyphidae* was a family of which dominated the identified HDMs. The most common mites of HDMs are *Dermatophagoides farinae* and *D. Pteronyssinus*. The common HDMs from other families were *Blomia tropicalis*, *Tyrophagus putrescentiae*, and *Aleuroglyphus ovatus*.^{23,24} The density of *Dermatophagoides sp.* were ranged between 0 - 37 HDM/g of dust, while the average density was 10.17 HDM/g of dust. Dust samples were collected from mattresses, where there was an accumulation of human squama on them, which can be used as a food source of HDMs. HDMs used squama as their food source.²⁵

The density of HDMs allergens in households is affected by multiple factors, including the physical environment and the households factors. The physical environment includes latitude, climate, seasonality, rainfall, altitude and distance from the coast. The household factors include the housing design, building age, materials, house orientation, type of furnishings, neighborhood location, ventilation, thermal system, family economic status and the number and behavior of the occupants. Our study was conducted in Jember, it is an area with the temperature range between 23-31°C and relative humidity $\geq 75\%$. These conditions support the growth and development of HDMs.^{7,23,26} The temperature and relative humidity may increase in slums and densely populated areas.²⁷ The negative finding of HDMs in some dust samples possibly be due to the occupants' hygiene behavior. HDMs density can be controlled by the way and frequency of the occupants to clean the house.²⁸ Our study found a lower average HDMs density than the previous studies.^{24,25} We only calculated *Dermatophagoides sp.* density, instead of all species of HDMs.

The SFAR score showed that 11 out of 30 respondents had a score of ≥ 7 , and none of the respondents had the maximum score. None of the respondents experienced nasal symptoms in certain months, known as seasonal AR,²⁸ causing 1 point reduction of the SFAR's score. In four-season countries, nasal symptoms can appear in certain months when pollen season comes. Since Indonesia is a two-season country, it is not easy to find seasonal AR. As we know that the SFAR was developed in France, a four-seasons country, so its application in Indonesia should be modified because the possible different allergens of both countries.

The study used the SFAR questionnaire to determine the AR. The SFAR questionnaire provided better information and more sensitive than the ISAAC questionnaire, which can not distinguish between AR and infectious rhinitis.¹⁰ But, the application of the SFAR questionnaire in Indonesia should be modified since it was established in France, a developed country, which possibly has different AR management from Indonesia. As previous study reported, socio-economic status results in differences in perception, reporting and AR symptoms management.¹⁰ Indonesia has not yet implemented allergic tests such as Skin Prick Test (SPT) or IgE serum routinely in AR's diagnosis algorithm. Hence, many AR patients know the disease without knowing the exact allergens of his/her disease. Respondents who had a total score of 5 or 6 in this study could get a total score of ≥ 7 when carried out an allergy test and had positive results.

Analysis using the Pearson Test showed no significant correlation between the density of *Dermatophagoides sp.* in households and the SFAR among housewives in Jember ($p=0.555$; $r=0.136$). AR is a multifactorial disease caused by the interaction between the complex environmental factors and the genetic factor of individual characters rather than a single factor, such as HDMs density. Each individual has a different level of sensitization and atopic person possesses a complex and subjective immune system.^{9,23,29} *Blomia tropicalis* was another species which dominantly found in previous studies. Its allergens can cross-react with *Dermatophagoides sp.* allergens, although the incidence reported was relatively low. *Lepydoglyphus sp.*, *B. tropicalis*, *B. kulagini*, *Acarus siro*, and *Glycyphagus sp.* were found in several Indonesia studies. Previous studies showed that AR patients were sensitive to some of those species' allergens. The insignificant correlation between variables could be due to another HDMs allergens that affect the AR symptoms as identified by the SFAR.^{22,30,31} Some HDMs were found to have incomplete body parts, made difficulties in identification process, whether *Dermatophagoides sp.* or other species. Eggs, larvae, and

nymphs were also found in dust samples. However, it was difficult to identify its morphological characteristic as *Dermatophagoides* sp. or other species. Therefore, they were not identified in this study. Those factors were the limitation of this study which may affect the results.

CONCLUSION

This study revealed no significant correlation between the density of *Dermatophagoides* sp in households and the SFAR score among housewives in Jember, East Java.

ACKNOWLEDGMENTS

The author would like to thank all housewives who consented to participate in this study

REFERENCES

1. European Academy of Allergy and Clinical Immunology. *Global Atlas of Allergic Rhinitis and Chronic Rhinosinusitis*. 2015. Zurich: EAACI Headquarters.
2. Kemenkes RI. Riset Kesehatan Dasar. *Badan Penelitian dan Pengembangan Kemenkes RI* Jakarta. 2017. p. 118.
3. Bachet C, van Cauwenberge P & Khaltaev N. Allergic Rhinitis and its Impact on Asthma. In Collaboration with the World Health Organization. Executive Summary of the workshop report, 7-10 December 1999, Geneva, Switzerland. *Allergy*, 2002; 57: 841-855. <https://doi.org/10.1034/j.1398-9995.2002.23625.x>
4. Habsari JT, Aryati & Pawarti DR. Perbandingan Nilai Diagnostik IgE Spesifik Tungau Debu Rumah, Metode ELISA, dan Immunoblot pada Rinitis Alergi. *Indonesian J Clin Pathol Med Lab*. 2016; 22 (2): 119-126. <https://doi.org/10.24293/ijcpml.v22i2.1113>
5. Dullaers M, De Bruyne R, Ramadani F, Gould HJ, Gevaert P & Lambrecht BN. The Who, Where, and When of IgE in Allergic Airway Disease. *J Allergy Clin Immunol*. 2012; 129 (3): p. 635-645. <https://doi.org/10.1016/j.jaci.2011.10.029>
6. Portnoy J, Miller JD, Williams PB, Chew GL, Miller JD, Zaitoun F. Environmental Assessment and Exposure Control of Dust Mites: a Practice Parameter. *Ann Allergy Asthma Immunol*. 2013; 111 (6): p. 465-507. <https://doi.org/10.1016/j.anai.2013.09.018>
7. Natalia D. Peranan Alergen Tungau Debu Rumah (Der p1 dan Der p2) dalam reaksi alergi. *CDK*; 2015; 42 (4): p. 251-255.
8. Mantu, B.G., Wahongan, G.J. & Bernadus, J.B. Hubungan Kepadatan Tungau Debu Rumah dengan Derajat Rinitis Alergi. *eBiomedik*. 2016; 4 (1). <https://doi.org/10.35790/ebm.4.1.2016.11056>
9. Annesi-Maesano I, Didier A., Klossek M, Chanal I, Moreau D. & Bousquet J. The Score for Allergic Rhinitis (SFAR): a Simple and Valid Assessment Method in Population Studies. *Allergy: European J Allergy and Clin. Immunol*. 2002; 57 (2): 107-114. <https://doi.org/10.1034/j.1398-9995.2002.103170.x>
10. Piau JP, Massot C, Moreau D, Ait-Khaled N, Bouayad Z, Mohammad Y, Khaldi F, et al. Assessing Allergic Rhinitis in Developing Countries. *IJTLD*. 2010; 14 (4): 506-512.
11. Ologe FE, Adebola SO, Dunmade AD, Adeniji, KA & Oyejola BA. Symptom Score for Allergic Rhinitis. *J Otolaryngol Head Neck Surg*. 2013 April; 148 (4): 557-563. <https://doi.org/10.1177/0194599813477605>
12. Purba ISE, Pijoh VD, Runtuwene J. Survey Perilaku Masyarakat terhadap Populasi Tungau Debu Rumah di Kelurahan Titiwungen Selatan Kecamatan Sario Kota Manado. *Jurnal e-Biomedik*. 2013 Mar; 1 (1): p. 319-324. <https://doi.org/10.35790/ebm.1.1.2013.4365>
13. Mantu BG, Wahongan GJ, Bernadus JB. Hubungan Kepadatan Tungau Debu Rumah dengan Derajat Rhinitis Alergi. *Jurnal e-Biomedik*. 2016 Jan; 4 (1). <https://doi.org/10.35790/ebm.4.1.2016.11056>
14. Yolazenia Y. Harianto H, Riady IT. Gambaran kepadatan tungau debu rumah sebagai pencetus rhinitis alergi pada anak panti asuhan Pekanbaru. *Jurnal Ilmu Kedokteran*. 2019 Mar; 13 (1): 23-27. <https://doi.org/10.26891/JIK.v13i1.2019.23-27>
15. Colloff MJ. *Dust mites*. Dordrecht: Springer Netherland; 2009. <https://doi.org/10.1071/9780643100497>
16. Kawulur YC, Tuda JS, dan Wahongan GJ. Jenis dan Kepadatan Tungau Debu Rumah yang Ditemukan di Kelurahan Teling Bawah Kecamatan Wenang Kota Manado. *eBiomedik*. 2013; 1 (3): 1081-1084. <https://doi.org/10.35790/ebm.1.3.2013.3263>
17. Rambe AY, Fadhli M, Munir D, Haryuna, TSH, dan Eyanoe PC. Hubungan Rinitis Alergi dan Disfungsi Tuba Eustachius dengan menggunakan Timpanometri. *ORLI*. 2013; 43 (1): p. 80-89. <https://doi.org/10.32637/orli.v43i1.20>
18. Novitasari, Sorisi A, dan Wahongan GJP. Profil Penderita Alergi dengan Hasil Skin Prick test TDR Positif di Poliklinik Alergi-Imunologi RSUP Prof. Dr. R. D. Kandou Manado periode 2007-2009. *eBiomedik*. 2013; 1 (2): p. 1014-1018. <https://doi.org/10.35790/ebm.1.2.2013.3254>

19. Reinhard E, Palandeng OI, dan Pelealu OC. Rinitis alergi di Poliklinik THT-KL BLU RSU Prof. dr. RD Kandou Manado periode Januari 2010–Desember 2012. *e-CliniC*. 2013; 1 (2). <https://doi.org/10.35790/ecl.1.2.2013.3280>
20. Susanti E, Pawarti DR, dan Soeprijadi S. Hubungan Kadar RANTES Sekret Hidung dengan Skor Gejala Total Penderita Rinitis Alergi. *ORLI*. 2016; 46 (2): p. 110-120. <https://doi.org/10.32637/orli.v46i2.158>
21. Rahmawati N, Suprihati S, dan Muyassaroh M. Faktor Risiko yang Mempengaruhi Disfungsi Tuba eustachius pada Penderita Rinitis Alergi Persisten. *ORLI*. 2011; 41 (2): 142-146. <https://doi.org/10.32637/orli.v41i2.51>
22. Cazzoletti L, Ferrari M, Olivieri M, Verlato G, Antonicelli L, Bono R., et al. The Gender, Age and Risk Factor Distribution Differs in Self-reported Allergic and Non-allergic Rhinitis: a Cross-sectional Population-based Study. *Allergy Asthma Clin Immunol*. 2015; 11 (1): 36-44. <https://doi.org/10.1186/s13223-015-0101-1>
23. Yu JM, Luo QH, Sun JL, Shi CL, Yin J, Zhou YL., et al. Diversity of House Dust Mite Species in Xishuangbanna Dai: a Tropical Rainforest Region in Southwest China. *BioMed research international*. 2015; p. 1-6. <https://doi.org/10.1155/2015/421716>
24. Ponggalunggu WF, Pijoh VD, & Wahongan GJ. Jenis dan Kepadatan Tungau Debu Rumah pada Beberapa Habitat di Rumah Penderita Penyakit Alergi. *eBiomedik*. 2015; 3 (1): p. 254-260. <https://doi.org/10.35790/ebm.3.1.2015.6734>
25. De Breving RFR., Tuda JSB & Wahongan GJP. Tungau Debu Rumah yang Ditemukan di Kelurahan Perkamil Kecamatan Paal 2 Kota Manado. *eBiomedik*. 2013; 1 (2): p. 859-862. <https://doi.org/10.35790/ebm.1.2.2013.5471>
26. Badan Perencanaan Pembangunan Daerah Provinsi Jawa Timur. *Potensi dan Produk Unggulan Jawa Timur*. 2013. Surabaya: Badan Perencanaan Pembangunan Daerah Provinsi Jawa Timur.
27. Badan Pusat Statistik Kabupaten Jember. *Kecamatan Sumbersari dalam Angka*. Jember: Badan Pusat Statistik Kabupaten Jember. 2018.
28. Soltani A, Azizi K, Saleh V & Dabaghmanesh T. The Fauna and Distribution of House Dust Mites in Residential Homes of Bandar Abbas District, Southern Iran. *Exp Appl Acarol*. 2011; 54 (3): p. 269-276. <https://doi.org/10.1007/s10493-011-9436-6>
29. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens W, Togias A, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA2LEN, and AllerGen). *Allergy*. 2008; 63 (Suppl. 86): p. 8-160.
30. Haqi, S.A. 2016. *Prevalensi dan Derajat Infestasi Tungau Debu Rumah di Asrama Mahasiswa* [disseration]. Bogor: Institut Pertanian Bogor.
31. Sade K, Roitman D & Kivity S. Sensitization to Dermatophagoides, Blomia tropicalis, and other mites in atopic patients. *J Asthma*. 2010; 47 (8): 849-852. <https://doi.org/10.3109/02770903.2010.506683>