

PUBLIKASI JURNAL


**Study on Activity of Cocoa Ethanolic Extract
Against *Shigella dysenteriae***

Dr. dr. Enny Suswati, MKes

NIP 197002141999032001

. Staf Pengajar Lab. Mikrobiologi

. Fakultas Kedokteran Universitas Jember



**KEMENTERIAN PENDIDIKAN, KEBUDAYAAN,
RISET DAN TEKNOLOGI
UNIVERSITAS JEMBER**

Karya Ilmiah dipublikasikan pada:

Jurnal Pelita Perkebunan

Volume 32 No. 2 2016 Hal 130-137

ISSN: 0215-0212 / e-ISSN: 2406-9574

Accredited by Degree of LIPI Chairman
Nomor 638/AU3/P2MI-LIPI/07/2015


ISSN 0215-0212
e-ISSN 2406-9574

Pelita **PERKEBUNAN**

A COFFEE AND COCOA RESEARCH JOURNAL

Volume 32 Number 2

August 2016



INDONESIAN COFFEE AND COCOA RESEARCH INSTITUTE (ICCRI)

Pelita Perkebunan	Vol. 32	No. 2	PP. 83 - 149	Jember August 2016	ISSN 0215-0212 e-ISSN 2406-9574
-------------------	---------	-------	--------------	-----------------------	------------------------------------

CONTENT

	Page
● Determination of selection index of cocoa (<i>Theobroma cacao</i> L.) yield traits using regression methods. Bayu Setyawan, Taryono and Suyadi Mitrowihardjo	83–90
● Abundance of pollinator insect (<i>Forcipomyia</i> spp.) of cocoa under some shade trees. Fakhrusy Zakariyya, Endang Sulistyowati and Dwi Suci Rahayu	91–100
● Application of copper fungicide and row covers to control <i>Phytophthora</i> seedling blight disease on cocoa. Febrilia Nur ‘Aini, Teguh Iman Santoso and Sudarsianto	101–108
● Red kidney bean powder substituted milk in cinnamon herbal coffee: Consumer perception, sensory properties and nutrition content. Noor Ariefandie Febrianto, Khalimatus Sa’diyah and Tejasari	109–119
● Performance of roasted cocoa bean winnower for smallholder chocolate producers. Hendy Firmanto, Lya Aklimawati and Bayu Setyo Abdurrizal	120–129
● Study on activity of cocoa ethanolic extract against <i>Shigella dysenteriae</i> . Ariza Budi Tunjung-Sari, Enny Suswati, Diana Chusna Mufida, Alfa Rianul Setiawan, Alvin Isnaini, Alfa Rianul Setiawan, Mochamad Rizal, Teguh Wahyudi and Misnawi	130–137
● Factors affecting micro and small-scale industries income: A case study on coffee processing industry. Lya Aklimawati, Djoko Soemarno and Surip Mawardi	138–148
● Invited reviewers of Pelita Perkebunan, Volume 32, Number 2, August 2016 Edition	149

Study on Activity of Cocoa Ethanolic Extract Against *Shigella dysenteriae*

Ariza Budi Tunjung-Sari^{1*}, Enny Suswati², Diana Chusna Mufida², Alfa Rianul Setiawan², Alvin Isnaini², Alfa Rianul Setiawan², Mochamad Rizal², Teguh Wahyudi¹, and Misnawi¹

¹Indonesian Coffee and Cocoa Research Institute (ICCR), Jl. PB Sudirman 90, Jember 68118, East Java, Indonesia.

²Faculty of Medicine, Jember University, Jl. Kalimantan 37, Jember 68121, East Java, Indonesia.

*Corresponding author: arizabudi@yahoo.com

Received: 16 June 2016 / accepted: 24 July 2016

Abstract

Shigella dysenteriae is a gastrointestinal pathogen which shows resistance to antibiotics. A study has been conducted to investigate alternative antibacterial agents, due to the emerging resistance of *S. dysenteriae* to ciprofloxacin and other antibiotic classes. In this study, antibacterial properties of cocoa ethanolic extract (CEE) and its impact on growth and morphology of *S. dysenteriae* were evaluated. The effect of CEE on bacterial growth was assayed by using agar-well diffusion method and by observing morphological changes of bacterial cells through the use of scanning electron microscopy (SEM). Furthermore, CEE was also applied orally to mice infected with *S. dysenteriae*. The intestinal fluids was cultured in selective medium to evaluate growth of *S. dysenteriae* colonies. This study demonstrated that CEE at concentrations of 15.6 mg/mL inhibited *S. dysenteriae* growth, and at concentrations of 500 mg/mL and 1,000 mg/mL exhibited equal activity to 6.5 µg/mL of ciprofloxacin. SEM showed that *S. dysenteriae* cells had formed filaments, indicating that CEE caused cellular stress to *S. dysenteriae*. In *in vivo* assay, CEE showed suppression of *S. dysenteriae* colony in the mice intestine. This research suggests that CEE could potentially be used as antibacterial agent againsts *S. dysenteriae*.

Keywords: antibacterial, dysentery, cocoa, active compound, antibiotic

INTRODUCTION

The World Health Organization (WHO) reported that diarrhea killed 1.9 million people each year, including 760,000 children under five-year of age. In fact, the highest incidence of diarrhea was among infants 6-11 months of age, who typically suffered 4.5 diarrhea incidents per year (Walker *et al.*, 2012). Among other diarrheal pathogens, *Shigella dysenteriae* produces the most severe disease and complications, because its Type-1 species releases toxins.

Instead of vaccines, *Shigella* infection is treated by using antibiotics particularly from the class of β -lactams, quinolones or

macrolides (Christopher *et al.*, 2010). Among these, ciprofloxacin from quinolone class is the first line of treatment, while pivmecillinam, ceftriaxone and azithromycin are the most used second options (Christopher *et al.*, 2010; WHO, 2005). However, Shiferaw *et al.* (2012) has reported resistance of *Shigella* towards ampicillin, streptomycin, trimethoprim-sulfamethoxazole, sulfamethoxazole-sulfisoxazole, and tetracycline, and decreasing susceptibility against ciprofloxacin and nalidixic acid. Alternative antimicrobial agents for *Shigella* has been investigated from plant extract, for instance garlic and allicin (Andualem, 2013; Gull *et al.*, 2012). Native medicinal plants, such as *Channa striatus* and *Picralima nitida*

also demonstrated promising antimicrobial activity (Haniffa *et al.*, 2013; Kouitcheu *et al.*, 2013).

Cocoa bean (*Theobroma cacao* L.) was anciently utilized as a medicine, despite the current use for confectioneries. It is a valuable commodity for the Aztecs due to its nutrition, energetic, and aphrodisiac benefits. Cocoa beans were used to treat stomach and intestinal pains including dysenteriae as described in “Florentine Codex,” a record of indigenous Aztec medical practices (Dillinger *et al.*, 2000; Lippi, 2009). In recent times, health properties of cocoa beans have been extensively investigated, particularly for the prevention of cardiovascular diseases. We have found antibacterial properties against *Escherichia coli*, *Staphylococcus aureus* and *Staphylococcus epidermidis*, as well as immunomodulatory and anticancer properties (Sari *et al.*, 2015; Sari *et al.*, 2014; Sari *et al.*, 2016). This study investigated the effects of cocoa ethanolic extract (CEE) on *Shigella dysenteriae* employing *in vitro* assay scanning electron microscopy (SEM) and *in vivo* testing in mice.

MATERIALS AND METHODS

Six kilograms of cocoa beans dried under the sun without prior fermentation, were ground into five-mesh particle size. The paste was pressed at 80–90°C to remove cocoa butter, and was soaked overnight in n-hexane for thorough fat removal. The defatted mass was then filtered to remove liquid and the solid fraction was soaked in ethanol overnight. Finally, the ethanol extract was concentrated and spray dried to produce powder. Total polyphenol content (TPC) was measured using Folin-Ciocalteu method and was expressed as catechin equivalent and gallic acid equivalent.

Assessment of antibacterial activity was carried out by performing agar-well diffusion assay (Rojas *et al.*, 2006). The CEE solution was prepared by diluting in sterile distilled water to create concentrations of 1,000; 500; 250; 125; 62.5; 31.3; 15.6 and 7.8 mg/mL. Positive control suspension was prepared from ciprofloxacin 6.5×10^{-3} mg/mL, while negative control was prepared from sterile distilled water. Culture of *S. dysenteriae* (1×10^8 CFU/mL) were streaked on plates containing Mueller Hinton agar. The agar was perforated to form well (diameter 10 mm, 4 wells/plate) and the CEE solutions were poured into each well. Plates were incubated at 37°C for 24 hours, and development of clear area surrounding each well was observed. The inhibition zone was visible as a clear area surrounding each well. The inhibition zone diameter (IZD) was the diameter of the visible clear zone subtracted from the well diameter. The relative inhibition zone diameter (RIZD) of CEE was $RIZD_{CEE}(\%) = [(IZD_{CEE} - IZD_{\text{Negative control}}) / IZD_{\text{positive control}}] \times 100\%$ (Rojas *et al.*, 2006). The assay was done triplicate.

To observe morphological changes of the bacteria cells, 10 µL of *S. dysenteriae* culture was added to vials containing CEE at concentrations of 7.8 mg/mL, 15.6 mg/mL, and 31.2 mg/mL, and was incubated at 35°C for 20 hours. The bacterial cells were fixated in glutaraldehyde 2% at 4°C, and were immersed in osmic acid 1% at 4°C and were subsequently dehydrated with ethanol. Preservation of the cells were made by using absolute amyl acetate solution. The cells then were dried on glass prior to gold glazing and microscopic observation process.

Male Swiss-Webster mice, aged 2–3 months with average weight of 25–30 g underwent acclimatization for one week prior to treatment. Mice were infected through intraperitoneal

route with 1.5 mL suspension of *S. dysenteriae* culture (10^5 CFU/mL). Twenty four hours post infection, mice orally received CEE (0.65; 1.3; and 2.6 mg/mL), ciprofloxacin (1.3 mg/mL), or sterile water (1mL). Application was done twice a day for seven consecutive days. On the 8th day, mice were sacrificed and the intestine wash was cultured in SS agar. After incubation for 20 days, colonies of *S. dysenteriae* shown in agar was counted. This trial has acquired ethical approval from committee from Faculty of Medicine, Jember University.

Experiment was done in triplicates and statistical analysis of the data used t-test, Kruskal-Wallis analysis, and Mann-Whitney test for post-hoc analysis.

RESULTS AND DISCUSSION

Secondary metabolites of plants have attracted studies on their antibacterial potency. Among these, polyphenol is considered as the most potent antibacterial agent, since its function in plant defense against predators. Polyphenol content in cocoa beans is higher than in fruits, such as pomegranates, cranberries, blueberries and acai berries (Crozier *et al.*, 2011). Compared to other beverages, cocoa provides more phenolic content per serving than coffee, tea and wine. This studies showed that total phenolics (gallic acid equivalent) and flavonoids (epicatechin equivalents) in cocoa per serving were about two to three fold greater than that of black tea, green tea and red wine (Lee *et al.*, 2003). Polyphenol accumulates in the nonfat solids of cocoa bean, thus the concentration is higher in chocolate products with less added ingredients. Chocolate, for instance, has less polyphenol content than cocoa powder and cocoa liquor (Gu *et al.*, 2006).

Cocoa polyphenol is composed largely of monomers, with a smaller number of trimers and tetramers (Misnawi *et al.*, 2002). Extract of cocoa bean may contain tannin, (-)-epicatechin and (+)-catechin, and some of phenolic acids, flavonoids (quercetin, kaempferol, naringenin, myricetin, luteolin, and apigenin) and alkaloids (caffeine, theophylline and theobromine) (Elwers *et al.*, 2009; Ortega *et al.*, 2008; Sánchez Rabaneda *et al.*, 2003). The alkaloids are distributed in various parts of cocoa (bean, pulp, and shell), where the amount is gradually decreasing during fermentation (Brunetto *et al.*, 2007). Proportion between theobromine and caffeine content is different in each cocoa variety, whereas Criollo bean has lower theobromine/caffeine concentration, while Forastero bean has high theobromine/caffeine concentration (Brunetto *et al.*, 2007).

Fermentation of cocoa beans may reduce bitterness and astringency caused by polyphenol, hence its concentration. Misnawi *et al.* (2003) demonstrated that fermentation not only reduced polyphenol concentration but also altered the composition of anthocyanin, epicatechin and catechin. Alkalization step also degrades polyphenols in cocoa. By mixing cocoa powder with an alkaline solution (a practice known as “Dutch process”), one will produce a dark, yet mild tasting, cocoa powder. Degree of alkalization may be determined based on cocoa appearance, as light alkalization converts natural cocoa powder colors from light brown to red/brown and heavy alkalization results in dark red/black colors (Miller *et al.*, 2008).

In this experiment, cocoa ethanolic extract (CEE) was produced from unfermented, unroasted, natural cocoa powder. The CEE contained TPC 140 mg/g (+)-catechin equivalent and 397 mg/g (gallic acid equivalent),

amounts that are almost 10-fold greater than in natural roasted cocoa powder (TPC 40.59–63.20 mg/g) and 40-fold greater than in heavily alkalized powder (TPC 9.54 mg/g) (Miller *et al.*, 2008).

In *in vitro* assay, the inhibition zone diameter (IZD) of CEE was ranging from 0.4 to 11.1 mm. After applying regression on IZD towards CEE concentration, a logarithmic response was observed ($R^2 = 0.9675$). This experiment indicated that antibacterial activity was concentration-dependent, however the concentration higher than 500 mg/mL may bring less significant effect. A comparable effect was shown from CEE at 500 mg/mL, 100 mg/mL and ciprofloxacin 6.5×10^{-3} mg/mL ($P > 0.05$). On the other hand, there was no significant difference between the effect of CEE at 7.8 mg/mL and sterile water, indicating that in this concentration CEE was not effective as an antibacterial agent. RIZD, the ratio of IZD between the extract and positive control, is expressing the antimicrobial activity of CEE relative to ciprofloxacin (Rojas *et al.*, 2006). A strong antibacterial activity was shown by CEE at concentrations between 500–1,000 mg/mL, which featured a RIZD of 97–106% (Table 1).

The polyphenol in CEE may perform bacteriostatic and bactericidal activities through several actions, including disturbance of bacterial cell metabolism, inhibition of cell wall synthesis, disruption of cell membrane permeability, and interference with nucleic acid and gene expression (Cui *et al.*, 2012; Denyer & Maillard, 2002). Tannin works primarily on the bacterial outer membrane, since its protein-precipitation ability may interrupt membrane continuity (Liu *et al.*, 2013). Research on green tea polyphenol, (-)-epigallocatechin-3-gallate (EGCG), demonstrated destructive activity toward *E. coli*, by inducing leakage and providing gateways to the cell interior (Cui *et al.*, 2012). In addition to polyphenols, antibacterial activity of cocoa extract may be due to the presence of methylxanthines. Caffeine and derivate of theophylline were reported to decrease growth of gram-negative bacteria (Al-Janabi, 2011). Particularly, caffeine stimulates bacteria cell to undergo morphological changes and lysis (Dash & Gummadi, 2008).

Susceptibility of gram-negative or gram-positive bacteria is different, depending on characteristics of the cell wall. Gram-negative cells are equipped with a lipopolysaccharide

Table 1. Inhibition zone diameter (IZD) of CEE in various concentrations (mg/mL)

Treatment	Concentration (mg/mL)	IZD*, mm	RIZD*, %
CEE	7.8	0.0 ± 0.0 ^a	0
	15.6	0.4 ± 0.3 ^c	3
	31.2	2.3 ± 0.4 ^d	22
	62.5	5.1 ± 0.2 ^e	48
	125	8.1 ± 0.6 ^f	77
	250	9.0 ± 0.3 ^g	86
	500	10.1 ± 0.6 ^h	97
	1,000	11.1 ± 0.3 ^b	106
Ciprofloxacin	6.5×10^{-3}	10.5 ± 0.5 ^b	100
Sterile water		0 ± 0 ^a	0

Notes: Data are means of three replications. Data in the same column followed by the same letter are not significantly different based on Mann-Whitney post hoc ($\alpha = 0.05$).

outer membrane which determines virulence and resistance. Studies on tea (+)-catechin reported low bactericidal activity against gram-negative bacteria due to the negative charge of lipopolysaccharides (Kajiya *et al.*, 2004). Nevertheless, the contact between proanthocyanidins and lipopolysaccharides may disrupt cellular homeostasis (Johnson *et al.*, 2008).

SEM observation found that the *S. dysenteriae* cell in sterile water (negative control) was measured 2.04 μm in length. After CEE exposure for 20 hours, the bacterial cells extended and formed filaments. Filamentation began at the lowest concentration of CEE, 7.8 mg/mL (4.89 μm), and continued at concentrations of 15.6 mg/mL (6.39 μm) and 31.2 mg/mL (10.61 μm). The extended length of the cells indicates that the bacteria is receiving stress factors. Bacterial stress stimuli may include an unfavorable environment, nutritional imbalance, overpopulation, host immunity, and presence of toxins and antibiotics (Justice *et al.*, 2008; Laureti *et al.*, 2013; Poole, 2012). In response to stimuli, the stress response system (SOS response) is activated. This response system

leads to cell shape modification, biofilm formation, mutation, and shifted virulence (Poole, 2012). When a DNA damage occurs, bacteria will delay mitosis to prevent transfer of defective DNA to the new cells. This condition leaves cells extend without cleaving, thus manifest in form of filaments (Justice *et al.*, 2008).

In the *in vivo* assay, antibacterial potency was indicated from the ability of test sample to suppress the growth of the *S. dysenteriae* colony in the mice intestine. There were significant differences in number of *S. dysenteriae* colonies between sterile water and CEE treatments, except for CEE at the lowest dose (0.56 mg/mL). A similar effect was shown after application of CEE in equal dose with ciprofloxacin (1.3 mg/mL, $p > 0.05$). CEE at the highest dose (2.6 mg/mL) had significantly reduced the colony number (Figure 1).

Interestingly, while in the *in vitro* assay the effective dose of CEE was 7×10^4 -fold from ciprofloxacin (CEE 500 mg/mL vs. ciprofloxacin 6.5×10^{-3} mg/mL), in the *in vivo* assay the equal dose between CEE and ciprofloxacin had produced comparable effects. Doubling the dose of CEE from that




Figure 1. Number of *S. dysenteriae* colonies from mice intestine. CEE 1 = 0.56 mg/mL, CEE 2 = 1.3 mg/mL, CEE 3 = 2.6 mg/mL, and Cipro = ciprofloxacin 1.3 mg/mL as positive control. Data are means of three replications (bars indicated standard deviation of means)

of ciprofloxacin had significantly suppressed the colony growth. This indicates that CEE is not only attributed with antibacterial activity, but also with immuno-stimulation properties. Some studies revealed that cocoa polyphenols may induce changes composition and activity of immune cells as well as immune response. Ramiro-Puig *et al.* (2008) reported that a 10% cocoa diet had changed lymphocyte composition in peyer's patches and mesenteric lymph nodes. In addition, Vázquez-Agell *et al.* (2011) demonstrated through a clinical study that cocoa consumption had acutely decreased the concentration of NF- κ B, the signaling molecule mediating inflammation.

In an attempt of replacing conventional treatments, we could try to apply both CEE and an antibiotic, to utilize the benefits of both treatments. Similarly, studies have shown that a combination of (-)-epigallocatechin-3-gallate (EGCG) and cefotaxime induced permanent damage in an *E. coli* cells, though those results have not been achieved by EGCG alone (Cui *et al.*, 2012).

CONCLUSIONS

Antibacterial activity against *S. dysenteriae* was demonstrated by CEE, which had inhibited growth and triggered cellular stress. The effective dose in the in vivo assay was 1.3 mg/mL which equals to ciprofloxacin. This indicates that CEE may also stimulate immune system. This research suggests that CEE could potentially be used as a treatment of *S. dysenteriae* infection, though further investigation is necessary.

REFERENCES

Al-Janabi, A.A.H.S. (2011). Potential activity of the purine compounds caffeine and aminophylline on bacteria. *Journal of Global Infectious Diseases*, 3, 133–137.

- Andualem, B. (2013). Synergistic antimicrobial effect of Tenegn honey (*Trigona iridipennis*) and garlic against standard and clinical pathogenic bacterial isolates. *International Journal of Microbiological Research*, 4, 16–22.
- Brunetto, M.a.d.R.; L. Gutiérrez; Y. Delgado; M. Gallignani; A. Zambrano; Á. Gómez; G. Ramos & C. Romero (2007). Determination of theobromine, theophylline and caffeine in cocoa samples by a high-performance liquid chromatographic method with on-line sample cleanup in a switching-column system. *Food Chemistry*, 100, 459–467.
- Christopher, P.; K.V. David; S.M. John & V. Sankarapandian (2010). Antibiotic therapy for *Shigella dysentery*. *Cochrane Database of Systematic Reviews*, 8, 1–101.
- Crozier, S.J.; A.G. Preston; J.W. Hurst; M.J. Payne; J. Mann; L. Hainly & D.L. Miller (2011). Cacao seeds are a “Super Fruit”: A comparative analysis of various fruit powders and products. *Chemistry Central Journal*, 5, 1–6.
- Cui, Y.; S.H. Kim; H. Kim; J. Yeom; K. Ko; W. Park & S. Park (2012). AFM probing the mechanism of synergistic effects of the green tea polyphenol (-)-epigallocatechin-3-gallate (EGCG) with cefotaxime against extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli*. *Public Library of Science One*, 7, 1–6.
- Dash, S.S. & S.N. Gummadi (2008). Inhibitory effect of caffeine on growth of various bacterial strains. *Research Journal of Microbiology*, 3, 457–465.
- Denyer, S. & J.Y. Maillard (2002). Cellular impermeability and uptake of biocides and antibiotics in Gram negative bacteria. *Journal of Applied Microbiology*, 92, 35–45.
- Dillinger, T.L.; P. Barriga; S. Escárcega; M. Jimenez; D.S. Lowe & L.E. Grivetti (2000). Food of the gods: cure for humanity? A cultural history of the medicinal and ritual use of chocolate. *Journal of Nutrition*, 130, 2057–2072.

- Elwers, S.; A. Zambrano; C. Rohsius & R. Lieberei (2009). Differences between the content of phenolic compounds in Criollo, Forastero and Trinitario cocoa seed (*Theobroma cacao* L.). *European Food Research and Technology*, 229, 937–948.
- Gu, L.; S.E. House; X. Wu; B. Ou & R.L. Prior (2006). Procyanidin and catechin contents and antioxidant capacity of cocoa and chocolate products. *Journal of Agricultural and Food Chemistry*, 54, 4057–4061.
- Gull, I.; M. Saeed; H. Shaukat; S.M. Aslam; Z.Q. Samra & A.M. Athar (2012). Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. *Annals of Clinical Microbiology and Antimicrobials*, 11, 1–6.
- Haniffa, M.; P. Jeyasheela & M.J. Milton (2013). In vitro antibacterial activity of tissue extracts from four channids against enteric pathogens. *Journal of Agricultural Technology*, 9, 1437–1445.
- Johnson, B.J.; J.B. Delehanty; B. Lin & F.S. Ligler (2008). Immobilized proanthocyanidins for the capture of bacterial lipopolysaccharides. *Analytical Chemistry*, 80, 2113–2117.
- Justice, S.S.; D.A. Hunstad; L. Cegelski & S.J. Hultgren (2008). Morphological plasticity as a bacterial survival strategy. *Nature Reviews Microbiology*, 6, 162–168.
- Kajiya, K.; H. Hojo; M. Suzuki; F. Nanjo; S. Kumazawa & T. Nakayama (2004). Relationship between antibacterial activity of (+)-catechin derivatives and their interaction with a model membrane. *Journal of Agricultural and Food Chemistry*, 52, 1514–1519.
- Kouitcheu, L.B.; J.L. Tamesse & J. Kouam (2013). The anti-shigellosis activity of the methanol extract of *Picralima nitida* on *Shigella dysenteriae* type I induced diarrhoea in rats. *BMC Complementary and Alternative Medicine*, 13, 1–11.
- Laureti, L.; I. Matic & A. Gutierrez (2013). Bacterial responses and genome instability induced by subinhibitory concentrations of antibiotics. *Antibiotics*, 2, 100–114.
- Lee, K.W.; Y.J. Kim; H.J. Lee & C.Y. Lee (2003). Cocoa has more phenolic phytochemicals and a higher antioxidant capacity than teas and red wine. *Journal of Agricultural and Food Chemistry*, 51, 7292–7295.
- Lippi, D. (2009). Chocolate and medicine: Dangerous liaisons? *Nutrition*, 25, 1100–1103.
- Miller, K.B.; W.J. Hurst; M.J. Payne; D.A. Stuart; J. Apgar; D.S. Sweigart & B. Ou (2008). Impact of alkalization on the antioxidant and flavanol content of commercial cocoa powders. *Journal of Agricultural and Food Chemistry*, 56, 8527–8533.
- Misnawi; S. Jinap; B. Jamilah & S. Nazamid (2003). Effects of incubation and polyphenol oxidase enrichment on colour, fermentation index, procyanidins and astringency of unfermented and partly fermented cocoa beans. *International Journal of Food Science & Technology*, 38, 285–295.
- Misnawi; J. Selamat; J. Bakar & N. Saari (2002). Oxidation of polyphenols in unfermented and partly fermented cocoa beans by cocoa polyphenol oxidase and tyrosinase. *Journal of the Science of Food and Agriculture*, 82, 559–566.
- Ortega, N.d.; M.-P. Romero; A. Maciá; J. Reguant; N. Anglés; J.-R.n. Morello & M.-J. Motilva (2008). Obtention and characterization of phenolic extracts from different cocoa sources. *Journal of Agricultural and Food Chemistry*, 56, 9621–9627.
- Poole, K. (2012). Bacterial stress responses as determinants of antimicrobial resistance. *Journal of Antimicrobial Chemotherapy*, 67, 2069–2089.
- Ramiro-Puig, E.; F.J. Pérez-Cano; S. Ramos-Romero; T. Pérez-Berezo; C. Castellote; J. Permanyer; À. Franch; M. Izquierdo-

- Pulido & M. Castell (2008). Intestinal immune system of young rats influenced by cocoa-enriched diet. *The Journal of Nutritional Biochemistry*, 19, 555–565.
- Rojas, J.J.; V.J. Ochoa; S.A. Ocampo & J.F. Muñoz (2006). Screening for antimicrobial activity of ten medicinal plants used in colombian folkloric medicine: A possible alternative in the treatment of non-nosocomial infections. *BMC Complementary and Alternative Medicine*, 6, 1–6.
- Sánchez ,R.F.; O. Jáuregui; I. Casals; C. Andrés Lacueva; M. Izquierdo Pulido & R.M. Lamuela-Raventós (2003). Liquid chromatographic/electrospray ionization tandem mass spectrometric study of the phenolic composition of cocoa (*Theobroma cacao*). *Journal of Mass Spectrometry*, 38, 35–42.
- Sari, A.B.T.; D.C. Mufida; N.N. Fatima; T.I. Hendrayati; T. Wahyudi & Misnawi (2014). In vitro antibacterial activity of cocoa ethanolic extract against *Escherichia coli*. *International Food Research Journal*, 21, 935–940.
- Sari, A.B.T.; Mahriani; G.A.P. Tiningrum; T. Wahyudi & Misnawi (2015). Cocoa extract indicated has activity on selectively killing breast cancer cells. *The Journal of Tropical Life Science*, 5, 128–132.
- Sari, A.B.T.; T. Wahyudi; D.C. Mufida; M. Tamarizki; D. Perwitahtati; I.K. Syamsuddin & Misnawi (2016). The use of cocoa ethanolic extract for treatment of Staphylococcal infection in rabbit skin model. *Pelita Perkebunan*, 32, 34–42.
- Shiferaw, B.; S. Solghan; A. Palmer; K. Joyce; E.J. Barzilay; A. Krueger & P. Cieslak (2012). Antimicrobial susceptibility patterns of shigella isolates in foodborne diseases active surveillance network (foodnet) sites, 2000–2010. *Clinical Infectious Diseases*, 54, 458–463.
- Vázquez-Agell, M.; M. Urpi-Sarda; E. Sacanella; S. Camino-López; G. Chiva-Blanch; V. Llorente-Cortés; E. Tobias; E. Roura; C. Andres-Lacueva & R.Lamuela-Raventós (2011). Cocoa consumption reduces NF-κB activation in peripheral blood mononuclear cells in humans. *Nutrition, Metabolism and Cardiovascular Diseases*, 23, 257–263.
- Walker, C.L.F.; J. Perin; M.J. Aryee; C. Bosch-Pinto & R.E. Black (2012). Diarrhea incidence in low- and middle-income countries in 1990 and 2010: a systematic review. *BMC Public Health*, 12, 1–7.
- WHO (2005). *Guidelines for Control of Shigellosis, Including Epidemics Due to Shigella dysenteriae Type 1*. World Health Organization Press, Geneva, Switzerland.

0