A petri dish containing a red agar medium. The top half of the dish shows several small, circular, white bacterial colonies. The bottom half of the dish is covered by a large, dense, white bacterial growth that has spread across the surface, showing a textured, almost crystalline appearance.

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 21 - Number 7 - July 2020

Front cover: *Bacillus cereus*
(PHOTO: JOE RUBIN)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033788



9 772085 472782

BIODIVERSITAS

Journal of Biological Diversity
Volume 21 - Number 7 - July 2020

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaie** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohib Shah** (Pakistan), **Mohamed M.M. Najim** (Srilanka), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Seyed Aliakbar Hedayati** (Iran), **Seyed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as *.doc* or *.docx* files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

A charge Starting on January 1, 2019, publishing costs waiver is granted to authors of graduate students from **Least Developed Countries**, who first publish the manuscript in this journal. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (*.doc* or *.rtf*; not *.docx*). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m⁻², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "*cit*" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

Detection of virulence and specific genes of *Salmonella* sp. indigenous from Jember, Indonesia

RIA YULIAN^{1,2}, ERLIA NARULITA^{1,2,*}, MOCHAMMAD IQBAL¹, DESY ROCHMIYAH SARI^{1,2},
IRMA SURYANINGSIH^{1,2}, DIAN EKA APRILIA FITRIA NINGRUM³

¹Program of Biology Education, Faculty of Teacher Training and Education, Universitas Jember. Jl. Kalimantan X, Jember 68121, East Java, Indonesia.
Tel.: +62-331-330738, *email: erlia.fkip@unej.ac.id

²Center for Development of Advanced Sciences and Technology, Universitas Jember. Jl. Kalimantan X, Jember 68121, East Java, Indonesia

³Department of Islamic Primary School Teacher Education, Faculty of Tarbiyah and Teacher Training, Universitas Islam Negeri Maulana Malik Ibrahim Malang. Jl. Tarbiyah, Malang 65144, East Java, Indonesia

Manuscript received: 13 March 2020. Revision accepted: 3 June 2020.

Abstract. Yulian R, Narulita E, Iqbal M, Rochmiyah D, Suryaningsih I, Ningrum DEAF. 2020. Detection of virulence and specific genes of *Salmonella* sp. indigenous from Jember, Indonesia. *Biodiversitas* 21: 2889-2892. The bacterium *Salmonella* sp. is the most common cause of foodborne infection in humans. It has the ability to invade the host, Salmonellosis is highly influenced by the combination of chromosome and plasmid in which *Salmonella* Pathogenesis Islands (SPIs) are the biggest gene in the chromosome that are responsible to build specific interaction between *Salmonella* and the host. This research performed detection on virulence and specific genes in the SPIs area using PCR method for two *Salmonella* spp. bacterial isolates originated from Jember which were KP2 and P21D. Seven pairs of primers including *stn*, *fimA*, *spvR* (virulence genes), *invA*, *ivaB*, *spvC*, and *fliC-d* (specific genes) were used. The result revealed that P21D bacteria containing 3 virulence genes i.e. *stn*, *fimA*, and *spvR*. Meanwhile, only two virulent genes *stn*, and *spvR* were present in KP2 bacteria. For the detection of specific genes, three genes *invA*, *ivaB*, and *spvC* were present while *fliC-d* genes were not detected. Only two, *invA*, and *ivaB* genes were detected in P21D bacteria, while *spvC*, and *fliC-d* were not detected.

Keywords: *Salmonella*, SPIs, virulence gene, specific gene

INTRODUCTION

All genes in the *Salmonella* chromosome are in the area that is called *Salmonella Pathogenesis Island* (SPIs) (Chaudhary et al. 2015; Andesfha, et al. 2019). SPIs in the *Salmonella* chromosome is responsible for building interaction between the *Salmonella* and the host (Sabbagh et al. 2010). The genes in the SPIs area can be treated as molecular marks to identify *Salmonella* and also the virulence type of those bacteria (Kumar et al. 2006). Specific genes that could be used to identify *Salmonella* bacteria had been widely reported. One of them was the detection of *S.typhi* using PCR with primer pairs such as *invA*, *ivaB*, *fliC-d*, *spv* genes (Kumar et al. 2006). *viaB*, *prt*, and *fliC-d* genes, each of them, was used to encode in synthesizing Vi (Capsule), O (LPS), and H antigen (flagellar) (Shanmugasamy et al. 2011). The virulence genes, i.e. *stn*, *fimA*, and *spvR* had been widely reported on many research especially for detection virulence genes in bacteria caused of foodborne disease (Naravaneni and Jamil 2005), for classification of *Salmonella* (Alphons and Jaap, 2005; Card et al. 2016), as well as for knowing nontyphoidal *Salmonella* in gastroenteritis (Araque 2009; Aoki et al. 2017).

The amplification of *invA* genes has been internationally acknowledged for the detection of *Salmonella* genus. *invA* gene is the primary gene in the operon *invABC* which encodes the protein in the inner membrane of *Salmonella* that is responsible for *invA*ding

intestinal epithelial cells of the host (Phumkhachorn and Rattanachaiunsopon 2017). *fliC-D* gene-encoded flagellin protein genes in phase 1 (Shanmugasamy et al. 2011). Genes *ivaB* and *fliC-d* encodes the synthesis of Vi (capsule), O (LPS), and antigen H (flagellar). Each antigen becomes the basis of detection and classification for *Salmonella* by Kauffmann-White scheme (Kumar et al. 2006; Mthembu 2019; Figueira and David 2012). *spv* gene holds the role in systemic infection by infecting host in the macrophage. *spv* gene allows to increase the virulence of *S. enteritidis* in the internal organs because one main function of *spv* operon is to potentiate the systemic spread of pathogen (Chaudhary et al. 2015). This research used *spvC* which was virulence-related gene in plasmid that is needed to survive in the host cell (Tekale et al. 2015).

Regarding the existence of virulence and specific genes of *Salmonella*, this research used two Jember *Salmonella* indigenous bacteria namely P21D and KP2 which were isolated from Puger and Kencong. There were 255 cases of food poisoning recorded in 2016 in Kencong and Puger (Diah 2016). Even though the status was decreasing, the bacterial contamination in Kencong area and surrounding was still detected in the early of March 2017 from the sampling conducted by Jember Health Department in which, in the sampling process, *Salmonella* sp. bacteria were still found. The recent case of food poisoning in Jember was in early of January 2020 (Health Crisis Center). Thus, this study aimed to detect virulence and specific gene contained in Jember *Salmonella* indigenous bacteria.

MATERIALS AND METHODS

Procedures

Salmonella sp. culture

Two *Salmonella* isolates (KP2, and P21D) were obtained from fish and vegetable wastes from Puger and Kencong, respectively. Reactivation was done by inoculating the bacteria from SSA medium to *Luria-Bertani* medium (LB). The bacteria were cultured for 24 hours in a shaker.

Isolation and extraction DNA of Salmonella sp.

The bacteria were grown in LB medium for 24 hours which then 1000µl aliquot was retrieved and centrifuged at 4°C, 1000rpm for 10 minutes. The obtained supernatant contained DNA transferred to other Eppendorf tubes and stored at 4°C. DNA extraction by boiling was done with a modification. The collected material was placed into a tube containing 40 µL, then boiling for 10 minutes. The mixture was centrifuged at 12000 rpm for 10 minutes. The supernatant was collected and 70µL volumes of cold absolute ethanol were added, then centrifuged for 20 minutes. The pellet was washed in EtOH, dried the genome, and resuspended in TE buffer (Dashti et al. 2009).

DNA amplification

The first PCR amplification process was conducted using gradient temperature to obtain optimum temperature for each primer pairs (Table 1). The gradient temperature setting was based on melting temperature (T_m) of each primer at 5°C below of T_m with 35 cycles. The PCR condition was denaturated at 95°C for 2 minutes, followed by 35 cycles, denaturation at 95°C for 30 seconds, annealing temperature at 46.9°-50.8°C for 30 seconds, and extension at 72°C for 45 seconds, and final extension at 72°C for 7 minutes (Sunar et al. 2014). The amplification products were separated by electrophoresis in 1% agarose gel stained with 5µg-/ml ethidium bromide using 1kb DNA ladders as molecular-weight size marker.

RESULTS AND DISCUSSION

The research step initiated by conducting genomic DNA extraction using colony and boiling PCR method which was then checked using electrophoresis in 1% agarose gel. The genomic visualization results of two isolates were seen for >1kbp band (Figure 1). Next, in PCR process, the amplification result of both *Salmonella sp.* samples which were KP2 and P21D showed difference in annealing temperature for each primer. The *stn* gene was amplified at annealing temperature of 46,9°C. Meanwhile, both *spvR* gene, and *fimA* gene were amplified at the annealing temperature of 50,8°C. For specific genes, the optimum annealing temperatures were *invA* gene (50,8°C), *ivaB* gene (47°C), *spvC* gene (57°C), while *fliC-d* gene did not show any amplified band at the optimum temperature gradient between 47°C - 54°C. The amplification result of *stn* gene showed there were 260 bp DNA bands on both samples (Figure 2). *Salmonella enterotoxin (stn)* is a gene encoding *stn* protein that causes gastroenteritis leading to nausea, vomiting, abdominal cramps, fever, and diarrhea. *stn* gene is detected in all *Salmonella* strains from broilers (Fekry et al. 2018). The virulence genes detected in this study were the *stn* gene, the *fimA* gene, and the *spvR* gene. The three genes in the *Salmonella sp.* are pathogenic. Virulence genes encode products that assist organisms in expressing virulence in host cells (Muthu et al. 2014). According to Thomas and Wigneshweraraj (2014) products of genes that facilitate successful colonization and bacterial survival or cause damage to the host are considered as determinants of virulence. The presence of virulence genes in *Salmonella* is related to the causes of salmonellosis and foodborne diseases in humans. *Salmonella* virulence is a combination of chromosome and plasmid factors (Chaudhary et al. 2015).

Table 1. Primers for detection of virulence and specific genes of *Salmonella sp.*

Gene target	Primer sequence (5-3')	Target size (bp)	Reference
<i>fimA</i>	F: CCT TTC TCC ATC GTC CTG AA R: TGG TGT TAT CTG CCT GAC CA	85	Chaudhary et al. (2015)
<i>Stn</i>	F: CTT TGG TCG TAA AAT AAG GCG R: TGC CCA AAG CAG AGA GAT TC	260	Chaudhary et al. (2015)
<i>spvR</i>	F: CAG GTT CCT TCA GTA TCG CA R: TTT GGC CGG AAA TGG TCA GT	310	Chaudhary et al. (2015)
<i>invA</i>	F: GTG AAA TTA TCG CCA CGT TCG GGC AA R: TCA TCG CAC CGT CAA AGG AAC C	284	Chaudhary et al. (2015)
<i>ivaB</i>	F: GTT ATT TCA GCA TAA GGA G R: ACT TGT CCG TGT TTT ACT C	599	Phumkhachorn and Rattanachaikunsopon (2017)
<i>fliC-d</i>	F: ACT CAG GCT TCC CGT AAC GC R: GGC TAG TAT TGT CCT TAT CG G	763	Phumkhachorn and Rattanachaikunsopon (2017)
<i>spvC</i>	F: ACT CCT TGC ACA ACC AAA TGC GGA R: TGT CTT CTG CAT TTC GCC ACC ATC A	571	Chaudhary et al. (2015)




Figure 1. Electroforegram of *fliC* amplification showed no band at all. M: marker; line 1-8: *Salmonella* sp. P21D; and line 9-16: *Salmonella* sp. KP2




Figure 3. *fimA* gene with size of 85 bp. Line M: marker; line 1: *Salmonella* sp. P21D; and line 2: *Salmonella* sp. KP2




Figure 2. *stn* gene with size of 339 bp. Line M: marker; line 1: *Salmonella* sp. P21D; and line 2: *Salmonella* sp. KP2




Figure 4. *spvR* gene with size of 310 bp. Line M: marker; line 1: *Salmonella* sp. P21D; and line 2: *Salmonella* sp. KP2

Salmonella sp. is a member of Enterobacteriaceae, Gram-negative, rod-shaped, non-spherical bacteria, and generally moves using peritrich flagellum (Araque 2009). *Salmonella* bacteria have facultative anaerobic properties to grow at a temperature range of 5-45°C with the optimum at 35-37°C. Other characteristics of *Salmonella* are that they multiply by dividing, easy to grow on simple mediums, and resistant to certain chemicals like. The DNA band of *fimA* gene only appeared on P21D while KP2 produced no bands (Figure 3).

The appearance of DNA band on the P21D indicated the presence of *fimA* gene while bacteria KP2 have no *fimA* gene. *fimA* is a fimbriae encoding gene in *Salmonella* (Zeiner et al. 2019). Fimbriae are found a lot on the enteric bacteria surface and mediating the attachment to eukaryotic cells. Type-1 fimbriae protein of *Salmonella typhi* was encoded by *fimA* gene group, with *fimA* assigned as single transcription unit. Based on amplification result of sample KP2, these bacteria have no type 1 fimbriae protein which contain *fimA* gene. Sample KP2 might have another fimbriae gene from *fimbrial gene cluster*. *FimA* is a fimbria coding gene in *Salmonella*. Fimbria is found in many

surfaces of enteric bacteria and functions for attachment to eukaryotic cells.

Bacteria can be classified based on phylogeny (to see the relationship between *Salmonella* bacteria). A phylogenetic tree can be made by looking at the genome sequence of the bacteria using 16S rRNA analysis. Around 2,463 *Salmonella* serotypes were placed under 2 species due to differences in the genome sequence of the 16S rRNA analysis results, including *Salmonella enterica* (2443 serotypes) and *Salmonella bongori* (20 serotypes).

spvR gene could be amplified on the two samples (Figure 4). The size of amplified DNA was 310 bp. The appearance of single band on the amplification result revealed that both samples possessed gen *spvR* with thick band displaying a high DNA concentration. *spv* gene (*Salmonella plasmid virulence*) is a 7.8 kb locus consisting of 5 genes, *spvR*, located in a plasmid which is generally associated with some serotypes. *spv* gene possesses the ability to increase the severity of enteritis, infection, and persistence at extra-intestinal sites (Oliveira et al. 2003; Alphons and Jaap 2005; Card et al. 2016). The *spv* gene has the ability to increase the severity of enteritis and

infection and persistence at the extraintestinal location (Oliveira et al. 2003).

invA gene was amplified on the two *Salmonella* samples revealed a 244 bp band (Figure 5A). On the P21D bacteria sample, the band was slightly thinner than the band on KP2. The thickness of the bands may be due to the concentration of DNA in the bacteria. *invA* gene is the specific primer that owns solely by genus *Salmonella* which means that the two bacterial samples used in this research belong to *Salmonella* genus (Phumkhachorn and Rattanachaiakunsopon 2017). *invA* gene makes the epithelial cell *invA* sion process, so when *salmonellosis* occurs, this gene will appear. Forty isolates of *Salmonella* had been studied for detection of *invA* gene using PCR, and *invA* appeared in all of the isolates (Sharma and Das 2016). The amplification result of *ivaB* primer gene on the two samples could be seen from the 599 bp band (Phumkhachorn and Rattanachaiakunsopon 2017). Sample P21D has a thinner band than sample KP2 (Figure 5B). *ivaB* gene is especially owned by *Salmonella typhi* dan *Salmonella paratyphi*. The function of this gene is to operate in *viaB* area where locus on chromosome contains specific structural genes to express antigen Vi (Virulence) (Xiong et al. 2017).

From the result of the amplification, *spvC* primer gene only appeared in sample 2 (KP2) at 571 bp (Figure 6), while sample 1 (P21D) did not contain *spvC* gene. Gene *spv* (*Salmonella plasmid virulence*) is a 7.8 kb locus consisting of 5 genes namely *spvRAB* (Alphons and Jaap 2005; Card et al. 2016). P21D bacteria did not contain *spvC* gene, so they do not possess the ability in partial virulence. However, it is possible for P21D sample to have the other 5 *spv* genes. According to the research, *spv* primer gene, *spvR*, and *psvC*, on 102 *Salmonella enteritidis* samples from poultry, human feces and foodborne disease plague were successfully amplified (Oliveira et al. 2003). The outcome was detection of 91.2% *spvR* and 90.2% *spvC*

virulence genes from all *Salmonella enteritidis* samples. *spvR* gene has a role to regulate LysR protein to express other *spv* genes (Alphons and Jaap 2005; Card et al. 2016). The absence of *spvR* gene might result in avirulence while the absence of *spvC* gene could show partial virulence. *Salmonella Enterica Serovar Typhimurium* strain, found on the rat, was inoculated subcutaneously, and contained *spvB* gene (Matsui et al. 2001). *spvB* gene expressed ADP-ribosylation enzyme which inhibited the process of polymerization actin in host cell, so it could inhibit the fusion between phagocyte and lysosome (Chu et al. 2016). *spvD* gene reduced the pro-inflammation regulation by inhibiting promotor activation which was controlled by NF- κ B (Silva et al. 2017; Liu et al. 2017). The *spvR* gene functions to regulate the LysR essential protein to express other *spv* genes. Without the *spvR* gene, it may be avirulent while the absence of the *spvC* gene can show partial virulence.




Figure 6. Electroforegram of *spvC* gene amplification of 571 bp. Line M: marker; line 1-8: *Salmonella* sp. P21D; and line 9-16: *Salmonella* sp. KP2




Figure 5. Electroforegram of *invA* (A) volume 25 μ L and *ivaB* (B) volume 3 μ L genes with size of 599 and 244 bp, respectively. Line M: marker; line 1: *Salmonella* sp. P21D; and line 2: *Salmonella* sp. KP2

In summarize, the detection result of virulence and specific genes on two *Salmonella* sp. bacterial isolates from Jember showed the presence of virulence genes namely *stn*, *fimA*, and *spvR* in P21D isolate. Meanwhile, the specific genes were *invA*, dan *ivaB*. In the KP2 isolate, virulence genes were *stn*, dan *spvR* while the detected specific genes were *invA*, *ivaB*, dan *spvC*. Many virulence factors have been shown to play various roles in the pathogenesis of *Salmonella* infections. These factors include flagella, capsules, plasmids, adhesion systems, and type 3 secretion systems (T3SS) encoded on the pathogenic islands of *Salmonella* (SPI)-1 and SPI-2 and other SPI.

ACKNOWLEDGEMENTS

This work was supported by the Research and Community Service Institute (LP2M), University of Jember, Indonesia through Hibah Mendukung IDB contract number of 3235/UN25.3.1/LT/2019.

REFERENCES

- Alphons JAMV, Jaap EVD. 2005. Distribution of “classic” virulence factors among *Salmonella* spp. FEMS. Immunol Med Microbiol 44: 251-259.
- Andesfha E, Agustin I, Mayasari NLPI, Rahayuningtyas I, Istiyaningsih J. 2019. Detection of *Salmonella* pathogenicity island and *Salmonella* plasmid virulence genes in *Salmonella enteritidis* originated from layer and broiler farms in Java Island. J Adv Vet Anim Res 3 (6): 2311-7710.
- Aoki Y, Katsuhiko K, Hinorobu K, Masayoshi S, Yukie A, Riu H, Yudai W, Akihito H. 2017. Clinical features of children with nontyphoidal *Salmonella* bacteremia: A single institution survey in rural Japan. PLoS One 12 (6): e0176990. DOI: 10.1371/journal.pone.0176990.
- Araque M. 2009. Nontyphoidal *Salmonella* Gastroenteritis in Pediatric Patients from urban Area in the city of Merida, Venezuela. J Infect Dev Control 3 (1): 28-34.
- Card R, Kelly V, Mary B, John S, William C, Tony S, Rob D, Muna FA. 2016. Virulence Characterisation of *Salmonella enterica* Isolates of Differing Antimicrobial Resistance Recovered from UK Livestock and Imported Meat Samples. Front Microbiol 2: 1-11.
- Chaudhary JH, Nayak JB, Brahmabhatt MN, Makwana PP. 2015. Virulence Genes Detection of *Salmonella* Serovars Isolated from Pork and Slaughterhouse Environment in Ahmedabad, Gujarat. Vet World 8 (1): 121-124.
- Chu Y, Gao S, Wang T, Yan J, Xu G, Li Y, Niu H, Huang R, Wu S. 2016. A novel contribution of *spvB* to pathogenesis of *Salmonella typhimurium* by inhibiting autophagy in host cells. Oncotarget 7 (7): 8295-8309.
- Dashti A, Jadaon MM, Abdulsamad AM, Dhasti. 2009. Heat Treatment of Bacteria: A Simple Method of DNA Extraction for Molecular Techniques. Kuwait Med J 41 (2): 117-122.
- Fekry E, Abdeen E, Ammar AM, Hussien AE. 2018. Molecular Detection of *InvA*, *OmpA* and *Stn* Genes in *Salmonella* Serovars from Broilers in Egypt. Alex J Vet Sci 56 (1): 69-74.
- Figueira R, David WH. 2012. Functions of the *Salmonella* Pathogenicity Island 2 (SPI-2) Type III Secretion System Effectors. Microbiol 128: 1147-1161.
- Health Crisis Center. 2016. Kejadian Luar Biasa (KLB)-Keracunan di Jember, Jawa Timur. <http://pusatkrisis.kemkes.go.id/kejadian-luar-biasa-klb--keracunan--di-jember-jawa-timur-04-09-2016> [Indonesian]
- Health Crisis Center. 2020. Kejadian Luar Biasa (KLB)-Keracunan di Jember, Jawa Timur. [http://www.pusatkrisis.kemkes.go.id/Kejadian-Luar-Biasa-\(KLB\)---Keracunan--di-JEMBER-JAWA-TIMUR-02-01-2020-42](http://www.pusatkrisis.kemkes.go.id/Kejadian-Luar-Biasa-(KLB)---Keracunan--di-JEMBER-JAWA-TIMUR-02-01-2020-42). [Indonesian]
- Kumar S, Balakrishna K, Batra H. 2006. Detection of *Salmonella enterica* serovar *typhi* (*S. typhi*) by selective amplification of *invA*, *viaB*, *fliC-d* and *prt* genes by Polymerase Chain Reaction in multiplex format. Lett Appl Microbiol 42 (2): 149-154.
- Liu T, Lingyun Z, Donghyun J, Shao-Cong, S. 2017. NG-kB signaling in inflammation. Sig Transduct Target Ther 2: 17023.
- Matsui H, Bacot CM, Garlington WA, Doyle TJ, Roberts S, Gulig PA. 2001. Virulence plasmid-borne *spvB* and *spvC* Genes can replace the 90 kilobase plasmid in conferring virulence to *Salmonella enterica* serovar *typhimurium* in subcutaneously inoculated mice. J Bacteriol 183 (15): 4652-8. DOI: 10.1128/JB.183.15.4652-4658.2001.
- Mthembu TP, Oliver TZ, Mohamed EEZ. 2019. Detection and molecular identification of *Salmonella* virulence genes in livestock production systems in South Africa. Pathogens 1 (24): 1-17.
- Muthu G, Suresh A, VishnuPrabu D, Munirajan AK, Mary SE, Sathishkumar E, Gopinath P, Srivani S. 2014. Detection of virulence genes from *Salmonella* species in Chennai, India. CIBTech J Microbiol 3 (1): 11-14.
- Oliveira SD, Rodenbusch CR, Michae GB, Cardoso MI, Canal CW, Brandelli A. 2003. Detection of virulence genes in *Salmonella enteritidis* isolated from different sources. Braz J Microbiol 34 (1): 123-124.
- Phumkhaichorn P, Rattanachaikunsopon P. 2017. Detection of Viable *Salmonella typhi* using Three Primer Pairs Specific to *invA*, *ivaB*, and *fliC-d* Genes. Emir J Food Agr 29 (4): 312-316.
- Sabbagh SC, Forest CG, Lapage C, Leclerc JM, Daigle F. 2010. Uncovering distinctive features in the genomes of *Salmonella enterica* serovars *typhimurium* and *typhi*. FEMS Microbiol Lett 305 (1): 1-13.
- Shanmugasamy M, Velayutham T, Rajeswar J. 2011. *InvA* gene-specific PCR for detection of *Salmonella* from broilers. Vet World 4 (12):562-564.
- Sharma I, Das K. 2016. Detection of *invA* gene in isolated *Salmonella* from marketed poultry meat by PCR assay. J Food Process Technol 2 (1):1-4.
- Silva C, Puente JL, Calva E. 2017. *Salmonella* virulence plasmid pathogenesis and ecology. Pathog Dis 75: 6. DOI: 10.1093/femspd/ftx070.
- Sunar NM, Stentiford, Stewart DI, Fletcher LA. 2014. Molecular techniques to characterize the *invA* Genes of *Salmonella* spp. for pathogen inactivation study in composting. Orbit 13 (1): 43-46.
- Tekale AA, Savalia CV, Kshirsagar DP, Brahmabhatt MN, Chatur YA. 2015. Detection and virulence gene characterization of *Salmonella* isolates from fish by conventional and molecular methods. J Vet Pub Hlth 13 (1): 43-46.
- Xiong K, Chunyue Z, Zhijin C, Chunping Z, Yong T, Xiancai R, Yanguang C. 2017. Vi capsular polysaccharide produced by recombinant *Salmonella enterica* serovar *paratyphi* a confers immunoprotection against infection by *Salmonella enterica* serovar *typhi*. Front Cell Infect Microbiol 7: 135. DOI: 10.3389/fcimb.2017.00135.
- Zeiner SA, Dwyer BE, Clegg S. 2019. *FimA*, *FimF*, and *FimH* are necessary for assembly of type 1 fimbriae on serovar *typhimurium*. J Infect Immun 80 (9): 3289-3296.