

**VOLUME 4 ISSUE 1 : JAN - FEB 2023**

Digital Repository Universitas Jember



# **INTERNATIONAL JOURNAL OF SCIENTIFIC ADVANCES**

**ISSN : 2708 - 7972**

[WWW.IJSCIA.COM](http://WWW.IJSCIA.COM)

# EDITORIAL BOARD

## Editor-in-Chief

Dr. T. J. Ashaolu

Email: editor@ijscia.com, ijsciaeditor@gmail.com

## Assistant Editor-in-Chief

Joseph O. Ashaolu

National Yang-Ming University, Taipei, Taiwan

## EDITORIAL MEMBERS

- Prof. Emeritus MNO Sadiku**, Prairie View A&M University, Texas, USA  
**Professor Ramesh C. Gupta**, Former Vice Chancellor, Radha Govind University, Jharkhand, India  
**Professor Massoud Kaykhali**, University of Sistan & Baluchestan, Zahedan, Iran  
**Professor Elżbieta Macioszek**, Silesian University of Technology, Gliwice, Poland  
**Professor Dariusz Jacek Jakobczak**, Koszalin University of Technology, Koszalin, Poland  
**Asst. Prof. Esam AL Lawati**, A' Sharqiyah University, Ibra, Sultanate of Oman  
**Dr. S. A. O. Adeyeye**, Ton Duc Thang University, Ho Chi Minh City, Vietnam  
**Asst. Prof. Ibrahim Khalifa**, Benha University, Moshtohor, Egypt  
**Professor Dr. Ho Soon Min**, INTI International University, Nilai, Malaysia  
**Dr. B. J. Alegbeleye**, St Elizabeth Catholic General Hospital, Kumbo-Nso, Cameroon  
**Dr. Ridwan Babatunde Ibrahim**, Academia Sinica, Taipei, Taiwan  
**Prof. Dr. Eng. Ahmed kadhim Hussein**, Babylon University, Babylon City, Hiila, Iraq  
**Dr. Murat Basar**, IVF Laboratories Bahçeci Sağlık Grubu, Istanbul, Turkey  
**Dr. Bassam Saeed**, Farah Association for Child with Kidney Disease, Damascus, Syria  
**Asst. Prof. Selezneva Irina Stanislavovna**, Ural Federal University, Yekaterinburg, Russia  
**Professor Michael Gr. Voskoglou**, GTES Institute of Western Greece, Patras, Greece  
**Assoc. Prof. Thoudam Paraskumar Singh**, MGM College of Engineering and Technology Kamothe, Navi Mumbai, India  
**Dr. Isaac Delali Kottoh**, Ghana Atomic Energy Commission, Accra, Ghana  
**Assoc. Prof. Godfred A Menezes**, RAK Medical & Health Sciences University, Ras Al Khaimah, United Arab Emirate  
**Assoc. Prof. Syed Kamran Sami**, Balochistan University of Information Technology, Quetta, Pakistan  
**Asst. Prof. Muhammad Amin**, Balochistan University of Information Technology, Quetta, Pakistan  
**Ma. Gladys B. Aquino**, De La Salle Medical and Health Sciences Institute, Dasmariñas, Philippines  
**Assoc. Prof. Vahe Davtyan**, Brusov State University, Yerevan, Armenia  
**Dr. Sylvain Somé Marie**, National Yang-Ming University, Taipei, Taiwan  
**Professor Teresa Orbera**, University of Oriente, Santiago de Cuba, Cuba  
**Dr. Magdaline Joseph Kwaji**, Bioresources Development Center, Odi, Bayelsa, Nigeria  
**Dr. Sampson Twumasi-Ankrah**, Kwame Nkrumah University of Science and Technology Kumasi, Ghana  
**Asst. Prof. Nisarg G. Gandhewar**, SB Jain Institute of Technology, Nagpur, India  
**Dr. Manoj Khandelwal**, Federation University, Ballarat, Victoria, Australia  
**Dr. Taimur Sharif**, Newman University, Birmingham, UK  
**Dr. Tinni Dutta**, University of Calcutta, Kolkata, India  
**Professor Dr. Jacob Oluwoye**, Alabama A&M University, Normal, Alabama, USA  
**Dr. Benjamin D. Sookhoo**, University of South Florida, Tampa, Florida, USA,  
**Victor Eyo Assi**, University of Uyo, Uyo, Akwa Ibom, Nigeria  
**Dr. Bid D. Dhruvprasad**, Sarvajanic College of Physiotherapy, Gujarat, India  
**Dr. Ankit Bhargava**, Consultant Physio & Fitness Expert-Wendt India Ltd. India  
**Dr. Manoranjan Tripathy, Dev Sanskriti Vishwavidyalaya**, Uttarakhand, India  
**Asst. Prof. Abbas Ar. Mohamed**, PMBAH National Guard Health Affairs, Al Madinah Al Munawwarah, Kingdom of Saudi Arabia

# Digital Repository Universitas Jember

- Asst. Prof. Nurul Mohammad Zayed**, Daffodil International University, Dhaka, Bangladesh  
**Prof. Dr. Hasibun Naher**, Brac University, Dhaka, Bangladesh  
**Davis Ojima J.P. Bursar**, Ignatius Ajuru University of Education, Port Harcourt, Rivers, Nigeria  
**Dr. Okogwu Antonia**, University of Port Harcourt, Port Harcourt, River, Nigeria  
**Dr. Garyfalia Charitaki**, Hellenic Open University ( University of Thessaly), Volos, Greece  
**Dr. Munish Kumar Sharma**, University of Alberta, Edmonton, Canada  
**Dr. Niyonzima N. Francois**, Deputy Vice Chancellor Academic and Research INES Ruhengeri, Rwanda  
**Dr. Srishti Sharma, M. Patel** College of Physiotherapy, Gandhi Nagar, India  
**Professor R. K. Mathukia**, Junagadh Agricultural University, Junagadh, Gujarat, India  
**Professor Hamid Ali Abed Al-Asadi**, Basra University, Basra, Iraq  
**Dr. Firas Abdullah Thweny Al-Saedi**, Nahrain University, Baghdad, Iraq and Private Researcher, New Jersey, USA  
**Dr. George Fuh Kum**, University of Yaoundé I, Yaoundé, Cameroon  
**Professor Mohamed El Houseny El Sebeay Shams**, Mansoura University, Mansoura, Egypt  
**Dr. Peter Ashlame Agu**, Nasarawa State University, Keffi, Nigeria  
**Dr. Asadullah Baloch**, Mir Chakar Khan Rind University, Quetta, Pakistan  
**Professor Ignatius Isaac Dambudzo**, Zimbabwe Open University, Harare, Zimbabwe  
**Professor Hamid Ali Abed Alasadi**, Iraq University College, Basra, Iraq  
**Dr. Ammar Daher Bashatweh**, Independent Scholar, Irbid, Jordan  
**Subrat Kumar Mahapatra**, Central University, Santiniketan, West Bengal  
**Dr. Xiong Chen**, Hubei University of Medicine, Shiyan City, China





# IJSCIA

## INTERNATIONAL JOURNAL OF SCIENTIFIC ADVANCES

### Indexing and Abstracting



ResearchGate

refseek\*



HARVARD  
UNIVERSITY



## TABLE OF CONTENTS

Risk Factors of Low-Birth-Weight Baby in Indonesia

Characteristics of Plate and Screw Fixation in Tibial Shaft Fracture at Sanjiwani General Hospital Gianyar in 2021: A Retrospective Descriptive Study

Sex Education: A Review of Sexual Behaviors in Sub-Saharan African Adolescent

Smoking as A Risk Factor of Low Superoxide Dismutase (SOD) Levels and Early Aging in Women Aged 20 to 35 Years

High Serum Level of Matrix Metalloproteinase-1 (MMP-1) Correlates Positively with The Severity of Photoaging Facial Wrinkles

Characteristic Cases of Snake Bite in Sanjiwani Hospital

Potential of Renewable Energy Sources for Distributed Generations: An Overview

Internal and External Behavior Outcomes in Children with Congenital Adrenal Hyperplasia: A Systematic Review

Buerger's Disease (Thromboangiitis Obliterans) among Smokers: A Literature Review

The Correlation Between Frequency of Transfusion and Alloimmunization in Transfusion-Dependent Thalassemia Patients

Management of Heart Failure Complications in Sepsis Patients

The Effect of Mental Posyandu Community Health Worker Training on Knowledge and Attitudes about Handling Patients with Mental Disorders in Indonesia

Electrical Injuries: Cardiovascular Implication and Management

A Literature Review of Clinical Characteristics of Cerebral Malaria

Knowledge, Attitude and Perception of Assisted Reproductive Technology Among Patients Attending Health Clinics at Babcoclishan-Remo Ogun Statek University Teaching Hospital I

Statistical Analysis of 1890 Land-Grant Universities Future Undergraduate Fall Enrollment Using Time-series from 1996 to 2018

The Effect of Alpha Lipoic Acid (ALA) Supplementation on the in Vitro Maturation Media of Goat Oocyte on Tumor Alpha Necrosis Factor (TNF- $\alpha$ ) Expression and Malondialdehyde (MDA) Levels

Comparison of Total Number of Bacteria in Raw Milk Friesian Holstein Cow Based on Milking Techniques Using Total Plate Count (TPC) Test

Rumen Fermentation Optimization of Kacang Goats Fed Complete Silage-Based Feed Sorghum-Clitoria ternatea with Various Concentrate Levels Contains ZnSO<sub>4</sub> and Zn-Cu Isoleucinate

Relevancy Between Knowledge and Attitudes of Students to Human Papillomavirus (HPV) Vaccine: Literature Review

Phenotype of the Production Characteristics of Broiler Chickens Fed Feed Containing Fermented Cow Feces

The Correlation Between Gestational Weight Gain with Children Nutritional Status Age 0 – 2 Years in Simomulyo Health Center Sukomanunggal Village Surabaya City

Learning Disorder in Attention Deficit Hyperactivity Disorder (ADHD) Children: A Literature Review

Screening Asymptomatic Men for Prostate Cancer Using Prostate-Specific Antigen as An Early Detection Tool: A Review of Existing Literature

Adverse Effects of Linezolid in Tuberculosis Patients at Dr. Soetomo General Hospital Surabaya

Glaucoma Medication and its Relationship with Depression

# Certificate of Publication

INTERNATIONAL JOURNAL OF SCIENTIFIC ADVANCES (IJSCIA)

ISSN: 2708-7972

Is hereby awarding this certificate to

**Zahrah Febianti**

In recognition of the publication of the paper entitled

**Buerger's Disease (Thromboangiitis Obliterans)  
among Smokers: A Literature Review**

Published in IJSCIA ([www.ijscia.com](http://www.ijscia.com))

Volume 4 Issue 1, in 6 February 2023



Website: [www.ijscia.com](http://www.ijscia.com)

Email: [editor.ijscia@gmail.com](mailto:editor.ijscia@gmail.com)



# Buerger's Disease (Thromboangiitis Obliterans) among Smokers: A Literature Review

Ichlasul Mahdi Fardhani<sup>1</sup>, Zahrah Febianti<sup>2\*</sup>, and Wahyu Agung Purnomo<sup>3</sup>

<sup>1</sup>Medical Program, Faculty of Medicine, University of Jember, East Java, Indonesia

<sup>2</sup>Department of Biochemistry, Faculty of Medicine, University of Jember, East Java, Indonesia

<sup>3</sup>Department of Pulmonologist and Respiratory Medicine, Dr. Soebandi Regional Hospital-Faculty of Medicine, University of Jember, East Java, Indonesia

\*Corresponding author details: Zahrah Febianti; [zfebianti.fk@unej.ac.id](mailto:zfebianti.fk@unej.ac.id)

## ABSTRACT

Thromboangiitis obliterans (TAO) also referred to as Buerger's disease is a segmental, non-atherosclerotic, and progressive inflammatory vascular disease. The most frequently impacted arteries are the small and medium arteries in the upper and lower extremities. Smoking plays a significant role in the emergence and progression of TAO, despite the fact that its etiology and pathophysiology are uncertain. Anamnesis, physical examination, and supporting examination can all be used to confirm the diagnosis of TAO. The effective method of TAO management is smoking cessation. Besides smoking cessation, there are supportive therapies both pharmacological and non-pharmacological that can be carried out to maintain the maximum blood flow and prevent complications such as amputation and secondary infections.

**Keywords:** Buerger's disease; thromboangiitis obliterans; smoking; chronic limb ischemia; rare disease

## INTRODUCTION

Thromboangiitis obliterans (TAO), often known as Buerger's disease, is a segmental, nonatherosclerotic, progressive inflammatory vascular disease that most frequently affects small and medium arteries in the upper and lower extremities (1,2). Inflammation that occurs in TAO due to peripheral vascular occlusion causes the patients to usually complain of claudication or pain at rest in the arms, hands, legs, and feet. The patients perceive it as pain arising from joint or neuromuscular disorders. The term for this condition is Raynaud's phenomenon (3,4).

The term "thromboangiitis obliterans" was first used in 1879 by Von Winiwarter, an Austrian-Belgian surgeon. Leo Buerger was able to describe TAO disease in 1908 based on the pathological findings of the amputated extremities of a TAO patient. Since then, there have been further developments about TAO disease (5,6). Although there are cases of thromboangiitis obliterans worldwide, they are more prevalent in Eastern Europe, South-East Asia, and South America (7). The prevalence of thromboangiitis obliterans varies worldwide. Western Europe has the lowest prevalence, ranging from 0.5% to 5.6%. Eastern Europe, South-East Asia, and South America are the regions where this disease is more prevalent (7). India has the highest prevalence, with 45 to 63% of the population (8). TAO affects males three times more frequently than females, with the ratio being about three to one. However, there is also an increasing trend of cases among women, which is associated with smoking trends (9).

TAO has a strong relationship with smoking despite the fact that the etiology is still uncertain. Thromboangiitis obliterans is mostly found in young to middle-aged males (21-45 years) with a smoking history.

Direct and indirect exposure to cigarette smoke is associated with the progression of disease (6). According to research by Fazeli et al. (2012), 108 patients who were diagnosed with TOA based on Shionoya criteria have an average smoking age of 21 years and smoked an average of one pack or around twenty cigarettes per day. Based on the multivariate analysis of the study, smoking duration was significantly associated with an adverse outcome, which is the amputation of the affected extremity (10).

Only a small percentage of smokers worldwide eventually become TAO. Due to limited publication and research on TAO and smoking behavior, the authors decided to write about TAO and its relation to smoking.

## CLINICAL FEATURE

Males with a history of smoking and who are younger than 50 years old are more likely to have TAO (11). The classic symptom of TAO is Raynaud's phenomenon, also known as livedo reticularis, in which the patient experiences pain in their hands, feet, and fingers at rest. These symptoms indicate the ischemic processes of the peripheral vascular. Initially, TAO affects the distal extremity blood vessels, but as the disease progresses, it can also affect the proximal vessels (9). According to research by Sasaki et al. (2000), 74.7% of cases were confined to the lower extremities, whereas 20.2% involved both extremities. The most frequently affected vessels of the lower extremities were anterior tibial artery (41.4%) and posterior tibial artery (40.4%), while the ulnar artery (11.5%) was the most frequently affected in the upper extremities (12).

## ETIOLOGY

The etiology of thromboangiitis obliterans, an autoimmune vasculitis disease that affects medium and large-sized vessels, remains unknown.

Cigarette smoke exposure is considered the initial step in the emergence and progression of diseases (13). Moreover, it is thought to lead to immunological dysfunction and hypersensitivity, which are associated with increased levels of type-1 and type-3 collagen, impaired endothelial vasodilatation, and increased anti-endothelial cell antibody (AECA) titers. In addition to exposure to cigarette smoke, there are several possible etiologies for the occurrence of this disease, including genetic factors due to an increased amount of human leukocyte antigens (HLA-54, HLA-A9, and HLA-B5), hypercoagulability, infection, and immunological mechanisms (14,15).

#### **PATHOPHYSIOLOGY OF SMOKING-INDUCED BUEYER'S DISEASE**

Although the pathophysiology of TAO is still unclear, it is believed that thrombosis, infection, and the presence of some autoantibodies may contribute to the occurrence of the disease. (16). According to Liew et al. (2015), there were four hypotheses about the pathogenesis of TAO, including variants of immunologic arteritis, atherosclerosis, hyperhomocysteinemia, and dental bacterial thrombosis (17). Based on histopathological findings, TAO was classified into three phases, including acute, subacute, and chronic. The occurrence of minimal inflammation in the vessel walls of the affected vessels is the main characteristic of the acute phase, polymorphonuclear leukocytes (PMN) predominate and microabscess formation within the thrombus. In the subacute phase, which is also called granulomatous inflammation, the PMN cells surround the microabscess and have the potential to organize and recanalize the thrombus. At a chronic phase, which is the end stage of the disease, the thrombus and vascular fibrosis occlude the blood vessels (9).

From several existing pathophysiology, smoking has been discovered to have a significant part in the pathophysiology and incidence of TAO (18). Increased levels of neutrophil elastase (NE) and reactive oxygen intermediates (ROIs) in smoking may trigger an inflammatory response that damages tissue (19). As a result of the neutrophil antimicrobial response, neutrophil elastase, which is a proteolytic enzyme degrades collagen and elastin fibers in the PMN primary granules (20). Furthermore, Iwai et al. (2005) found bacteria from the genus *Porphyromonas* in arterial thrombosis, which are gram-negative bacteria and originate from the mouth (21). This finding is in line with the postulate that smoking can lead to chronic gingivitis, which in turn stimulates the proliferation of *Porphyromonas gingivalis* in the cavity and may eventually lead to the bacteria being engulfed by platelets, resulting in the formation of an infective thrombus in which it crosses the bloodstream and causes thrombosis of blood vessels (18). These findings are also consistent with Allen and Brown's hypothesis, which stated that the etiology of TAO may be related to pathogenic microbes or viruses (22). These gram-negative bacteria can induce an immune response from Th1 and Th17 through the TLR4 receptor due to increased levels of cytokines interleukin-8 (IL-8) and NE. This series of events causes inflammatory cells to release vasoactive mediators, including histamine, bradykinin, and prostaglandins (19). An inflammatory thrombus occludes blood vessels as a result of neutrophils' inflammatory response, which also result in granulomatous formation (9).

Smoking can also damage vascular walls, in which affects the formation of prostacyclin and increases interactions between blood vessel walls and platelets (23). It is presumed that carbon monoxide and nicotine both damage blood vessels (23,24). The vascular endothelium may get injured as a result of this damage to the vessel wall resulting in an inflammatory response and affecting the progression of the disease. (25).

Endothelial cells have the ability to initiate and enhance the development of vasculitic lesions, which impairs endothelium-dependent vasodilation (26,18). In addition, exposure to cigarette smoke triggers the release of inflammatory cytokines that activate cellular signaling pathways through the adhesion of immunocytes and endothelial cells, leading to intravascular thrombosis and vasculitis (23). Several molecules serve significant roles in mediating cell-to-cell adhesion, including cell-to-cell adhesion molecule-1 (ICAM-1) and vascular cell adhesion molecule-1 (VCAM-1) (27,28). The presence of both molecules in excess in blood vessels that are affected suggests that they play a part in the development of the disease (29).

#### **RELATIONSHIP BETWEEN SMOKING AND BUEYER'S DISEASE**

Smoking has become a risk factor for the occurrence of TAO, several existing diagnostic criteria include smoking or smoking history as one of the points in assisting the diagnosis of TAO. Research on the relationship or effect of smoking on TAO is still very limited. Moreover, the majority of the patients in Sasaki et al. (2000) study, which included 850 patients (771 males and 79 females), had a history of smoking (93.2%). Patients who failed to quit smoking had a much higher chance of developing ulcer (odds ratio 51.71, 95%CI51.19–2.47; P50.004) and having an amputation (odds ratio 52.73, 95% CI51.86–4.01; P<0.0001) (11). Another study conducted by Fazeli et al. (2012), found that the duration of smoking had a significant relationship with the complications in the form of major amputation (P=0.004) in 108 (103 males and 5 females) patients diagnosed with TAO using the Shionoya criteria based on multivariate analysis. However, the results of multivariate analysis could not demonstrate the effect of daily cigarette consumption, age of disease onset, sex, and limb salvage from amputation (10). Furthermore, in a retrospective study of 224 patients (173 males and 51 females), Joncour et al. (2018), discovered that patients who quit smoking have a lower risk of amputation than those who continued to smoke (P=0.001). Based on 48 patients who underwent their first amputation, 5 (10.4%) were former smokers, and 43 (89.6%) were active smokers. Meanwhile, a total of 34 patients had undergone subsequent amputations; 8 (23.5%) were former smokers, and 26 (76%) were patients who continued to smoke. These findings suggest that smoking cessation can improve amputation-free survival in TAO patients. Further research is needed to fully understand the effects of smoking and its relationship to the extremities (30).

#### **DIAGNOSIS**

The diagnosis of TAO can be established through anamnesis, physical examination, and supporting examinations. A smoking history is usually obtained during anamnesis, along with information on any claudication in the fingers, feet, or hands. Most patients diagnosed with TAO are usually smokers. Thromboangiitis obliterans can also occur to those who have previously consumed chewing tobacco or tobacco products. TAO is very likely to occur for those who smoke at least one and a half packs per day (31,32).

Physical examination in TAO patients usually shows Raynaud's phenomenon, namely changes in skin color to become paler when in a cold environment (33). Raynaud's phenomenon occurs in about 40% of patients with Buerger's disease (15). Allen's test is performed to test the extent of the initial disease and assess the vascularity of the hand region (9). The procedure for Allen's test is that the patient is asked to clench his hand and the examiner presses on the patient's wrist intending to obstruct blood flow to the hand. TAO patients usually negative or have slowing of blood flow to the hands (5,34).



There are several criteria that can be used to diagnose TAO, including Shionoya, Olin, and Papa (36,37,38). First, there are five Shionoya criteria listed as follows: smoking history; onset of disease before the age of 50; infrapopliteal artery occlusion; upper extremity involvement or phlebitis migrans; and no additional risk factors for atherosclerosis besides smoking (36). Next, there are Olin's criteria, which include the following: onset under 45 years; a history of tobacco use; the presence of distal limb ischemia with indications for claudication, rest pain, ischemic ulceration or gangrene, and documented by non-invasive vascular testing; excluding autoimmune diseases, hypercoagulability, and diabetes mellitus; excluding emboli originating proximally using echocardiography or arteriography; and findings confirmed using arteriography of clinically associated and unrelated extremities (37). The last set of criteria is Papa's criteria, which have the same criteria as the previous ones but a lower age of onset, specifically between 30-40 years, as well as Raynaud's syndrome, which is characterized by intermittent foot claudication, single limb involvement (a negative criterion), and female gender (a negative variation) (38). Among the aforementioned criteria, the Shionoya clinical criteria are one of the most widely utilized and acknowledge worldwide (16).

Supporting examinations for TAO disease investigations are angiogram, vascular biopsy, and histopathological examination. Despite the fact that a number of supporting examinations have already been mentioned, TAO cannot be diagnosed by any particular investigations (9,32). Angiographic findings in TAO patients are corkscrew-shaped collaterals or Martorell's sign, indicating compensatory changes in the vasa vasorum due to segmental lesions or due to occlusion of the distal extremities (39). Martorell's sign can be detected in systemic lupus erythematosus (SLE), scleroderma, and other small vessel occlusive diseases, hence it is not a pathognomonic sign of TAO (40). A biopsy is used in atypical patients, such as elderly patients or patients with large artery disease. A histopathological examination typically indicates thrombus infiltration, polymorphonuclear leukocytes (PMN), and the presence of multinuclear giant cells (MN) in the associated vessels (6).

### TREATMENT

There is currently no cure for TAO. The most effective method of TAO management is smoking cessation. The use of smokeless tobacco, such as chewing tobacco or applying nicotine patches, can affect the development of disease (6,42). It is essential to educate about how exposure to cigarettes or tobacco products affects the development of TAO (42,43). In addition to smoking cessation, supportive therapies can be used if there are ulcers on the extremities to maintain maximum blood flow and prevent secondary infection (15). Platelet inhibitors, vasodilators, anticoagulants, anti-inflammatories, thrombolytics, and prostacyclin analogues are a few examples of the medications that can be used to provide pharmacological therapy. Calcium channel blockers and prostaglandin E1 are vasodilators that are commonly utilized and effective in TAO. Meanwhile, the most effective inhibitor of platelet aggregation is ticlopidine (44). The use of anti-inflammatory steroids has not demonstrated a significant effect (15,44). Prostacyclin analogs, such as iloprost and intra-arterial thrombolytics with streptokinase, are effective at managing pain that occurs at rest and reducing the risk of amputation (44). Furthermore, spinal cord stimulation, sympathectomy, vascular endothelial growth factor gene therapy, and amputation are examples of non-pharmacological therapy for TOA disease (15,45,46).

### PROGNOSIS

Data on mortality of TAO is few or difficult to find because TAO deaths are uncommon.

The US CDC reported that 117 deaths in the United States were caused by TAO disease between 1999 and 2007 (9). The risk of long-term amputation due to TAO disease is 25% every 5 years, 38% every 10 years, and 46% every 20 years (41). Furthermore, in another study, the prevalence of amputation was 26% and 34% at 10 years and 15 years later, respectively (30). Smoking cessation has a preventive effect since it is associated with reducing the risk of amputation. Moreover, if the patient is able to quit smoking, surgery is also less likely to be required (6,32).

### CONCLUSIONS

Thromboangiitis obliterans is an inflammatory vascular disease with no identifiable cause and is closely associated with smoking and the use of tobacco products. This disease is characterized by ischemia due to occlusion of the distal vessels in the form of intermittent claudication, which can progress to persistent ischemic pain at rest. The development of the disease may be influenced by smoking and using smokeless tobacco. Smoking cessation has a preventive effect since it is associated with reducing the risk of amputation and surgery.

### CONFLICT OF INTEREST

There is no conflict of interest to declare.

### REFERENCES

- [1] Olin, J. W. (2018). Thromboangiitis obliterans: 110 years old and little progress made. *Journal of the American Heart Association*, 7(23), 1-3.
- [2] Fazeli, B. & Rezaee, S. A. (2011). A review on thromboangiitis obliterans pathophysiology: thrombosis and angiitis, which is to blame? *Vascular*, 19(3), 141-53.
- [3] Mohareri, M., Mirhosseini, A., Mehraban S., & Fazeli, B. (2018). Thromboangiitis obliterans episode: autoimmune flare-up or reinfection? *Vascular Health and Risk Management*, 14, 247-251.
- [4] Jamadi, G. J., Wirka, I. M., & Sarifuddin. (2019). Thromboangiitis obliterans (buerger's disease). *Jurnal Medical Profession (MedPro)*, 1(1), 32-38.
- [5] Ramin, M., Salimi, J., & Meysamie, A. (2014). An Iranian scoring system for diagnosing buerger's disease. *Acta Med Iran*, 52(1), 60-65.
- [6] Rivera-Chavarría, I. J. & Brenes-Gutierrez, J. D. (2016). Thromboangiitis obliterans (buerger's disease). *Annals of Medicine and Surgery*, 7, 79-82.
- [7] Joviliano, E. E., Dellalibera-Joviliano, R., Dalio, M., Evora, P. R., & Piccinato, C. E. (2009). Etiopathogenesis, clinical diagnosis and treatment of thromboangiitis obliterans—current practices. *International Journal of Angiology*, 18(3), 119-125.
- [8] Olin, J. W. (2000). Thromboangiitis obliterans (buerger's disease). *The New England Journal of Medicine*, 343(12), 864-869. [3] Mohareri, M., Mirhosseini, A., Mehraban S., & Fazeli, B. (2018). Thromboangiitis obliterans episode: autoimmune flare-up or reinfection? *Vascular Health and Risk Management*, 14, 247-251.
- [9] Qaja, E., Muco, E., & Hashmi, M. F. (2022). *Buerger Disease*. Florida, USA: StatPearls Publishing.
- [10] Fazeli, B., Ravari H., & Assadi R. (2012). Natural history definition and a suggested clinical approach to buerger's disease: a case-control study with survival analysis. *Vascular*, 20(4), 198-202.

- [11] Sasaki, S., Sakuma, M., & Yasuda, K. (2000). Current status of thromboangiitis obliterans (Buerger's disease) in Japan. *International Journal of Cardiology*, 75(1), 175-181. [9] Qaja, E., Muco, E., & Hashmi, M. F. (2022). *Buerger Disease*. Florida, USA: StatPearls Publishing
- [12] Sasaki, S., Sakuma, M., Kunihara, T., & Yasuda, K. (2000). Distribution of arterial involvement in thromboangiitis obliterans (Buerger's disease): results of a study conducted by the intractable vasculitis syndromes research group in Japan. *Surgery Today*, 30(7), 600-605.
- [13] Cacione, D. G. (2018). *Buerger's Disease: Clinical Aspects and Evidence-Based Treatments*. Pp. 89-101. Cairo, Egypt: IntechOpen.
- [14] Vijayakumar, A., Tiwari, R., & Prabhuswamy V. K. (2013). Thromboangiitis obliterans (Buerger's disease)-current practices. *International Journal of Inflammation*, 13, 1-7.
- [15] Arkkila, P. E. T. (2006). Thromboangiitis obliterans (Buerger's disease). *Orphanet Journal of Rare Diseases*, 1(14), 1-5. [9] Qaja, E., Muco, E., & Hashmi, M. F. (2022). *Buerger Disease*. Florida, USA: StatPearls Publishing
- [16] Fazeli, B., Ligi, D., Keramat, S., Maniscalco, R., Sharebiani, H., & Mannello, F. (2021). Recent updates and advances in Buerger disease (thromboangiitis obliterans): biomolecular mechanisms, diagnostics and clinical consequences. *Diagnostics*, 11(10), 1736.
- [17] Liew, N. C., Lee, L., Hanipah, Z. N., Gee, T., & Jabar, M. F. (2015). Pathogenesis and management of Buerger's disease. *The International Journal of Lower Extremity Wounds*, 14(3), 231-235.
- [18] Igari, K., Inoue, Y., & Iwai, T. (2016). The epidemiologic and clinical findings of patients with Buerger's disease. *Annals of Vascular Surgery*, 30, 263-269. [9] Qaja, E., Muco, E., & Hashmi, M. F. (2022). *Buerger Disease*. Florida, USA: StatPearls Publishing.
- [19] Arekhi, S., Ghodsi, A., Omranzadeh, A., & Rahimi, H. R. (2021). Does adaptive T cell immunity have any role in the pathophysiology and histopathology of Buerger's disease? *Journal of Basic Research in Medical Sciences*, 8 (1), 1-9.
- [20] Kasuma, U. (2014). Relation of zinc consumption as cofactor in Minangkabau traditional food with neutrophil elastase level in gingival crevicular fluid in periodontal disease. *Dentika Dental Journal*, 18(2), 105-109.
- [21] Iwai, T., Inoue, Y., Umeda, M. H. Y., Kurihara, N., & Koike, M. (2005). Oral bacteria in the occluded arteries of patients with Buerger disease. *Journal of Vascular Surgery*, 42(1), 107-115.
- [22] Williams, G. (1969). Recent views on Buerger's disease. *Journal of Clinical Pathology*, 22(5), 573-8.
- [23] Ketha, S. S., & Cooper, L. T. (2013). The role of autoimmunity in thromboangiitis obliterans (Buerger's disease). *Annals of the New York Academy of Sciences*, 1285:15-25.
- [24] Leone, A. (2005). Biochemical markers of cardiovascular damage from tobacco smoke. *Current Pharmaceutical Design*, 11(17), 2199-2208.
- [25] Małeck, R., Kluz, J., Przędziecka-Dołyk, J., & Adamiec, R. (2015). The pathogenesis and diagnosis of thromboangiitis obliterans: is it still a mystery? *Advances in Clinical and Experimental Medicine*, 24(6), 1085-1097.
- [26] Azizi, M., Boutouyrie, P., Bura-Rivière, A., Peyrard, S., Laurent, S., Fiessinger, J. N. (2010). Thromboangiitis obliterans and endothelial function. *European Journal of Clinical Investigation*, 40(6), 518-526.
- [27] Palmefors, H., DuttaRoy, S., Rundqvist, B., & Borjesson, M. (2014). The effect of physical activity or exercise on key biomarkers in atherosclerosis: a systematic review. *Atherosclerosis*, 235(1), 150-61.
- [28] Long, E. O. (2011). ICAM-1: getting a grip on leukocyte adhesion. *Journal of Immunology*, 186(9), 5021-5023.
- [29] Caridi, D. G., Bitto, A., Massara, M., Pallio, G., Pizzino, G., Serra, R., Altavilla, D., Squadrito, F., & Spinelli, F. (2016). Increased serum HMGB-1, ICAM-1 and metalloproteinase-9 levels in Buerger's patients. *Current Vascular Pharmacology*, 14(4), 382-387.
- [30] Joncour, A. L., Soudet, S., Dupont, A., Espitia, O., Koskas, F., Cluzel, P., Hatron, P. Y., Emmerich, J., Cacoub, P., Rigon, M. R., Lambert, M., Saadoun, D., & Network, F. B. (2018). Long-term outcome and prognostic factors of complications in thromboangiitis obliterans (Buerger's disease): a multicenter study of 224 patients. *Journal of the American Heart Association*, 7(23), 1-9.
- [31] Oktaria, D., & Samosir, R. K. (2017). Kriteria diagnosis dan tatalaksana pada Buerger's disease. *Journal of Majority*, 6(2), 126-131.
- [32] Apriliana, S. (2021). Thromboangitis obliterans (TAO): diagnosis dan tatalaksana. *Cermin Dunia Kedokteran*, 48(12), 713-717.
- [33] Musa, R., & Qurie, A. (2022). *Raynaud Disease*. Florida, USA: StatPearls Publishing.
- [34] Foreman, A., Almeida, J. R. D., Gilbert, R., & Goldstein, D. P. (2015). The Allen's test: revisiting the importance of bidirectional testing to determine candidacy and design of radial forearm free flap harvest in the era of trans radial endovascular access procedures. *Journal of Otolaryngology-Head and Neck Surgery*, 44(47), 1-5.
- [35] Fazeli, B., Ligi, D., Keramat, S., Maniscalco, R., Sharebiani, H., & Mannello, F. (2021). Recent updates and advances in Buerger disease (thromboangiitis obliterans): biomolecular mechanisms, diagnostics and clinical consequences. *Diagnostics*, 11(10), 1736.
- [36] Shionoya, S. (1998). Diagnostic criteria of Buerger's disease. *International Journal of Cardiology*, 66(1), 243-45.
- [37] Olin, J. W., & Shih, A. (2006). Thromboangiitis obliterans (Buerger's disease). *Current Opinion in Rheumatology*, 18(1), 18-24.

- [38] Papa, M. Z., Rabi, I., & Adar R. (1996). A point scoring system for the clinical diagnosis of Buerger's disease. *European Journal of Vascular and Endovascular Surgery*, 11(3), 335-339.
- [39] Gallagher, K. A., Tracci, M. C., & Scovell, S. D. (2013). Vascular arteritides in women. *Journal of Vascular Surgery*, 57(4), 27-36.
- [40] Dimick, S., Goh, A., Cauzza, E., Steinbach, L., Baumgartner, I., Stauffer, E., Voegelin, E., & Anderson S. E. (2012). Imaging appearances of buerger's disease complications in the upper and lower limbs. *Clinical Radiology*, 67(12), 1207-1211.
- [41] Cacione, D. G., Novaes, F. D. C., & Moreno, D. H. (2018). Stem cell therapy for treatment of thromboangiitis obliterans (buerger's disease). *Cochrane Database of Systematic Review*, 10(10), 1-29.
- [42] Piazza, G., & Creager, M. A. (2010). Thromboangiitis obliterans. *Circulation*, 121(16):1858-1861.
- [43] Hernawan, Heri. (2016). Tromboangiitis obliterans dengan komorbid DVT. *Syifa' Medika: Jurnal Kedokteran dan Kesehatan*, 6(2), 66-73.
- [44] Khanna, A. K., & Puneet, M. S. (2011). *Manual of Vascular Surgery*. New Delhi, India: Jaypee Medical Ltd.
- [45] Klomp, H. M., Steyerberg, E. W., Habbema, J. D. F., & Eses S. G. (2009). What is the evidence on efficacy of spinal cord stimulation in (subgroups of) patients with critical limb ischemia? *Annals of Vascular Surgery*, 23(3), 355-363.
- [46] Gersbach, P. A., Argitis, V., Gardaz, J. P., Segesser, V. L. K., & Haesler, E. (2007). Late outcome of spinal cord stimulation for unreconstructable and limb-threatening lower limb ischemia. *European Journal of Vascular and Endovascular Surgery*, 33(6), 717-724.

