




NOW on PubMed

Saudi Dental Journal



The official publication of
Saudi Dental Society

ISSN 1013-9052

Saudi Dental Journal

Editor-in-Chief

HAMDAN ALGHAMDI

College of Dentistry, Dept. of Periodontics and Community Dentistry, King Saud University, Riyadh,
Kingdom of Saudi Arabia

Scientific Editor

CHALINI SUNDAR

College of Dentistry, King Saud University, Riyadh, Kingdom of Saudi Arabia

Editorial Board

THAKIB ALSHALAN

College of Dentistry, Al Farabi University

IBRAHEEM AL SHIDDI

College of Dentistry, Dept. of Prosthetic Dental Sciences,
King Saud University, Riyadh, Kingdom of Saudi Arabia

NADIR BABAY

College of Dentistry, Dept. of Periodontics and Community Dentistry,
King Saud University, Riyadh, Kingdom of Saudi Arabia

HANAN BALTO

College of Dentistry, Princes Noura University, Riyadh, Saudi Arabia

HESHAM KHALIL ALMOALLIM

College of Dentistry, Dept. of Oral and Maxillofacial Surgery,
King Saud University, Riyadh, Kingdom of Saudi Arabia

RITA KHOUNGANIAN

College of Dentistry, Division of Oral Pathology,
Biology and Microbiology,
Dept. of Oral Medicine and Diagnostic Sciences,
King Saud University, Riyadh, Kingdom of Saudi Arabia

ABDULGHANI MIRA

Faculty of Dentistry,
King Abdulaziz University (KAUFD) Jeddah,
Saudi Arabia

Advisory Board

ALEXANDER HEMPRICH

Dept. of Oral, Maxillofacial and Facial Plastic Surgery,
Faculty of Dentistry, Universitat Leipzig, Leipzig, Germany

JOHN JANSEN

Dept. of Periodontology and Biomaterials, Radboud University,
Nijmegen, The Netherlands

REMESH K. KUBA

School of Dentistry, Dept. of Diagnostic and Biological Sciences,
University of Minnesota, Minneapolis, USA

EDWIN A. MCGLUMPHY, JR.

Restorative Science and Prosthodontics, The Ohio State University,
Ohio, USA

STEVE MORGANO

Restorative Dentistry, Rutgers School of Dental Medicine, Newark, USA

COLIN ALEXANDER MURRAY

Restorative Dentistry and Director Advanced Resto and Prosth
Programme, European University College, Dubai, UAE

YOSHIKI ONO

Dept. of Pediatric Dentistry Graduate School,
Tokyo Medical and Dental University, Tokyo, Japan

STEPHEN. F. ROSENTIEL

College of Dentistry, Restorative and Prosthetic Dentistry,
The Ohio State University, Health Sciences Center Columbus, USA

AXEL RUPRECHT

College of Dentistry, Dept. of Oral Pathology,
Radiology and Medicine, The University of Iowa, Iowa, USA

LAKSHMAN SAMARANAYAKE

Dean of Dentistry and Chair of Oral Microbiology,
The University of Hong Kong,
Prince Philip Dental Hospital, Sai Ying Pun, Hong Kong

WOOSUNG SOHN

Dept. of Health Policy and Health Services,
Advanced Specialty Education Program in Dental Health,
Henry M. Goldman School of Dental Medicine, Boston University,
Boston, USA

THOMAS VAN DYKE

Research and Chair of Periodontology,
The Forsyth Institute, Cambridge, USA

Biostatistician

NASSR AL-MAFLEHI

College of Dentistry, King Saud University, Riyadh, Kingdom of Saudi Arabia

The Saudi Dental Journal

Open access

[Submit your article](#)


[Guide for authors](#)

Menu

 Search in this journal

Volume 35, Issue 4

Pages 283-386 (May 2023)

 [Download full issue](#)

[< Previous vol/issue](#)

[Next vol/issue >](#)

Receive an update when the latest issues in this journal are published

[Sign in to set up alerts](#)

Open access

Editorial Board

Page ii

 [View PDF](#)

FEEDBACK 

The Saudi Dental Journal

Open access

[Submit your article](#)

[Guide for authors](#)

Menu



Fatih S. Almorad, Khaled Alkows, Abdurrahman Almutbarak, Foziah Waheed Almadhoon, Naima Asim

Pages 283-293



[View PDF](#)

[Article preview](#)

Review article *Open access*

Periodontal tissue changes after crown lengthening surgery: A systematic review and meta-analysis

Samantha C. Smith, Rayner Goh, Sunyoung Ma, Getulio R. Nogueira, ... Andrew Tawse-Smith

Pages 294-304



[View PDF](#)

[Article preview](#)

Review article *Open access*

Factor XIII deficiency in the Saudi population, an underestimated bleeding risk. Review article and an illustrative case report with dental complications

Mansour Aljabry

Pages 305-309



[View PDF](#)

[Article preview](#)

ORIGINAL RESEARCH ARTICLES

Research article *Open access*

Novel modified quantitative index for cone-beam CT assessment of medication-related osteonecrosis of the jaw - comparative analysis with original index

Ghaidaa Badabaan, Maxine Strickland, Steven R Singer, Adriana G. Creanga, Gayathri Subramanian

Pages 310-316

[FEEDBACK](#)

The Saudi Dental Journal

Open access

[Submit your article](#)[Guide for authors](#)

Menu



Siti Kurnati, Devi Eka Juniarti, Tulanti Kartini Sunari, Melidi Kurnia Ariani

Pages 317-321

[View PDF](#)[Article preview](#) Research article [Open access](#)

Inhibiting the growth of periopathogenic bacteria and accelerating bone repair processes by using robusta coffee bean extract

Desi Sandra Sari, Peni Pujiastuti, Dwi Warna Aju Fatmawati, Mega Ayu Mardiyana, ... Yuliana Mahdiyah Daat Arina

Pages 322-329

[View PDF](#)[Article preview](#) Research article [Open access](#)

Orofacial anatomical and occlusal changes in patients with sickle cell disease in Kuwait

Hussain Dashti, Devipriya Sundaram, Aref Ghayyath, Hassan Al-Jafar, Mariam Baghdady

Pages 330-336

[View PDF](#)[Article preview](#) Research article [Open access](#)

Vitamin C influences antioxidative, anti-inflammatory and wound healing markers in smokers' gingival fibroblasts *in vitro*

Ruaa Alyami, Reham Al Jasser, Fahad Ali Alshehri, Nouf Alshibani, ... Abdurahman A. Niazy

Pages 337-344

[View PDF](#)[Article preview](#) [FEEDBACK](#)

The Saudi Dental Journal

Open access

[Submit your article](#)

[Guide for authors](#)

Menu



[View PDF](#) [Article preview](#)

Research article *Open access*

A radiographic analysis of the location of the mental foramen

Abdalmalik O. Ghandourah, Mohd. B. Badaoud, Anmar Dahlawi, Abdullah Alghamdi, ... Abrar K. Demyati
Pages 354-358

[View PDF](#) [Article preview](#)

Research article *Open access*

Effect of staining and whitening systems on color stability of computer aided design/computer aided manufacturing lithium disilicate glass ceramic

Nadia S. Al-Angari, Shahad Quwayhis, Nouf Almayouf, Shahad Meaigel, ... Sarah S. Al-Angari
Pages 359-364

[View PDF](#) [Article preview](#)

Research article *Open access*

Comparative evaluation of the accuracy of 3D-printed denture teeth

Haidar Alalawi, Faisal D. Al-Qarni, Mohammed M. Gad
Pages 365-371

[View PDF](#) [Article preview](#)

Research article *Open access*

FEEDBACK

The Saudi Dental Journal

Open access

[Submit your article](#)

[Guide for authors](#)

Menu



Research article *Open access*

Effect of multidrug solution for the treatment of chemotherapy-induced oral mucositis *in vivo*

Rebecca Rhuanny Tolentino Limeira, Isabella Lima Arrais Ribeiro, Paulo Rogério Ferreti Bonan, Danielle da Nóbrega Alves, ... Ricardo Dias de Castro

Pages 378-385



[View PDF](#)

[Article preview](#)

[← Previous vol/issue](#)

[Next vol/issue >](#)



Copyright © 2023 Elsevier B.V. or its licensors or contributors.
ScienceDirect® is a registered trademark of Elsevier B.V.



FEEDBACK



ORIGINAL ARTICLE

Inhibiting the growth of periopathogenic bacteria and accelerating bone repair processes by using robusta coffee bean extract



Desi Sandra Sari^{a,*}, Peni Pujiastuti^a, Dwi Warna Aju Fatmawati^b,
Mega Ayu Mardiyana^c, Ayu Tri Wulandari^c, Yuliana Mahdiyah Daat Arina^a

^a Department of Periodontology Faculty of Dentistry, University of Jember, Jember 68121, Indonesia

^b Department of Restorative Dentistry, Faculty of Dentistry, University of Jember, Jember 68121, Indonesia

^c Faculty of Dentistry, University of Jember, Jember 68121, Indonesia

Received 22 November 2022; revised 19 March 2023; accepted 20 March 2023

Available online 24 March 2023

KEYWORDS

Periodontitis;
Remodelling;
Coffee;
Osteoblast;
Osteoclast

Abstract *Background:* Periodontitis is an inflammatory disease of the teeth-supporting tissues caused by microorganisms. Robusta coffee bean extract has antibacterial properties due to its caffeine, flavonoids, trigonelline, and chlorogenic acid contents. The robusta coffee bean extract also regulates alveolar bone healing through bone remodelling.

Aim: The study aimed to investigate robusta coffee bean extract to inhibit bacterial growth and accelerate bone repair in vitro and in vivo.

Methods: This study used the paper disc diffusion method with the research group of robusta coffee bean extract with concentrations of 50%, 25%, 12.5%, 6.25%, and negative control, as much as 20 and dripped onto the disc paper then placed on the surface of the agar media that had been inoculated with bacteria. The diameter of the inhibition zone was measured. Twenty periodontitis rat models were given 0.05 ml of the robusta coffee bean extract on the molars and put in a periodontal pocket for seven days. Rats were decapitated, and alveolar bone tissues were stained with HE and IHC staining. The number of osteoclasts, osteoblasts and BMP-2 was counted using a microscope. Statistical test with *Kruskal Wallis* followed by *Mann Whitney* showed a p-value of < 0.05.

* Corresponding author at: Jln, Kalimantan 37, Jember 68121, Indonesia.

E-mail addresses: desi_sari.fkg@unej.ac.id (D.S. Sari), Peni.pujiastuti@unej.ac.id (P. Pujiastuti), dwiwarna.fkg@unej.ac.id (D.W.A. Fatmawati), 181610101138@mail.unej.ac.id (M.A. Mardiyana), 181610101068@mail.unej.ac.id (A.T. Wulandari), yuliana.mda@unej.ac.id (Y.M.D. Arina).

Peer review under responsibility of King Saud University. Production and hosting by Elsevier



Production and hosting by Elsevier

Results: The average diameter of the inhibitory zone of robusta coffee bean extract showed that the *P. gingivalis* group of bacteria was higher than that of *A. actinomycetemcomitans* and *S. viridans* ($p < 0.05$) with a concentration of 50%. The average number of osteoblast cells increased, and the average number of osteoclast cells decreased in the 50% concentration group compared to the other groups ($p < 0.05$). BMP-2 expression in the robusta coffee bean extract group was 50% higher than in the other groups.

Conclusion: Robusta coffee bean extract has a periopathogenic antibacterial and accelerates alveolar bone repair.

© 2023 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Periodontitis is an inflammatory disease of the teeth-supporting tissues caused by *P. gingivalis*, *A. actinomycetemcomitans*, a periopathogenic bacterium and *S. viridans* that cause pulp-periapical abnormalities. Among the major periopathogenic bacteria, *P. gingivalis* is one of the main etiological bacteria in the pathogenesis and development of inflammation in periodontal disease because of its ability to adhere to the oral epithelium, invade the oral epithelium, and possess virulence factors (Shewale et al., 2016; Wahyukundari et al., 2017).

Periodontitis resulting in progressive destruction of the periodontal ligament and alveolar bone resorption shows an imbalance between osteoclasts and osteoblasts (Park et al., 2017). These cellular activities include the resorption of old bone by osteoclasts and the formation of new bone by osteoblasts (Fakhry et al., 2013; Florencio-Silva et al., 2015; Koga et al., 2016).

Treatment of periodontitis can be surgical or non-surgical. Scaling and root planing (SRP) is a non-surgical treatment which is the gold standard in treating periodontal disease. However, this treatment also has limitations. Therefore, additional antimicrobial therapy is used to eliminate or reduce the number of bacterial pathogens. Other antimicrobial treatment can be given locally as it does not cause side effects. People are now turning to use herbs for treatment (Gigi et al., 2022; Panduwawala et al., 2017).

Coffee is a popular plant consumed by Indonesian people, including those living in Jember Regency. Jember Regency has abundant Robusta coffee. The part of coffee plants commonly consumed is the coffee beans. Coffee beans are naturally rich in caffeine, flavonoids, trigonelline and chlorogenic acid, which have an antibacterial properties. The active compounds contained in robusta coffee beans are antibacterial, which destroy amino acids, building blocks of cell walls and DNA, causing changes in bacterial genesis and lysis and disrupting the stability of the bacterial cytoplasmic membrane (Anand et al., 2015; Ojeh et al., 2016).

This study aimed to investigate the effect of coffee extract in inhibiting the growth of periopathogenic bacteria and in accelerating bone repair processes in vitro and in vivo.

2. Materials and methods

This study was divided into 2, in vitro to see the inhibition of periopathogenic bacteria and in vivo to see the accelerating bone growth.

2.1. Ethical clearance

The ethical clearance for the procedure for treating animals in this study was approved Ethics Committee of the Faculty of Dentistry, the University of Jember, with letter number No.1274/UN25.8/KEPK/DL/2021.

2.2. Robusta coffee beans extract preparation

Five hundred grams of Robusta coffee beans were extracted into powder using the maceration method. The powder was put into a closed container and then soaked using 96% ethanol in a ratio of 1:5 for 72 h and stirred twice a day. The immersion results were filtered using filter paper to produce filtrate and residue. The filtrate was evaporated using a rotary evaporator to obtain viscous extract brown robusta coffee bean extract. The extract with a concentration was diluted using aqua dest with the serial dilution method with a concentration of 50%, 25%, 12.5%, 6.25%, and aqua dest (Macedo et al., 2015).

2.3. Antibacterial sensitivity

The antibacterial test treatment stage was carried out according to the modified NCCLS (National Committee for Clinical Laboratory Standards) standard protocol. Bacterial inoculation was prepared on agar media with a sterile cotton swab. 1 ml of bacterial suspension was taken using a syringe. Inoculate with streaking motion (zig-zag) on the entire media surface. 20 L of robusta coffee bean extract was dropped onto the paper disc with concentrations of 50%, 25%, 12.5%, 6.25%, and negative control (equates) using a micropipette. The Petri dish was closed and then put in an inverted position into a desiccator to create anaerobic conditions. Then, the desiccator was put into the incubator to be incubated for 24 h at 37 °C (Alcayaga-Miranda et al., 2017; Rurenga et al., 2013).

2.4. Inhibition zone diameter

After incubation for 24 h, the Petri dish was removed from the desiccator, and a zone of inhibition was seen, indicated by a clear area around the paper disc. The diameter of the inhibition zone was measured by turning the Petri dish. Observation and measurement of the diameter of the inhibition zone around the paper disc in each research group were conducted using a digital calliper (Magnani et al., 2014).

2.5. Periodontitis rat model

Twenty male Wistar rats aged 2–3 months with a body weight of ± 200 –250 g were modelled for periodontitis by inducing *P. gingivalis* in the buccal gingival sulcus of the lower molars as much as 0.05 ml and given once every three days for 14 days using a tuberculin syringe with a needle size of 30 gauge. Periodontitis in rats can be seen by several examinations, such as through clinical and radiographic features. Clinically, there was swelling at the induction site, a redder colour of the gingiva and spontaneous bleeding. (Hienz et al., 2015; Sugihartini et al., 2017).

2.6. The administration of robusta coffee bean

Robusta coffee bean extract was rinsed in the periodontal pocket of the lower molars with as much as 0.05 ml using a syringe. Group I was made a periodontitis model and given robusta coffee bean extract with a concentration of 6.25%, Group II was made a periodontitis model and given robusta coffee bean extract with a concentration of 25%, Group III was made a periodontitis model and was offered robusta coffee bean extract with a concentration of 12.5%, Group III IV was made a periodontitis model and given robusta coffee bean extract with a concentration of 50%. The administration was carried out for seven days, after which the rats were euthanized.

2.7. Sample preparation

First, the experimental animals (2–3 rats) were put into a closed jar (16L size) which contained 5 ml of cotton/tissue moistened with ether. After the rat was unconscious, it was decapitated to take the mandibular bone; then, the mandible was put into a 10% formalin buffer solution for ± 8 h. The goal was to prevent the bone specimens from being damaged and to fix the bone specimens.

2.8. Hematoxylin-Eosin staining

The tissue was cut using a microtome. After that, the cutting thickness setting of the incision was changed to a size of 6 μ m, and then the paraffin block containing the tissue was cut. The cut was transferred with a brush to the surface of the water bath with a temperature of 37 °C–40 °C so that the incision expands well. The incision results were then

transferred to an object Tissue staining was carried out using Mayer's Hematoxylin-Eosin method with the phases of defaration, dehydration, core staining, cytoplasmic staining, dehydration, clearing, and mounting (Sugihartini et al., 2017).

2.9. Cell count stage

Cells were counted using a light microscope (Olympus) with 1000x magnification. The counting area consisted of three visual fields in the section of the rat alveolar bone for each sample and was calculated using the computer program Adobe Photoshop CC 2021.

2.10. Immunohistochemical staining (IHC)

The immunohistochemical staining used in this study was for the examination of BMP-2. The method for colouring BMP-2 was by slicing the tissue with a microtome placed on a glass object and then deparaffinized, i.e. pulling/removing the paraffin present in the tissue.

2.11. Statistical analysis

The results of the test show that the data were not normally distributed and not homogeneous. The Kruskal-Wallis non-parametric test was carried out, and then the Mann-Whitney test with a 95% confidence level ($\alpha = 0.05$; SPSS Version 23).

3. Results

The results of observations on the inhibitory ability of robusta coffee bean extract on the growth of *P. gingivalis*, *A. actinomycetemcomitans* and *S. viridans* are presented in Table 1.

The results showed that all bacteria had an inhibitory effect on robusta coffee bean extract. The growth inhibition of *P. gingivalis* bacteria was highest at 50% concentration of robusta coffee bean extract, which was 19.18 ± 0.18 mm compared to other bacteria. The lowest bacterial growth inhibition with a concentration of 12.5% was 8.40 ± 0.22 in *A. actinomycetemcomitans* bacteria compared to other bacteria (see Table 2).

The negative control had no inhibition with Robusta coffee bean extract, with a concentration of 6.25% in all bacterial groups. The results of the Mann-Whitney test showed that all treatment groups were significantly different from the negative control, with a concentration of 6.25% robusta coffee

Table 1 Average of Inhibitory zone and standard deviation of robusta coffee bean extract on periopathogenic bacteria.

Num	Group	N	Mean (mm) \pm Standart Deviation		
			<i>P.gingivalis</i>	<i>A. actinomycetemcomitans</i>	<i>S. viridans</i>
1	Negative control	4	0,00 \pm 0,00*	0,00 \pm 0,00*	0,00 \pm 0,00*
2	Concentration 50%	4	19,18 \pm 0,18*	16,15 \pm 0,12*	18,15 \pm 0,19*
3	Concentration 25%	4	16,59 \pm 0,17	12,20 \pm 0,10	14,95 \pm 0,10
4	Concentration 12,5%	4	13,14 \pm 0,24	8,40 \pm 0,22	12,15 \pm 0,25
5	Concentration 6,25%	4	0,00 \pm 0,00 ^{ns}	0,00 \pm 0,00 ^{ns}	0,00 \pm 0,00 ^{ns}

Description: * significant, ^{ns} non-significant.

Table 2 The number of Osteoblast cells and Osteoclast cells.

No	Groups	N	Mean ± Standart Deviation	
			Osteoclast cells	Osteoblast cells
1	Negative control	4	3.1 ± 8,1*	99,4 ± 5,9*
2	Concentration 50%	4	4,5 ± 5,9*	133 ± 5,1*
3	Concentration 25%	4	4 ± 7,5 ^{ns}	101 ± 9,7 ^{ns}
4	Concentration 12,5%	4	4,8 ± 6,7 ^{ns}	94,5 ± 8,2 ^{ns}
5	Concentration 6,25%	4	4,1 ± 9,1 ^{ns}	93 ± 7,8 ^{ns}

Description: * significant, ^{ns} non-significant.




Fig. 1 Rat alveolar bone tissue: (A) Negative control, (B) Concentration 50%, (C) Concentration 25%, (D) Concentration 12,5%, (E) Concentration 6.25% osteoblast cells (blue arrow), osteoclast cells (green arrow) (Hematoxylin-Eosin staining 400× magnitude).




Fig. 2 Expression of bone morphogenetic protein 2 (BMP-2) in rat alveolar bone: Description: * significant, ^{ns} non-significant.

extract. The results showed an inhibition zone (clear zone) seen around the wells that had been given Robusta coffee extract 50%, 25%, 12.5%, 6.25%, and negative control (see Fig. 1).

Fig 2 shows the expression of BMP-2 in the 50% concentrated Robusta coffee bean group and the control group. There was no significant difference ($p > 0.05$), but in the 25%, 12.5%, and 6.25% treatment groups there were significant differences ($p < 0.05$) (see Fig. 3).

4. Discussion

The results of this study indicate that not all concentrations of robusta coffee bean extract have inhibitory power against periopathogenic bacteria. A concentration of 6.25% did not show any inhibition result, which was indicated by the absence of a clear zone around the wellbore (inhibition zone = 0). It was possible because the active compound of the antibacterial substance of robusta coffee bean extract in toothpaste was too small, so it was unable to inhibit bacterial growth. (Sukohar et al., 2011).

The ability of robusta coffee bean extract to inhibit bacteria is due to the presence of antibacterial compounds such as caffeine, flavonoids, trigonelline, and chlorogenic acid. These ingredients have antibacterial activity with different mechanisms of inhibiting bacterial growth (Eldin et al., 2016; Macedo et al., 2015).

Caffeine in robusta coffee beans works because it significantly influences the ability of alkaloid compounds; when in contact with bacteria, it will react with amino acid compounds that make up cell walls and bacterial DNA. This reaction will cause changes in the structure and arrangement of amino acids, which will change the performance of the DNA chain so that DNA damage will occur in the bacterial cell nucleus and support the lysis of the bacterial cell nucleus. Thus the bacteria will become inactive and destroyed (Kim et al., 2016; Li et al., 2016).

The above statement is supported by previous research that the caffeine and trigonelline content in the robusta coffee bean extract at concentrations of 1%, 1.25%, 1.5%, and 3% had

antibacterial effects against *P. gingivalis*. The mechanism of phenol compounds in killing bacteria is by denaturing cell proteins. Flavonoids play a role in inhibiting the function of bacterial cell membranes by forming complex compounds against extracellular and dissolved proteins that damage the bacterial cell membrane and are followed by the release of intracellular compounds that result in cell death. The antibacterial activity of chlorogenic acid increases the permeability of the outer membrane and plasma membrane, thereby reducing the defence function, nucleotide leakage and cytoplasmic contents. These compounds also reduce levels of Reactive Oxygen Species (ROS). The decrease in ROS levels results in the disruption of intracellular signalling in bacteria and accumulates Ca^{2+} levels as a proapoptotic agent, causing bacterial cell death due to apoptotic signals (Martin et al., 2022; Shushtari & Froushani, 2017).

On day 7, the group of periodontitis rats and the treatment group were given Robusta coffee bean chlorogenic acid gel had differences in the amount of OCN expression. This is in line with the results of research from Yamamoto showing that chlorogenic acid from coffee can increase the synthesis of IL-6 in osteoblasts. It can initiate bone formation (Nurman et al., 2019; Struppek et al., 2022). Other studies have shown that chlorogenic acid can enhance the osteogenic differentiation of human adipose tissue-derived mesenchymal stem cells (MSC) by increasing the mineralization of bone tissue. This indicates that the acid chlorogenic acid can increase the potential for osteogenesis (Maalik et al., 2016; Tian et al., 2019; Yamagata et al., 2018).

BMP can effectively induce mesenchymal cells to differentiate into osteoblasts and mesenchymal cell differentiation and initiate bone formation. Furthermore, BMP-2 plays a crucial role in bone formation, growth, and repair. A study by Sari. (2019) showed that BMP-2 could increase the concentration of Ca^{2+} and activate osteoblasts to differentiate the enhanced formation and integration of mineralized bone nodules. BMP-2 in vitro and in vivo studies induce bone formation by expressing bone markers (Ernie et al., 2017; Sari et al., 2019; Setiawatie et al., 2022).




Fig. 3 Rat alveolar bone tissue: expression BMP-2 (red arrow) (A) Negative control, (B) Concentration 50%, (C) Concentration 25%, (D) Concentration 12,5%, (E) Concentration 6.25% (Immunohistochemistry Staining 400x magnitude).

5. Conclusion

Robusta coffee bean extract has a periopathogenic antibacterial and alveolar bone repair ability.

Research funding

This research was funded by the University of Jember through Reworking Research Grant 2022 contract No.6332/UN25.3.1/LT/2022.

Author contributions

All authors have accepted responsibility for the entire content of this manuscript and approved its Submission.

CRedit authorship contribution statement

Desi Sandra Sari: Conceptualization, Data curation, Formal analysis, Supervision, Writing – review & editing. **Peni Pujiastuti:** Methodology, Writing – original draft. **Dwi Warna Aju Fatmawati:** Investigation, Supervision. **Mega Ayu Mardiyana:** Formal analysis. **Ayu Tri Wulandari:** Investigation, Visualization. **Yuliana Mahdiyah Daat Arina:** Investigation.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgment

The author would like to extend our uttermost gratitude to the Institute of Research and community Services of the University of Jember.

Appendix A. Supplementary material



Supplementary data to this article can be found online at <https://doi.org/10.1016/j.sdentj.2023.03.007>.

References

- Alcayaga-Miranda, F., Cuenca, J., Khoury, M., 2017. Antimicrobial activity of mesenchymal stem cells: Current status and new perspectives of antimicrobial peptide-based therapies. *Front. Immunol.* 8 (MAR), 1–15. <https://doi.org/10.3389/fimmu.2017.00339>.
- Anand, J., Upadhyaya, B., Rawat, P., Rai, N., 2015. Biochemical characterization and pharmacognostic evaluation of purified catechins in green tea (*Camellia sinensis*) cultivars of India. *3 Biotech* 5 (3), 285–294. <https://doi.org/10.1007/s13205-014-0230-0>.
- Eldin, I., Elgailani, H., Eldin, I., Elgailani, H., 2016. Spectrophotometric and Phytochemical Analysis of Black Tea (*Camellia sinensis* Leaves) Spectrophotometric and Phytochemical Analysis of Black Tea (*Camellia sinensis* Leaves). *J. Appl. Ind. Sci.* 3 (March), 1–6.
- Ernie Maduratna, Ulfah Noer, Wahjuningrum Dian Agustin, Desi Sandra Sari, Rubianto. 2017. Viability Bovine Tooth Hydroxiapatite on Bone Marrow Mesenchymal Stem Cells. In: International Medical Device and Technolgt Conference, pp. 100–104.
- Fakhry, M., Hamade, E., Badran, B., Buchet, R., Magne, D., 2013. Molecular mechanisms of mesenchymal stem cell differentiation towards osteoblasts. *World J. Stem Cells* 5 (4), 136–148. <https://doi.org/10.4252/wjsc.v5.i4.136>.
- Florencio-Silva, R., Sasso, G.R.D.S., Sasso-Cerri, E., Simões, M.J., Cerri, P.S., 2015. Biology of bone tissue: structure, function, and factors that influence bone cells. *Biomed Res. Int.* 2015. <https://doi.org/10.1155/2015/421746>.
- Gigi, M.K., Ramadhani, Y., Rahayu, R., Rahmasari, P., Prajnasari, K.N., Alhakim, M.M., Aljunaid, M., Mohammed, H., 2022. Review article A mucoadhesive gingival patch with Epigallocatechin-3-gallate green tea (*Camellia sinensis*) as an alternative adjunct therapy for periodontal disease : A narrative review. *114(158)*, 114–119. <https://doi.org/10.20473/j.djmg.v55.i2.p114>.
- Hienz, Stefan A., Paliwal, Sweta, Ivanovski, S., 2015. Mechanisms of bone resorption in periodontitis. *J. Immunol. Res.* 2015, 1–10. [https://doi.org/10.1016/8756-3282\(85\)90328-X](https://doi.org/10.1016/8756-3282(85)90328-X).
- Kim, H.J., Yoon, B.K., Park, H., Seok, J.W., Choi, H., Yu, J.H., Choi, Y., Song, S.J., Kim, A., Kim, J.W., 2016. Caffeine inhibits adipogenesis through modulation of mitotic clonal expansion and the AKT/GSK3 pathway in 3T3-L1 adipocytes. *BMB Rep.* 49 (2), 111–115. <https://doi.org/10.5483/BMBRep.2016.49.2.128>.
- Koga, T., Minamizato, T., Kawai, Y., Miura, K., Takashi, I., 2016. Bone regeneration using dentin matrix depends on the degree of demineralization and particle size. *PLOS o* 1–12. <https://doi.org/10.1371/journal.pone.0147235>.
- Li, A., Wu, N., Zou, H., Zhu, B., Xiong, S., Xiao, G., 2016. Low concentration of caffeine inhibits cell viability, migration and invasion, and induces cell apoptosis of B16F10 melanoma cells. *Int. J. Clin. Exp. Path.* 9 (11), 11206–11213.
- Maalik, A., Bukhari, S.M., Zaidi, A., Shah, K.H., Khan, F.A., 2016. Chlorogenic acid: A pharmacologically potent molecule. *Acta Poloniae Pharmaceutica - Drug Research* 73 (4), 851–854.
- Macedo, R.M., Brentegani, L.G., de Lacerda, S.A., 2015. Effects of coffee intake and intraperitoneal caffeine on bone repair process – A histologic and histometric study. *Braz. Dent. J.* 26 (2), 175–180. <https://doi.org/10.1590/0103-6440201300219>.
- Magnani, C., Galdorfini Chiari, B., Lucia Borges Isaac, V., Correa, M. A., & Nunes Salgado, H. R. 2014. In Vitro Safety Evaluation of Caffeic Acid. *Athens J. Health*, 1(3), 181–188. <https://doi.org/10.30958/ajh.1-3-2>.
- Martin, M., Sari, D. S., Praharani, D., Ayu, M. roslan. 2022. Combination of Dental Pulp Stem-Cell Secretome and Robusta Coffee Bean Extract (*Coffea canephora*) in Enhancing Osteocalcin and Alkaline Phosphatase Expression in Periodontitis-Induced Wistar Rats. *J. Orofacial Sci.* 13(2), 136–141. <https://doi.org/10.4103/jofs.jofs>.
- Nurman, S., Yulia, R., Irmayanti, Noor, E., Sunarti, T.C., 2019. The optimization of gel preparations using the active compounds of arabica coffee ground nanoparticles. *Sci. Pharm.* 87 (4). <https://doi.org/10.3390/scipharm87040032>.
- Ojeh, N., Stojadinovic, O., Pastar, I., Sawaya, A., Yin, N., Tomic-Caneh, M., 2016. The effects of caffeine on wound healing. *Int. Wound J.* 13 (5), 605–613. <https://doi.org/10.1111/iwj.12327>.
- Panduawala, C.P., Zhan, X., Dissanayaka, W.L., Samaranyake, L. P., Jin, L., Zhang, C., 2017. In vivo periodontal tissue regeneration by periodontal ligament stem cells and endothelial cells in three-dimensional cell sheet constructs. *J. Periodontal Res.* 52 (3), 408–418. <https://doi.org/10.1111/jre.12405>.
- Park, S. Min, Kim, D.H., Pang, E.K., 2017. Bone formation of demineralized human dentin block graft with different demineralization time: In vitro and in vivo study. *J. Cranio-Maxillofacial Surgery.* <https://doi.org/10.1016/j.jcms.2017.03.007>.
- Rurenga, P., Raangs, E., Singadji, Z., Wekema-Mulder, G., Veloo, A. C.M., van Winkelhoff, A.J., 2013. Evaluation of three selective media for isolation of Aggregatibacter actinomycetemcomitans. *J. Periodontal Res.* 48 (5), 549–552. <https://doi.org/10.1111/jre.12037>.
- Sari, D.S., Maduratna, E., Ferdiansyah, Latief, F.D.E., Satuman, Nugraha, A.P., Sudiana, K., Rantam, F.A., 2019. Osteogenic Differentiation and Biocompatibility of Bovine Teeth Scaffold with Rat Adipose-derived Mesenchymal Stem Cells. *Eur. J. Dentis.* 13 (2), 206–212. <https://doi.org/10.1055/s-0039-1694305>.
- Setiawatie, E.M., Gani, M.A., Rahayu, R.P., Ulfah, N., Kurnia, S., Augustina, E.F., Sari, D.S., 2022. Nigella sativa toothpaste promotes anti-inflammatory and anti-destructive effects in a rat model of periodontitis. *Arch. Oral Biol.* 137, (March). <https://doi.org/10.1016/j.archoralbio.2022.105396>.
- Shewale, A.H., Gattani, D.R., Bhatia, N., Mahajan, R., Saravanan, S. P., 2016. Prevalence of periodontal disease in the general population of India-A systematic review. *J. Clin. Diagnostic Res.* 10 (6), ZE04–ZE09. <https://doi.org/10.7860/JCDR/2016/17958.7962>.
- Shushtari, N., Froushani, S.M.A., 2017. Caffeine augments the instruction of anti-inflammatory macrophages by the conditioned medium of mesenchymal stem cells. *Cell J.* 19 (3), 415–424. <https://doi.org/10.22074/cellj.2017.4364>.
- Struppek, J., Walther, C., Bunte, K., Zyriax, B.C., Wenzel, J.P., Senftinger, J., Nikorowitsch, J., Heydecke, G., Seedorf, U., Beikler, T., Borof, K., Mayer, C., Aarabi, G., 2022. The association between coffee consumption and periodontitis: a cross-sectional study of a northern German population. *Clin. Oral Invest.* 26 (3), 2421–2427. <https://doi.org/10.1007/s00784-021-04208-9>.
- Sugihartini, N., Saridewi, R., M, U.R., Rahmawanti, F., Yuliani, S., Sophia, V. 2017. Anti-inflammatory Activity of *Camellia sinensis*, I. Extract Cream Combined with Vitamin C as Antioxidant on Croton Oil-induced Inflammation in Male Mice Strain BALB/C. *Majalah Obat Tradisional*, 22(2), 73. <https://doi.org/10.22146/trad-medj.27915>.
- Sukohar, A., Setiawan, Wirakusumah, F., Sastramihardja, H., 2011. Isolation and characterization cytotoxic compounds caffeine and chlorogenic acid isolasi dan karakterisasi. *Jurnal Medika Planta* 1 (4), 11.
- Tian, L., Su, C.P., Wang, Q., Wu, F.J., Bai, R., Zhang, H.M., Liu, J. Y., Lu, W.J., Wang, W., Lan, F., Guo, S.Z., 2019. Chlorogenic

- acid: A potent molecule that protects cardiomyocytes from TNF- α -induced injury via inhibiting NF- κ B and JNK signals. *J. Cell Mol. Med.* 23 (7), 4666–4678. <https://doi.org/10.1111/jcmm.14351>.
- Wahyukundari, M.A., Sari, D.S., Pujiastuti, P., Praharani, D., Arina, Y.M.D., Ermawati, T., Latief, F.D.E. 2017. Virulence of periodontopathogens *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*: micro-computed tomography and inflammatory cytokine analysis. In: 11th International Dentistry Scientific Meeting (IDSM 2017), 4, 312–321. <https://doi.org/10.2991/idsm-17.2018.42>.
- Yamagata, K., Izawa, Y., Onodera, D., Tagami, M., 2018. Chlorogenic acid regulates apoptosis and stem cell marker-related gene expression in A549 human lung cancer cells. *Mol. Cell. Biochem.* 441 (1–2), 9–19. <https://doi.org/10.1007/s11010-017-3171-1>.



Title	Type	↓ SJR	H index	Total Docs. (2022)	Total Docs. (3years)	Total Refs. (2022)	Total Cites (3years)	Citable Docs. (3years)	Cites / Doc. (2years)	Ref. / Doc. (2022)	
69 Saudi Dental Journal 	journal	0.446 	33	100	300	3395	627	295	1.92	33.95	