













NATIONAL CONGRESS OF THE INDONESIAN SOLOGY SOCIETY INTERNATIONAL WORKSHOP, SYMPOSIUM AND SEMINAR PHYSIOLOGY 2018)

13-30, 2018, Bandung, Indonesia

PROGRAMME BOOK (ABSTRACT)





Raya Bandung-Sumedang Km. 21, Sumedang 45363

1022) 84288828, Email: iaifibandung@gmail.com

TABLE OF CONTENTS

	1
LREACE	2
2. RUNDOWN	6
IL ORGANIZING COMMITTEE	8
4. WAP OF PLACE	9
E. E.E.T DESCRIPTION	17
SPEACERS ABSTRACT	10
ABSTRACT	ه بله









The 17th National Congress of The Indonesian Physiology Society and 27th International Physiology Seminar 2018 (BIPSC 2018)

Bandung, 28th -30th of June 2018

Preface

Dear colleagues,

Welcome to the 17th National Congress of Indonesian Physiology Society and the 27th International Physiology Seminar (BIPSC) in Bandung, West Java, Indonesia. BIPSC 2018 is an annual meeting of the Indonesian Physiology Society. This year we welcome all researchers and lecturers to share an interesting report in all aspects of Physiological Sciences, including advanced research and translational physiology including in human, plant and animal study area. The theme for the congress is "Harmony of life: Integrated Physiology for one health". We believe that harmony of animal health, environment (plant health) and human health as one triad health will be a key point to improving human quality of life.

We had received more than 100 abstracts and more than 150 registrations from Indonesia and foreign country especially Asian and Oceanian countries. Guest speakers are coming from many renowned universities in London, Japan, Singapore, Malaysia, Australia, and also from Indonesia itself. They will share their knowledge and experiences on related topics during the event. The congress will also be associated with Teaching Workshops and Training Courses for young researchers. There will be numerous opportunities to gain new knowledge, share technical expertise and the latest research discoveries, and to enjoy scientific and educational interactions.

Through the workshop, symposium and seminar, the Indonesian Physiological Society aims to help physiologist, exercise physiologist, specialist, general practitioner, coaches and other related professionals by sharing the latest science and technology to prevent Non-Communicable Diseases (NCD) and Improve Athlete Performance.

Finally, the output from this event will be reviewed and the selected full manuscript will be published in national or international indexed Scopus partner journal.

Whether you are attending the workshop, symposium or seminar, we wish you a successful event and hope you will enjoy your visit and leave with many happy memories from BIPSC2018.

Chairman of BIPSC 2018,

BIPSC 20

Dr Vita Murniati Tarawan, dr., Sp.OG., M.Kes., AIFO., SH.

Norkshop 1		•	Speaker Askes PhD		
/enue Eijkman Building	09.00- 16.00	Topic Western Blot	Ronny Lesmana, dr.,MKes.,PhD Andri Rezano, dr., M.Kes, PhD M.Rizky Anggun dr.,Mkes.,PhD Susianti		
Workshop	2		Speaker		
Venue Time Topic Grand 09.00- Exercise Physiology I: Exercise Physi		Exercise Physiology I: Exercise endocrinology and Doping assessment	prise Prof. Dr Ieva B. Akbar.,dr.,AIF Dr.Vita M.Tarawan, dr.,SpOG.,Mkes.,SH.,AIF dr.,MS.,AIF		
	13.00- 16.00		Prof Beltasar Tarigan Infandiary Ihrahim, MS (UI)		
Worksho	р3		Speaker		
Venue Eijkman Building	09.00 16.00		dr Edward Jayahadi., M.sc		
Worksho	op 4		Speaker		
Venue Grand Tjokro	7ime 09.0 16.0	0- Scientific Writing	Ronald Hamidie.,dr.,PhD Kasno Pamungkas SS.,MHum Ahmad Faried.,dr.,SpBS.,PhD Rizky Abdullah, Apt., PhD Dr. Dra. Triadiati M.Si		
Marks	hon 5		Speaker		
Workshop 5 Venue Grand Tjokro		Teaching Physiology I: Cr Organ Principles and Ir Organ Physiology	Prof. Cheng Hwee Ming		
	18	Organ Physiology .30- Physiology Experiments 8.00- Welcome Dinner	. I-I Mar ())IIIOga ()		
1		1.00	E na		

Day 2	(Friday,	29th	June	2018	١
-------	----------	------	------	------	---

Venue	Time	Activity	Topics	Speaker		
Grand	07.30-	Registration			- Contract of the state of the	
Tjokro	08.30					
Ballro	08.30-	Opening Ceremony	Sundanese T	raditional Dance		
om	09.00		Indonesia Ra	aya		
			Hymne IAIFI			
			Sambutan Ke	Sambutan Ketua Panitia		
			Sambutan PP IAIFI			
			Pembukaan oleh Rektor UNPAD			
	09.00-	Plenary Lecture		Menkes Indonesia (atau yang mewakili)		
	10.00	4 4		Prof Tri Hanggono Achmad (Rektor		
				UNPAD)	(,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
	10.00-	Keynote Speaker: Ph	avsiology for		MSc.Agr.,	
	11.30	One Health	lysiciogy ioi	AIFH (IPB)		
	33.150			Prof.Dr. A. Purba, dr., AIF (Unpad)		
				Dylan Morrissey (Queen Mary Univers		
		40		London)	ry Omversity	
	11.45-	LUNCH + Break - Frida	av Prav	Londony	+	
	13.00	Editor Filedical Trial	ayıtay			
Baralle		a Ruang I				
Venue	Time	Theme	Topics	Speaker	Moderator	
A	111111	THETHE	TOPICS	Speaker	Wioderator	
	13.00-	Symposia I:	Physiology	Prof. Dr. leva Baniasih	Astrid	
	14.30	Physiological	of Aging	Akbar, dr.,Mkes.,AlF	Feinisa	
	Titian	updates on Aging,	Of Aging	(Unisba)	remisa	
		Metabolic and		Paul M Yen (Duke NUS)		
		Endocrine		Paul W Fell (Duke NOS)		
		endocrine	Cargononia	Laguardi de CaDD		
			Sarcopenia	Lazuardi, dr, SpPD		
			Obesity:			
	-		New			
			Updates			
	3.0 3.0				Nova	
	14.30-	Symposia II: Update	s on Cardiac	Dr. Rizky M.Akbar, dr.,SpJP		
	18.50	Symposia II: Update Physiology	s on Cardiac	(Unpad)	Sylviana	
	Section 1		s on Cardiac	(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	
	10.00	Physiology		(Unpad)	Sylviana	
	10.00			(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	
	18.00 216.00 216.00	Physiology Caffee Break and Pos		(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	
	18.00 18.00 18.00 18.00	Physiology		(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	
	18.00 246.00 246.00 18.00	Cral Presentation		(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	
	18.00 18.00 18.00 18.00	Physiology Caffee Break and Pos		(Unpad) Badai Tiksnadi, dr., SpJP	Sylviana	

Digital Repository University 3 (Saturday, 30th June 2018)

arallel S	Symposia	Ruang II		Moderator
	Time	Theme	Speaker	
3	13.00- 14.30	Symposia III: Animal Physiology	Dr. Endang Yuni Setyowati, drh.,MSc.Ag (Unpad) Prof Dr. Roostita B Prof.Ir.Wasmen Menalu PhD (IPB)	Pranyata Tanggul Waskita, dr (Unpad)
	14.30- 16.00	Symposia IV: Neurophysiology and Sport Biomechanics	Prof.Dr.dr.I Putu Gde Adiatmika, Mkes (Unud) Hanna, dr., Mkes.,PhD (Unpad) Siti Baitul Mukaromah	dr. Renaldi Praseti Mkes.,SpOT (Unpad)
	16.00-	Coffee Break and Po	ster Session	
	16.30- 18.00	Oral Presentation		
Paralle	Sympos	a Ruang III		Moderator
Venue		Theme	Speaker	Titing Nurhayati
Venue C	-	Symposium V Exercise Physiology	Lagnardo	WA
	14.30- 16.00	Exercise Physiolog for Health an Athletes	: Prof Irawan Yusuf (Unhas) Dr.dr.Ermita Isfandiary Ibrahim of MS (UI) Prof Gusbakti Reinaldi (UISU) d	Yosef Purwoko
	16.00	Coffee Break and		
	16.30	- Oral Presentation		

Venue	Time	ia Ruang I Theme	I a d	
A	ime	Theme	Speaker	Moderator
	08.00- 10.00	Symposia VII: Aging, Metabolic and Endocrine Physiology	Noriyuki Koibuchi (Gunma University) Annisa Widjaya (DUKE NUS) Prof. Naguib Saleh MD (UM)	Prof.Lovita Adriani.,Ir.,MS
	10.00-	Symposia VIII: Molecular Physiology	Dr.med.Setiawan, dr., AIF (Unpad) Winifred PhD (DUKE NUS)	Nur Atik
	Sympos	ia Ruang II		
Venue B	Time	Theme	Speaker	Moderator
	08.00- 10.00	Symposia IX: Plant Physiology	Dr. Dra. Triadiati M.Si (IPB) Noladhi Wicaksana, SP.,MP.,PhD (Unpad) Dr. rer nat Soeseno Amien	Kadapi
r,	10.00- 11.00	Symposia X: Psychophysiology	Prof Cheng Hwee Ming Aulia	Yuni Susanti
Paraffel	Sympos	ia Ruang III		
Venue	Time	Theme	Speaker	Moderator
	08.00- 10.00	Symposia XI: Physiological Prespective of Doping	Ronny Lesmana, dr.,MKes.,PhD (Unpad) Dr.Vita M.Tarawan, dr.,SpOG.,Mkes.,SH.,AIF (Unpad) Dr. Gaga Irawan, dr., Mkes (Unpad)	Titing Nurhayati
	10.00	Symposia XII: Pharmacophysiology	Keri Lestari, PhD (Unpad) Rizky Abdulah, PhD (Unpad) Melisa PhD (Unpad)	Dr. Irma Melyan Puspitasari
11 000	Oral Pri	sentation		
T m	Lunch B	reak		
1 (16)	Deal Pre	mentation		
101	kalling I	break A Poster Session		
	Elitting	Ramarki & Award		

SUSUNAN KEPANITIAAN

KONGRES NASIONAL IAIFI KE-XVII DAN 27th INTERNATIONAL WORKSHOP, SYMPOSIUM SEMINAR PHYSIOLOGY, BANDUNG, 28-30 JUNI 2018

Penasehat	:	Rek	tor Universitas Padjadjaran
	1:	Dr.	med. Setiawan, dr., M.Kes., AIFM (Dekan FK-
Pengarah:		LINI	PAD, Bandung)
		Pro	f. Dr. drh. Agik Suprayogi, MSc, AIF
		Pro	f. Adnyana Manuaba Hon.FErg.S., FIPS, SF
	1	Pro	of. Dr. H.R. Soedarso Djojonegoro, dr., AIF
		Pro	of. Dr. dr. A Purba, MSc, AIFO
		Pro	of, Dr. Wahyu Kahiwikarta, dr., MS.
		Pre	of. Dr. Ir. H. Mochamad Bintoro, M. Agr
		Pr	of. Dr. Beltasar Tarigan, MS, AIFO
	1	Pr	of, Dr. leva B. Akbar, dr., AIF.
	1	Pr	of. Dr. dr. I Ketut Tirtayasa
	1	Pr	of. Dr. Sayuti Syahara, MS., AIFO
		Pr	of. Dr. drh. Pudji Astuti, M.P
Ketua		: D	r. Vita Murniati Tarawan, dr., M.Kes., SpOG., AIFO,
Ketua		S	- Ph D
Sekretaris Umum		: R	onny Lesmana, dr., M.Kes., Ph.D.
Sekretaris I		: Y	uni Susanti Pratiwi, dr., M.Kes., AIFO
Sekretaris II		: 1	yagita, drh., MVSc
Staf Kesekretariatan		: 4	zziiz M. Rosdianto, Skep., Ns., MH.Kes., MSi.
Star Research			Aep Kusnandar, SSi.
		1	Susianti, SSi.
			dr. Naufal M. Nurdin. S.ked. M.Si
Bendahara Umum			Hanna, dr., M.Kes., PhD, AIFO
Bendahara			Julia Windi Gunadi, dr., M.Kes
			Titing Nurhayati, dr., M.Kes
Bidang Ilmiah		1:1	Ronald Hamidie, dr.,PhD
Koordinator		11	Lucky Angkawidjaya, dr., M.Pd
Sekretaris		1:1	Dr. Caecilia, drg.
Anggota			Dr. Sri Tjahajawati
			Dr. Reni Farenia,dr., M. Kes
		1:	DI REMIT WILLIAM
Bidang Workshop		1:	Hanna, dr., MKes., PhD
Koordinator		+	Stella Tinia dr, M.Kes
Sekretaris		+	Susianti, S.Si.
Anggota		+	
Bidang Sponsor dan Pameran		+	Dr. M. Rizki Akbar, dr., Sp.JP(K), M.Kes
Koordinator		+	- Luca-doop dr Sn.PD M.Kes.
Sekretaris			- Desuid dr SnOT
Anggota			Widayanti, dr., M.Kes.
		1	Apen, dr., SpPD

Bidang Program dan Protokoler		
(welcome dinner)		
Ketua	;	Nova Sylviana, dr., M.Kes.
Sekretaris	;	Decky dr., M.Kes
		Naufal M. Nurdin, dr., S.Ked., M.Si
		Dr. Triadiati, MSi, AIFT
		Dr. Drh. Aryani Sismin Satijaningtijas, MSc.
Bidang Sosial (Hospitality dan Ladies	;	
Program)		
Ketua	:	Kartika Indah Sari, drg., M.Kes
Sekretaris	:	Anggun, drg
Bidang Publikasi dan Dokumentasi	:	
Koordinator	:	Harijadi, dr.
Sekretaris	:	Decky, dr., M. Kes
Bidang Konsumsi	-	
Koordinator	:	Juliati, dr., AIF
Sekretaris	:	Yuni Susanti Pratiwi, dr., M.Kes., AIFO
Anggota	:	Eka, dr.
Bidang Logisitik	:	
Koordinator	:	Ronny, dr., M.Kes., PhD
Sekretaris	:	Fathul Huda, dr., PhD
Anggota	:	Dr. Leonardo Lubis, dr., M.Kes., AIFO
		Aziz MR
		Masri, S.Si
		Edi Sukmayadi
		Wanto

Tanggal

: 15 Februari 2018

Ketua Umum PP IAIFI

Prof. Dr. drh. Agik Suprayogi, M.Sc, AlF

Dislipidemia Prevention Caused by The Administration of Coconut Dregs Extra

The Effect of Monosodium Glutamate & High-Fat Diets on Ovarian and
Fulloplan Tube Histonathological Changes of Battus and Part Diets on Ovarian and in Rats Given A High-Fat Diet

Aris Prasetyo¹, Harjanto², Achmad Subagio³

- 1. Physiology Departement Medical Faculty University of Jember 2. Professor Physiology Departement Medical Faculty Airlangga University Surah
- 3. Professor Chemistry Departement Agrocultural Technology Faculty University of Jemb

Consumption of coconut dregs extract/ CDE were reported to have much benefit in preven dyslipidemia profile, such as total cholesterol, triglycerides, LDL cholesterol and HDL cholesterol CDE containing galactomannan is nondigestible carbohydrate in intestinum so fermentated in C and enhancing SCFA production (acetat, propionate, and butyrate). GLP-1 may serve as a poliprevention agent for dislipidemic. This study investigated the effect CDE prevent increasing of cholesterol, triglycerides, LDL cholesterol and prevent decreasing HDL cholesterol in male rule high fat diet. Wistar male rats were 3-4 months age and determination of the object of research completely randomized design with four treatments, namely (1) rats with a diet high in fat. with high-fat diet + CDE 70 mg/200gw, (3) rats with high-fat diet + CDE 140 mg/200gw, with high-fat diet + Simvastatin 0,18mg/200gw. Feeding rats administered orally, whereas sonde 1x/day. Plasma lipid profile was measured by by spektrophotometry. The results suggests CDE prevent significantly increased total cholesterol, trigliseride, LDL cholesterol in 140mg <0,05) and prevent significantly decreased HDL cholesterol (p <0,05). The administration is contributed to preventing the increase in LDL cholesterol and the decrease in HDL cholesterol

Keywords: CDE, Galactomannan, SCFA, GLP-1, high fat diet, dyslipidemia

Hantuti RD¹, Ariani NKD¹, Mustika D², Kurnianingsih N², Agustin D³, Indriani A⁴

- Undergraduate of Midwifery Program, Faculty of Medicine, Brawijaya University, Indonesia. Department of Physiology, Faculty of Medicine, Brawijaya University, Indonesia.
 - Department of Anatomy and Histhology, Faculty of Medicine, Brawijaya University, Department of Midwifery, Faculty of Medicine, Brawijaya University, Indonesia.
- prevalence of obesity and obesity-associated complications continues to be a major that the difference it is well accepted that both excessive consumption of monosodium glutamate high-fat diets lead to obesity and disorder of reproductive organ. However, there are and that showed the effect of daily consumption dose of MSG combined high-fat diets on The urgan. Therefore, this research focused at the effects high-fat diets combined with MSG, the the average daily consumption of humans, on ovarian and fallopian tube histopathological
 - winter rate were selected and divided into 6 groups, consists of control, high-fat diet high doze MSG (0.7mg/gBW) alone, combined HFD and MSG (0.05; 0.2; MSG was administrated once a day for 56 days and The rats were euthanized and the ovarium and fallopian tube 47.61 when reached the proestrus phase. Histopathologic preparation with HE the calculating the number of follicles de Graaf and atretic follicles of ovarium standard cells and smooth muscle of fallopian tubes by using a light microscope. Way ANOVA method followed by Post hoc Tukey test. The result showed partite beginning antly decreased the secretory epithelial cells number (p = 0.008) this have of fallopian tubes (p = 0.001) compared to control. It also significantly whereas insignificantly decreased Graafian Nevertheless, there were no significant differences
 - 1111 D and 111 D alone, accept for the number of atretic follicles. In this study, the same effects with high and tallopian tube histopathological changes. High-fat diets combined daily consumption of humans, have the same effect as the effects block on ovarian and fallopian tube histopathological changes of

D-4

The effect of Ambon Banana stems sap (Musa paradisiaca L var sapientum) as an anti-angiogenesis towards HUVEC induced by IL1 and CdCl2

Sofiana KD¹, Wulan PMY², Alinda R³, Dorothy MJ³, Prihardina B⁴, Permatasari N⁵, Husnul K⁵, Widodo MA⁵

¹ Physiology Department; Universitas Jember; Jember; Jawa Timur; Indonesia ,Electronic

² Biomedical Sciences; Universitas Brawijaya; Malang; Jawa Timur; Indonesia

³ Medical Faculty; Universitas Brawijaya; Malang; Jawa Timur; Indonesia

⁴ Biomedical Sciences Laboratory; Universitas Brawijaya; Malang; Jawa Timur; Indonesia

5 Pharmacology Department; Universitas Brawijaya; Malang; Jawa Timur; Indonesia,

Tumors are characterized by an increase in angiogenesis that is identical to cell growth, invasion, and metastasis. Contamination of heavy metal such as Cadmium causes cell transformed into malignancy VEGF levels and migration are markers of angiogenesis process. Ambon banana is a plant that became local wisdom of the Indonesian nation which is used as a treatment therapy of various diseases. The antioxidant contains inside this banana stem sap according to some studies are useful to inhibit the proliferative process or excessive migration in cases of tumor or cancer. The aim of this research to look at the anti-angiogenesis effect through migration and VEGF levels on HUVECS which induced by IL1 and CdCl2. About 70-80% confluent HUVECs were divided into five groups The control group was HUVEC were left within 12 hours then scratched and incubated in 24 hours Induction group was induced by IL1 (200ng / ml) in 12 hours then scratched and was given CdCl2 (24,154 µg / L) in medium and incubated in 24 hours. The treatment group was HUVEC induced by IL-1 (200 ng/ml) for 12 hours scratched and was given CdCl2 (24,154 µg/L) along with various concentrations of Ambon banana sap (0.125%, 0.25%, 0.5%) then incubated for 24 hours. Migration measurements were performed by scratch assay at 12 and 24 hours and the measurement of VEGF levels on culture medium was done at 12 and 24 using elisa method Data were analyzed using SPSS software. Ambon banana stem sap inhibit excessive migration on II 1 and CdCl2 induced HUVECs on certain time and concentration. Administration of Ambon banana stem sap is also able to downregulate VEGF levels at the certain time and concentration. Ambon banana stem sap can suppress excessive angiogenesis process due to induction of IL1 and CdC12 in in-vitro study on HUVEC.

Keyword: HUVEC, CdCl2, Ambon banana stem sap, migration, VEGF

D-5

The Effect of Glutathione to a Change of SGOT, SGPT, and ALP White Rated High Diets Diet

Eric Mayo Dagradi

Department of Physiology, Hang Tuah Medical School Surabaya

Hard work without rest will ultimately burden the heart. These conditions can damage cells including liver cells, resulting in a fourfold increase in aspartate plasma transaminase (AST / SGOT) and elevated bilirubin levels which are a sign of impaired liver function (Droge, 2002). Exercise done for a moment, also can increase AST / SGOT and Alanin amino transaminase (ALT / SGPT) in the blood (KoutLdakis, 1993). Liver damage is reversible as a result of accumulation of fat accumulation in liver cells from abnormal metabolism. Causes of fatty liver (FDL) include alcohol abuse, protein malnutrition, diabetes mellitus, obesity, hepatotoxins and drugs. Antioxidants are defined as compounds that can delay, slow down, and prevent lipid oxidation processes. In a special sense, antioxidants are substances that can delay or prevent the occurrence of free radical antioxidation reactions in lipid oxidation. High levels of free radicals such as heavy physical activity, resulting in the ability of endogenous antioxidants are not sufficient to neutralize free radicals resulting in an unbalanced state between free radicals and antioxidants called oxidative stress Glutathione is one of the endogenous thiol compounds that act as defense of anti-oxidants in the cells so it can also act as a powerful anti-oxidant. And one way is the use of anti-oxidants endogenous that should be produced in the body, but added in certain amounts to be exogenous antioxidant that is glutathione. (Ashtiani et al., 2011). Glutathione protects cells from damage caused by oxygen containing unstable molecules, which are a by-product of energy production (Kerksick C and Willoughby D. 2005). Proving Glutathione can improve SGOT, SGPT and ALP due to high-fat diet induction. Wistar strain male, 20 weeks old, weighing 180 - 200 grams, divided into 4 groups, each group of 8, with Separate pre-post test control groups design. The treatment group performed one intravenous adrenaline injection and was exposed to egg yolks as a high-fat diet. Moderate treatment controls without adrenaline injection and standard feed exposure were all performed within 14 days. Glutathione was administered intraperitoneally at a dose of 600 mg / kilogram BB and 1200 mg / KgBB, once daily for 7 days. Data was taken on day 14 by means of cardiac punture. Blood analysis unit taken from the heart is done by measurement of SGOT and SGPT enzyme and ALP by using photometer. Giving Glutathione due to the induction of a high-fat diet with a short time can not decrease the SGPT level is evidenced by anova assay that is greater than alpha value, the difference in decreased levels of SGOT on LSD parametric test, and decreased ALP levels due to high-fat diet induction is evidenced by significant P value 0.001 <5% in the non parametric test of Mann Whitney. Short-term glutathione with high-fat diet induction can not lower levels of SGPT but may lower levels of SGOT and ALP

Keywords: Glutathione, oxidative stress, fatty liver.

C-18

Assessment of Heavy Metal Pollution in Lung and Kidney of Broiler, and It's Correlation of Water and Animals Feed

Amelia RA1, Hera M2, Huda SD3

- Department of Anatomy, Physiology and Pharmacology, Bogor Agricultural University, Bogor, West Java, Indonesia.
- 2. Divion of Physiology, Department of Anatomy, Physiology and Pharmacology, Bogor Agricultural University, Bogor, West Java, Indonesia.
- 3. Divion of Pharmacology and Toxicology, Department of Anatomy, Physiology and Pharmacology, Bogor Agricultural University, Bogor, West Java, Indonesia.

Cadmium belongs to heavy metal with high anthropogenic pollutant and non-essential to physiological function. Cadmium present in soil, water, air as well as in food. Reguler intake of consuming cadmium leads to physiological disorders in poultry animal as well as humans as one who consume it. The aim of this study was to analyze the profile of cadmium content in environtmental samples (poultry feed and water) and it correlated with the degree of contamination in broiler tissue The study used 30 sample of poultry feeds and animal water, 30 sample of chicken lungs and 30 sample of chicken kidneys. Feed samples were taken at each floc where the chickne were previously taken. Testing and determining levels of heavy metal cadmium is referred to the Indonesian National Standard (SNI) 01-2354.5-2011 by using AAS (Atomic Absorption Spectrophotometry). The results showed level of cadmium in feed between 0.009-0.202 mg/kg and 0.0068-0.0096 mg/l of cadmium in the water. Cadmium in the lungs and kidneys was positively contaminated by cadmium with range of 0.438-0.655 mg / kg and it was significantly different (p < 0.05) among the organs. A strong positive correlation was found between cadmium and water levels in renal (r = 0.925). In contrast, weak positive correlation (r = 0.624) was found in the relationship between cadmium in feed and lung. The environmental quality of cadmium contamination should be aware for its effects to physigological status of animals as well as humans as the highest food chain. It is indicated by the relationship between cadmium in feed and water deposited in chicken tissues. Cadmium concentration in the poultry feed and water increased significantly to the concentration of cadmium deposited in organs.

Keyword: cadmium, water, pollutant, poultry, animal

C-19

The Role of Resistant Starch Type 3 Modified Cassava Flour (MOCAF) as an Alternative Prebiotic in Diabetes Mellitus Rats Model

Jauhar Firdaus ¹, Elly Nurus Sakinah ²

- 1. Departemen of Physiology, Faculty of Medicine, University of Jember
- 2. Departement of Pharmachology, Faculty of Medicine, University of Jember

Prebiotics has widely known to have benefit in human health. As an alternative prebiotics Resistant Starch type 3 (RS3) has has several advantages wich are do not cause constipation nor diarrhea, RS3 also decrease blood glucose and increase the level of Glucagon Like Peptide-1 (GLP-1) which has a major role in the mechanism of blood glucose control in patients with diabetes mellitus (DM). One that can be used as a source of RS3 is Modified Cassava Flour (MOCAF). Objective: The goal of this study is to analyze the role of RS3 MOCAF as an alternative probiotic in diabetes mellitus rats model. This study used twenty four rats randomly selected and grouped into 4 group (normal, MOCAF, RS3 MOCAF and Diabetes) given different food (standard, MOCAF and RS3 MOCAF) as much as 20 gram each day for 4 weeks. In the end of study blood taken to measure post prandial blood glucose and fasting blood glucose. Stool sample were taken from large intestine for Short Chain Fatty Acid (SCFA) and microbiological analysis. Dietary RS3 MOCAF decrease post-prandial blood glucose from 526 mg/dL to 96 mg/dL and fasting blood glucose from 494 mg/dL to 107 mg/dL. SCFA analysis showed that fermentation of RS3 MOCAF produces acetic acid (16:18 mmol), propionic acid (5:51 mmol) valeric acid (2:52 mmol) and butyric acid (0.57 mmol). Microbiological analysis showed that the most widely grown bacteria in the large intestine after dietary RS3 mocaf is Lactobacillus Sp. RS3 MOCAF plays role as prebiotics and has ability to decrease blood glucose in diabetes mellitus rats model.

Keyword: Prebiotics, Resistant starch type 3, MOCAF, Blood glucose, SCFA, Diabetes mellitus

D-16

Influence of Left Ventricular Pump Dysfunction on Estimated VO2max in Patients after ST-Segmen Elevation Myocardial Infarction.

P. Wulandari

Department of Physiology, Faculty of Medicine, University of Jember

Left ventricular (LV) remodeling and dysfunction started Early after ST-Segmen Elevation myocardial Infarction especially in a large Myocardial Infarction (MI). A model of the relationships among the concepts of level of left ventricular pump dysfunction is ability to perform exercise without an ischemic response and achievement of an exercise trained state and functional capacity. The 6 MWT as a tool to evaluate the estimated VO2max is a safe and reproducible measurement of functional capacity in stable patients after a noncomplicated MI, even when performed within a week of the event The greater the level of left ventricular pump dysfunction as measured by ejection fraction following an MI, the less the heart is able to pump blood at rest and during exercise. The present study investigated how LV pump dysfunction influence the patient after ST-Segmen Elevation Myocardial Infarction to perform exercise. This is a cross sectional study of 42 consecutive patients with STEMI Eligible patients measurement of Left Ventricular ejection fraction (LV EF) using echocardiography within 48 hour of admission and 6MWT pre-discharge or at the first follow-up visits (day 4-9) Conversion 6 MWT distance in estimated VO2max using Cahalin formula. The correlation of LVEF and VO2max was analyzed through Pearson's correlation test. There was a weak correlation between echocardiographic LVEF and VO2max (r=0.273, p=0.081). The mean of six-minute walk distance among this STEMI patiets was 368.36 (±58.90), mean VO2max was 10.56 mL/(kg.minutes) (±1.18) and mean of LVEF was 45.29% (±8.256). This weak correlation between LVEF and VO2max in patients after STEMI explain why post-MI patients differ in their response to exercise

D-17

Effect of The Problem-Focused Coping Stress Management Program on Self Efficacy of Students Semester I University of Udayana

Susy Purnawati, I P G Adiatmika, I Made Krisna Dinata

Department of Physiology Faculty of Medicine, Udayana University

The academic tasks or work load of medical students is heavy in quantitative or qualitatively. This is in accordance with the demands of the profession and their responsibilities which will have to deal with the risk of patient death and medical misconduct. The academic load factor combines with individual factors and another related factors resulting of academic stress. Academic stress must be manage early, as it may lead to a tendency of decreased academic performance and an increased risk of mental disorders. The problem-focused coping stress management program proved to be effective in improving self-efficacy in some professions. High score of self-efficacy indicate with reduces psychological distress in a reason that the person will see the demands of the task as a change to achieve the goal. Problem-focused coping stress management program can include personal skill component training and time management skills training. This experimental study aims to analyze the improvement of student self-efficacy after undergoing a problem-focused coping stress management program. From 40 first semester students of Faculty of Medicine Udayana University selected by a simple random sampling and have high academic stress score determined proportionally and grouped as control group (14 people) and treatment group (26 people), and selected by random allocation with simple random sampling technique using lottery method. This study conducted at Denpasar in April to December 2015. The intervention was in the form of problem-focused coping stress management training which was done 4 times each week with the duration of each session was 1 1/2 hours in each classroom training session. In the subject matter of study subjects was introduced with the impact of academic stress, the pathophysiology of academic stress on the tendency of physical or mental illness, and the benefits of academic stress management program of problem-focused coping methods. Participants was also trained in practicing time management techniques, and to be internal locus of control type to improve cognitive performance. Subjects are also given independent tasks that are done in the form of each home work and learn to fill the mood diary. The result of this study showed that the self-efficacy was significantly different between before and after treatment in the intervention group with p = 0.005 (using statistic paired sample t-test), with mean of 19.31 ± 2.396 to 21.27 ± 2.677 . This shows that there is an increase of self-efficacy by 10%. There was no significant difference between baseline self-efficacy data (pre) and post in the control group. Different self-efficacy test after treatment between those two group using independent sample t-test also showed a significant difference, with the p value = 0.029 with mean and SD of self-efficacy (post) respectively were 19.29 ± 1.541 and 21.27 ± 2.677. It can be concluded that the effect of problem-focused coping stress management improves student self-efficacy by 10%.

Keywords: problem-focused coping management program, student self-efficacy