
ISBN: 978-602-60569-5-5





PROCEEDINGS

The 1st International Basic Science Conference 2016
TOWARDS THE EXTENDED USE OF BASIC SCIENCE
FOR ENHANCING HEALTH, ENVIRONMENT,
ENERGY, AND BIOTECHNOLOGY

University of Jember, September 26 - 27, 2016



The 1st International Basic Science Conference 2016

(The 1st IBSC 2016)

"Towards the extended use of basic science for enhancing health, environment, energy and biotechnology"

Editor Fuad Bahrul Ulum M. Ziaul Arif Agung Tjahjo Nugroho Yudi Aris Sulistiyo

Faculty of Mathematics and Natural Sciences

The University of Jember

Jember

Indonesia

Email : ibsc.fmipa@unej.ac.id

Website : http://ibsc.fmipa.unej.ac.id/

Address: Jl. Kalimantan no 37 Kampus Tegal Boto, Jember 68121

First Published 2017

TITLE : PROCEEDINGS OF THE 1st INTERNATIONAL BASIC SCIENCE

CONFERENCE 2016

EDITORS: Fuad Bahrul Ulum, M. Ziaul Arif, Agung Tjahjo Nugroho, Yudi Aris

Sulistiyo



ISBN: 978-602-60569-5-5

Copyright © 2017 IBSC The University of Jember and the authors of the paper

All Rights Reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any means without the permission in writing of the copyright holders.

Preface

A Conference on the extended use of Basic Science was hosted by the Faculty of Mathematics and Natural Science (FMIPA), at CDAST building, Universitas Jember in 26-27th day of September 2016. This conference is intended to promote further developments of basic science for their tangible applications, especially health, environment, energy and biotechnology.

The conference posed the question "what biological, chemical, physical, geological, mathematical, statistical, medical, agricultural and other basic science field changes must be made in order to ensure better live quality in term of health, environment, energy and biotechnology. FMIPA was fortunate to welcome researchers, educators and engineers from various backgrounds representing a variety ways to extend the application of basic science in which safety, environmental friendly and energy efficiency were being pursued. More than two hundred contributors from fifty five different institutions presented the theory, methodology and application of the field and thus the 1st IBSC 2016 was very rich as the proceeding in this volume.

The major theme that emerged from the conference which was conducted by Faculty of Mathematics and Natural Science, Universitas Jember, and The Ministry of Technology and Higher Education (KEMENRISTEKDIKTI), Republic of Indonesia, was that basic science must extend in very fundamental way if high live quality is to become a stable standard of health, environment, energy and biotechnology. Ramkrishna Ramaswamy in his very inspiring talk, present the complexity and simplicity in biological systems; while Agus Salim discussed the big data of biostatistics with stressing on the quantity does not equal quality; Manabu Abe report the design and synthesis of a new Cromophore, and Bambang Sugiharto address the regularisation of sucrose-phosphate synthesis from sugarcane, and many more expert discus the application of basic science for improving live quality.

Having introduced the 1st IBSC 2016, we will introduce the 2nd Conference of basic science (The 2nd IBSC) in near future. This collaboration and link will be maintain. Hand in hand researcher, expert, educator and other professional in basic science is needed to improve live quality.

Jember January 26, 2017

Agung Tjahjo Nugroho

The Chairman of The 1st IBSC

Reviewers

Drs. Sujito, Ph.D

Drs. Yuda Cahyoargo Hariadi, M.Sc., Ph.D

Dr. Rike Oktarianti, M.Si.

Purwatiningsih, S.Si, M.Si, Ph.D

Ika Oktavianawati, S.Si, M.Sc

Dwi Indarti, S.Si., M.Si

Yeni Maulidah Muflihah, S.Si, M.Si

Dr. Mohamat Fatekurohman, S.Si., M.Si

Erlia Narulita, S.Pd., M.Si., Ph.D




Table of Contents

ACE	
E OF CONTENTS	
OGY	1
COMMUNITY STRATEGY FOR MANAGING TROPICAL FOREST RESOURCES IN THE AREA OF CAGAR ALAM PULAU SEMPU (NATURE RESERVE OF SEMPU ISLAND)	2
BIOREDUCTION ADSORBENT (BIOSORBENT): RECOVERY TECHNOLOGY OF HEAVY METAL POLLUTION (CADMIUM/	
CD) IN POLLUTED LAPINDO WATER SOURCES USING BACTERIA AND DURIAN LEATHER	7
COMPARATIVE STUDY OF THE MANAGEMENT OF VANAME SHRIMP (LITOPENAEUS VANNAMEI) BASED ON	
DEMOGRAPHIC FACTORS AT MOLANG BEACH TULUNGAGUNG	10
ANALYSIS OF THE INFLUENCE OF PUBLIC PARTICIPATION IN THE MANAGEMENT OF RESOURCES SUSTAINABLE	
WATER MALANG DISTRICTCONSERVATION COCCINELLA SP. AS PREDATOR OF GREEN PEACH APHID MYZUS PERSICAE SULZER ON POTATO INTERCROPPING	
THE EFFECT OF MYCORRHIZAL INOCULANT AND COMPOST OF VOLCANIC ASH ON GROWTH AND YIELD OF CHILLI (CAPSICUM ANNUM L.)	
THE POTENTIAL OF ARTHROPODE DIVERSITY FOR ECOTOURISM DEVELOPMENT IN WONOREJO MANGROVE ECOSYSTEM, SURABAYA	
THE EFFECTS OF WATER FRACTION OF BITTER MELON (MOMORDICA CHARANTIA) LEAF EXTRACT IN MAMMARY	
GLAND DEVELOPMENT OF BALB/C MICE (MUS MUSCULUS) WITH HISTOLOGICAL AND MOLECULAR	
BIOLOGICAL ANALYSIS OF PROTEIN APPROACHES	27
COMPETITIVENESS AND POTENTIAL OF SHEEP LIVESTOCK AS SOURCE INCREASING INCOME AND PROVIDER OF	20
MEAT ANIMAL IN NORTH SUMATRA	30
MORPHOLOGICAL AND PHYSIOLOGICAL CHARACTERS OF CASSAVA (MANIHOT ESCULENTA <i>CRANTZ</i>) WHICH WET	37
THE EFFECT OF SOY TEMPEH FLOUR EXTRACT ON VAGINA HISTOLOGICAL STRUCTURE OF SWISS WEBSTER	52
OVARIECTOMIZED MICE (MUS MUSCULUS)	36
THE TOXICITY OF SEEDS EXTRACT OF ANNONA SQUAMOSA L., LEAVES EXTRACT OF TERMINALIA CATAPPA L. AND	00
LEAVES EXTRACT OF ACACIA NILOTICA L. ON THE MORTALITY OF AEDES AEGYPTI L. LARVAE	39
ELEPHANTOPUS SCABER AND SAUROPUS ANDROGYNUS REGULATE MACROPHAGES AND B LYMPHOCYTE CELLS DURING SALMONELLA TYPHI INFECTION	42
The Effort To Increase Production of Super Red Dragon Fruit (Hylocereus costaricensis) By	
A <mark>rtificial Pollin</mark> ation	
EVALUATION OF ZONATION OF THE MANGROVE CONSERVATION AREAS IN PAMURBAYA	
CENTRAL JAVA	
RELATIONSHIP BETWEEN WATER QUALITY AND ABUNDANCE OF CYANOPHYTA IN PENJALIN RESERVOIR	
HEMATOLOGI <mark>CAL CHARACTERISTIC</mark> OF THE FEMALE ASIAN VINE SNAKE (AHAETULIA PRASINA BOIE, 1827)	
BIOSYNTHESIS SILVER NANOPARTICLE USING FRESH WATER ALGAE.	
EFFECT OF SAPONIN-PODS EXTRACT ACACIA (ACACIA MANGIUM) TO HEMATOCRIT, HEMOGLOBIN AT TILAPIA	05
(OREOCHROMI <mark>S NILOTICUS)</mark>	67
EFFECT OF DISSOLVED NUTRIENT CONCENTRATION (NITRATE AND ORTHOPHOSPHATE) ON ABUNDANCE OF	
CHLOROPHYTA IN PENJALIN RESERVOIR BREBES REGENCY	70
THE ANATOMY OF CAROTENE BIOSYNTHESIS IN BETA VULGARIS L., VAR. RUBRA USING SCAN ELECTRON	
MICROSCOPE	74
OPTIMIZATION OF YOGURT FERMENTED MILK PRODUCTS WITH THE ADDITION OF NATURAL STABILIZER BASED ON	77
LOCAL POTENTIAL OF TARO STARCH (COLOCASIA ESCULENTA)	
PTERIDOPHYTES OF ALAS PURWO NATIONAL PARK AND THEIR MEDICINAL POTENCY	
GENETIC VARIATION OF <i>AEDES AEGYPTI</i> (DIPTERA: CULICIDAE) BASED ON DNA POLYMORPHISM	
MATING BEHAVIOUR OF CROCIDOLOMIA PAVONANA F	
CULTURE	
THE DEVELOPMENT OF SUSTAINABLE RESERVE FOOD GARDEN PROGRAM'S VIDEO IN MALANG CITY	
EFFECT OF MEDIUM COMPOSITIONS ON THE GROWTH OF RICE (ORYZA SATIVA L. CV. CIHERANG) CALLUS	97
BLOOD FIGURE OF RAMBON CATTLE FED FORMULATED CONCENTRATE CONTAINING SOYBEAN CAKE, POLLARD	101
AND CORN OIL COMBINE WITH UREA XYLANASE MOLASSES CANDY	101
PROGRAM TO MEET SELF SUFFICIENCY IN MEAT	103

FOOD TECHNOLOGY	106
MODIFICATION OF BEAN SPROUT AND UREA MEDIA TO SPIRULINA PLATENSIS CULTURE	107
COLLAGEN FROM SEA CUCUMBER (STICHOPUS VARIEGATUS) AS AN ALTERNATIVE SOURCE OF HALAL COLLAGEN	111
DEVELOPMENT OF NEW PRODUCT "COCOA SPIRULINA AS FUNCTIONAL FOOD"	
THE PROTEIN AND WATER CONTENT OF TEN VARIATIONS OF THE FEED CASSAPRO OF YEAST TAPE	. 120
MEDICAL, DENDTISTRY, AND PUBLIC HEALTH	123
EFFECT OF POMELO (CITRUS GRANDIS) ETHANOLIC EXTRACT ON ATHEROSCLEROTIC PLAQUE FORMATION	. 124
CLINICAL MANIFESTATION OF ORAL TUBERCULOSIS	127
IDENTIFICATION OF DERMATOPHYTES BY MULTIPLEX-POLYMERASE CHAIN REACTION, POLYMERASE CHAIN	
REACTION-RESTRICTION FRAGMENT LENGTH POLYMORPHISM ITS1-ITS4 PRIMERS AND MVAI, AND	122
POLYMERASE CHAIN REACTION (GACA) ₄ PRIMER	
ROLE OF REACTIVE OXYGEN SPECIES ON DEVELOPMENTS OF OSTEOCLASTOGENESIS IN AGING	
DETERMINANT FACTOR THAT INFLUENCED ANXIETY LEVEL AND ENERGY INTAKE AMONG ELDERLY	
P-CARE BPJS ACCEPTANCE MODEL IN PRIMARY HEALTH CENTERS	147
THE EFFORT OF TB CADRE IN THE IMPROVING OF THE SUCCESS OF TB THERAPY AND REDUCING SIDE EFFECTS OF	
ANTI TUBERCULOSIS DRUGS	
RISK FACTOR OF GREEN TOBACCO SICKNESS (GTS) AT THE CHILDREN ON TOBACCO PLANTATION	
PHYSICS	
DIRECT SCATTERING PROBLEM FOR MICROWAVE TOMOGRAPHY	158
MICROSTRUCTURE AND MECHANICAL PROPERTIES OF DISSIMILAR JOINT OF COLD ROLLED STEEL SHEETS 1.8	
SPCC-SD AND NUT WELD M6 BY SPOT WELDING	
FEATURE EXTRACTION OF HEART SIGNALS USING FAST FOURIER TRANSFORM	. 165
ANALYSIS OF EL NIÑO EVENT IN 2015 AND THE IMPACT TO THE INCREASE OF HOTSPOTS IN SUMATERA AND KALIMANTAN REGION OF INDONESIA	160
SYNTHESIS OF ZINC OXIDE (ZNO) NANOPARTICLE BY MECHANO-CHEMICAL METHOD	
MODELLING DYNAMICS OF ZNO PARTICLES IN THE SPRAY PYROLISIS REACTOR TUBE	
THE INFLUENCE OF EXTREMELY LOW FREQUENCY (ELF) MAGNETIC FIELD EXPOSURE ON THE PROCESS OF MAKING CREAM CHEESE	
Au Grade of Epithermal Gold Ore at Paningkaban ASGM, Banyumas District, Central Java Province, Indonesia	
RENEWABLE ENERGY CONVERSION WITH HYBRID SOLAR CELL AND FUEL CELL	188
RADAR ABSORBING MATERIALS DOUBLE LAYER FROM LATERITE IRON ROCKS AND ACTIVED CARBON OF CASSAVA	100
PEEL IN X-BAND FREQUENCY RANGE	
DEPLOYMENT POROSITY ESTIMATION OF SANDSTONE RESERVOIR IN THE FIELD OF HIDROCARBON EXPLORATION PENOBSCOT CANADA	
SEISMIC RESOLUTION ENHACEMENT WITH SPECTRAL DECOMPOSITION ATTRIBUTE AT EXPLORATION FIELD IN	
CANADA	199
SIMULATION OF I-V CHARACTERISTICS OF SI DIODE AT DIFFERENCE OPERATING TEMPERATURE: EFFECT OF IONIZED	204
Impurity Scattering	
THE STUDY OF ELECTRICAL CONDUCTANCE SPECTROSCOPY OF THE INNER MEMBRANE OF SALAK	
THE ACCURACY COMPARISON OF OSCILLOSCOPE AND VOLTMETER UTILIZATED IN GETTING DIELECTRIC CONSTANT	. 203
VALUES	211
WINDOW FILTER (WINTER) TO CAPTURE POLLUTION OF LEAD (PB) FOR HOUSES NEAR THE HIGHWAY TO	
PREVENT HEALTH PROBLEMS	. 214
SIMULATION OF SOLAR CELL DIODE I-V CHARACTERISTICS USING FINITE ELEMENT METHODE INFLUENCE OF P-LAYER THICKNESS	216
GEOLOGY	
	210
GIS-BASED OPTIMIZATION METHOD FOR UTILIZING COAL REMAINING RESOURCES AND POST-MINING LAND USE PLANNING: A CASE STUDY OF PT ADARO COAL MINE IN SOUTH KALIMANTAN	219
QUANTIFICATION MODEL OF QUALITATIVE GEOLOGICAL DATA VARIABLES FOR EXPLORATION RISK ASSESSMENT	226
IN PROSPECT CU-AU PORPHYRY DEPOSIT RANDU KUNING, WONOGIRI, CENTRAL JAVA	
RELOCATION OF HYPOCENTER USING JACOBIAN'S MATRIX AND JEFFREYS-BULLEN'S VELOCITY MODEL	
CHEMISTRY	
	. 239
SYNTHESIS AND CHARACTERIZATION MAGNETIC FE ₃ O ₄ NANOPARTICLE BY USING OLEIC ACID AS STABILIZING	2
AGENTSYNTHESIS OF ZEOLITES FROM LOMBOK PUMICE AS SILICA SOURCE FOR ION EXCHANGER	_
PREPARATION OF NANOBIOCATALYST MICROREACTOR USING IMMOBILIZED ENZYME ONTO NANOPOROUS	. 444
MONOLITHIC POLYMER FOR HIGH SPEED PROTEIN DIGESTION	248

	Analysis of protein profile of neem leaves juice (azadirachta indica L. Juss)	253
	HYDROPHOBIC AEROGEL-BASED FILM COATING ON GLASS BY USING MICROWAVE	256
	PREPARATION AND CHARACTERIZATION OF CACAO WASTE AS CACAO VINEGAR AND CHARCOAL	259
	THE EFFECT OF PHYSICO-CHEMICAL PROPERTIES OF AQUATIC SEDIMENT TO THE DISTRIBUTION OF GEOCHEMICAL	
	FRACTIONS OF HEAVY METALS IN THE SEDIMENT	262
	INCREASED CONCENTRATION OF BIOETHANOL BY RECTIFICATION DISTILLATION SIEVE TRAY TYPE	266
	DETERMINATION OF LEAD IN COSMETIC SAMPELS USING COATED WIRE LEAD (II) ION SELECTIVE ELECTRODE	
	Based On Phyropillite	270
	PYROLYSIS TEMPERATURE EFFECT ON VOLUME AND CHEMICAL COMPOSITION OF LIQUID VOLATILE MATTER OF	
	Durian Shell	273
	HIGH PERFORMANCE LIQUID CHROMATOGRAPHY OF AMINO ACIDS USING POTENTIOMETRIC DETECTOR WITH A	
	TUNGSTEN OXIDE ELECTRODE	276
	RAINWATER TREATMENT USING TREATED NATURAL ZEOLITE AND ACTIVATED CARBON FILTER	279
	FILTRATION OF PROTEIN IN TEMPE WASTEWATER USING CELLULOSE ACETATE MEMBRANE	282
4 A '	THEMATICS	205
1A		
	IMAGE ENCRYPTION TECHNIQUE BASED ON PIXEL EXCHANGE AND XOR OPERATION	286
	FUZZY ANP METHOD AND INTERNAL BUSINESS PERSPECTIVE FOR PERFORMANCE MEASUREMENT IN	
	DETERMINING STRATEGY SMES	
	APPLICATION OF FUZZY TOPSIS METHOD IN SCHOLARSHIP INTERVIEW	295
	The Effect of Inflation, Inter <mark>est Rate, and Indonesia Composite Ind</mark> ex (ICI) to the Performances of	
	MUTUAL FUND RETURN AND UNIT LINK WITH PANEL DATA REGRESSION MODELLING	299
	USING LOGISTIC REGRESSION TO ESTIMATE THE INFLUENCE OF ADOLESCENT SEXUAL BEHAVIOR FACTORS ON	
	STUDENTS OF SENIOR HIGH SCHOOL 1 SANGATTA, EAST KUTAI-EAST KALIMANTAN	303
	Application Cluster Analysis on Time Series Modelling with Spatial Correlations for Rainfall Data	
	IN JEMB <mark>er Rege</mark> ncy	
	A ZERO CROSSING-VIRUS EVOLUTIONARY GENETIC ALGORITHM (VEGA) TO SOLVE NONLINEAR EQUATIONS	311
	Analysis of Simultaneous Equation Model (SEM) on Non normally Response used the Method of	
	REDUCE RANK VECTOR GENERALIZED LINEAR MODELS (RR-VGLM)	
	THE RAINBOW (1,2)-CONNECTION NUMBER OF EXPONENTIAL GRAPH AND IT'S LOWER BOUND	319
	CONSTRUCTION OF SUPER H-ANTIMAGICNESS OF GRAPH BY USES A PARTITION TECHNIQUE WITH CANCELATION	225
	NUMBER	
	ON THE TOTAL R-DYNAMIC COLORING OF EDGE COMB PRODUCT GRAPH G D H	
	ON THE METRIC DIMENSION WITH NON-ISOLATED RESOLVING NUMBER OF SOME EXPONENTIAL GRAPH	
	ON TOTAL R-DYNAMIC COLORING OF SEVERAL CLASSES OF GRAPHS AND THEIR RELATED OPERATIONS	
	THE ANALYSIS OF R-DYNAMIC VERTEX COLOURING ON GRAPH OPERATION OF SHACKLE	
	HANDLING OUTLIER IN THE TWO WAYS TABLE BY USING ROBUST AMMI AND ROBUST FACTOR	
	AN EPIDEMIC MODEL OF VARICELLA WITH VACCINATION.	333
	THE CORRELATION BETWEEN PERCEPTION AND BEHAVIOR OF RIVER POLLUTION BY COMMUNITIES AROUND BRANTAS RIVERBANK IN MALANG	250
	ISOLATION AND SCREENING OF SPECIFIC METHICILLIN RESISTANT-STAPHYLOCOCCUS AUREUS BACTERIOPHAGE FRO	
	HOSIPTAL WASTE AT BANYUMAS	
	QUANTIFICATION MODEL OF QUALITATIVE GEOLOGICAL DATA VARIABLES FOR EXPLORATION RISK ASSESSMENT IN	
	PROSPECT CU-AU PORPHYRY DEPOSIT RANDU KUNING, WONOGIRI, CENTRAL JAVA CO (III) AS MEDIATOR IN PHENOL DESTRUCTION USING ELECTROCHEMICAL OXIDATION	
	DESIGN OF SYSTEM BATCH INJECTION ANALYSIS (BIA) FOR MONITORING THE PRODUCTION OF ALCOHOL (II)	
	THOP INDEX	
M 17	HOW INDEX	358



Proceeding The 1st IBSC: Towards The Extended Use Of Basic Science For Enhancing Health, Environment, Energy And Biotechnology ISBN: 978-602-60569-5-5

The Study of Electrical Conductance Spectroscopy of The Inner membrane of Salak

Wenny Maulina¹

¹Biophysics Division: Physics Department, Faculty of Mathematics and Natural Sciences, Jember University, Jember, Indonesia, e-mail: wenny@unej.ac.id

Abstract— Main objective of the present work was to investigate mechanism transport of the inner membrane of salak (*Salacca edulis*) using electrical conductance measurements. The inner thin membrane obtain after removing the outer cover of salak has been found to record electrical conductance when it separates two different concentration of electrolyte solution; NaCl, MgCl₂ and CaCO₃. Conductance values were measured for each concentration at different temperature. The temperature range studied is between 303 K and 333 K at 5 K intervals. The normal behaviour of these conductance shows that the values increase smoothly with increase in concentration. The magnitude follow the order Mg²⁺ > Na⁺ > Ca²⁺. The increase of temperature makes mobilisation of the ions was rapidly during the measurement of conductance.

Keywords— inner thin membrane, salak, conductance.

INTRODUCTION

The membrane is essentially a barrier, which separates two phases and restricts transport of various chemicals in a selective manner. A membrane can be homogeneous or heterogeneous, symmetric or asymmetric, solid or liquid; it can carry positive or negative charges or can be neutral. Transport through a membrane can be affected by convection or by diffusion of individual molecules, and induced by the chemical gradient or electrical gradient. A Membrane should have some basic requirements, such as high flux of the product; good mechanical strength for supporting the physical structure; and good selectivity for the desired substances. Membranes can be either dense or porous. There are various types of membrane processes: a) Microfiltration (pore sizes ranging from 0,1 to 10,0 μm); b) Ultrafiltration (0,001-0,1 μm); c) Reverse Osmosis (0,0001-0,001 μm); and d) Conventional Filtration (10-100 μm). Membranes consist of two types: biological membrane and synthetic membrane [1-2]. Focus on this study is biological membrane i.e. the inner membrane of Salak.

Salak (Salacca edulis) also known as snake fruit belongs to Arecaceae family. This palm is native in Indonesia and Malaysia. Nevertheless, cultivation of salak palm can also be found in Thailand and Philippines recently. Salak fruit is a good source for dietary fibres and carbohydrate. It also contains valuable bioactive antioxidants such as Vitamin A, Vitamin C and Phenolic compound. However, salak has a short shelf life of less than a week. Salak that stored in the room temperature could not be consumed after the day of 14 after harvesting. It is because rapid ripening and degradation of the bioactive ingredients. Numerous studies have been conducted to analysis thermo-physical properties and transport properties of fruit, such as garlic, pumpkin, grapes, apple and orange skin. Several mathematical models have been proposed for the evaluation of transport behaviour by either applying theoretical (Fick's law), semi-theoretical and empirical modelling [3]. Meanwhile, the theoretical models may not able to explain the exact mechanism of transport but it often gives good estimation by incorporating some values into the model parameters. Electrical measurements provide the opportunity to

Generally, electrical properties of the fruit were studied in order to develop a rapid and non-destructive assessment method or to characterize its ripening. Electrical properties are important in the cognitive aspect, especially to find out responses of the fruit to electric fields of variable frequency. Behaviors of the electrical properties are related to the nature of the material (composition, structure), conditions of the material (temperature, frequency), and the age or maturity stage of the material. Utilization of electrical measurements because simple, low cost, and quick assessment of product quality [4].

Transport processes through biological membranes, as well as in many synthetic membranes, are important because of their potential use in different separation processes. Studies on the conductance of electrolyte

solution through such polymeric networks are related to the ions transport through the pores. Extensive research work related to the conductance and diffusion of electrolyte solution through ion-exchange or porous membrane has been reported by different workers. Studies of the biological membrane are necessary to interpretation the mechanism of transport occurring in the biological system [5]. The purpose of the present paper is to report the conductance studies of the inner thin layer membrane of Salak with different electrolyte concentration and various temperatures. The observing of mechanism transport through this membrane has been calculated by using conductance spectroscopy.

METHODS

The inner thin layer membrane was separated manually from the outer of Salak. The membrane was throughly washed with deionised water to remove any adsorbed. It was always kept in the wet condition to avoid any disturbance arise due to the entrapped air within the pores and also to prevent the crack in dry condition. The membranes were cut into rectangular small piece and attached at a chamber. A chamber was filled with a different concentration solution. Solutions of different concentrations were prepared with analytical grade NaCl, MgCl2 and CaCO3 by using distilled water. Platinum electrode used to the electrical connections to the spectroscopy. Conductance measurements were carried out with an LCRmeter Hitester 5322-50. Conductance values were measured for each concentration at different temperature. The variation of concentration electrolyte solution were 0,1 mM, 1 mM, 10 mM, and 100 mM, respectively. The temperature range studied is between 303 K and 333 K at 5 K intervals.

RESULTS AND DISCUSSION

Based on the result of conductance spectroscopic measurements indicated that the inner membrane of salak as a semipermeable membrane. It is evidence by some specific ions could selective transport through the membrane. The nature of variations of conductance with concentrations for different electrolytes at room temperature, are shown in Fig 1. The common trend found for all the electrolytes is that the membrane conductance increases almost linearly with increase in concentration. The magnitude follow the order $Mg^{2+} > Na^+ > Ca^{2+}$. After certain concentrations, these two different concentrations may have a tendency to balance each other.

The increase in conductance of CaCO₃ is not too significant. Because of ion Ca²⁺ has been prevented from passing through the membrane. At this time, membrane shrinkage occurs due to the exclusion of water within the pores, showing steady conductance values. The linear correlation indicates that the inner membrane of Salak is able to respond various changes in increasing concentrations toward values of conductance that through the membrane.






Fig 1. Plot of conductance of different electrolytes through the inner membrane of Salak at room temperature

The effect of temperature on the membrane conductance is shown in Fig 2 represents the specific conductance values for MgCl₂ solutions of different concentrations in the temperature range 303-333 K at 5K intervals. The linear regression with a negative slope shows a correlation between Conductance and Temperature. The increase of temperature makes mobilisation of the ions was rapidly during the measurement of conductance.

By applying the Arrhenius equation in its basic form, where is:

$$G = G_0 exp(-dU/kT) \tag{1}$$

In G values of an various electrolyte are plotted against 1/T. Fig 3 shows the plots of the different electrolytes at the concentration of 100 mM. The smooth linear plots suggest that there may be no abrupt irreversible change in the membrane structure within the concentration and temperature range studied [5].




Fig 2. Plot of conductance of various concentration MgCl₂ through the inner membrane of Salak at different temperatures (303-333 K)




Fig 3. Arrhenius plots of ln G at one of the concentrations (100 mM) against 1/T at different electrolytes through the inner membrane of Salak

1

CONCLUSION

The inner membrane of Salak indicated as a semipermeable membrane. The normal behaviour of conductance shows that the values increase smoothly with increase in concentration. The slope shows correlation between changes of temperature with a conductance as a mechanism transport ion through the inner membrane.

REFERENCES

- [1] J.P. Chen, H. Mou, L.K. Wang, T. Matsuura, and Y. Wei, "Membrane Separation: Basics and Applications," in Handbook of Environmental Engineering, Volume 13: Membrane and Desalination Technologies. Springer Science, Business Media, LLC, 2011, pp. 271-332.
- [2] R.W. Baker, Membrane Technology and Applications, 2nd ed, England: John Wiley & Sons Ltd, 2004.
- [3] S.P. Ong and C.L. Law, "Mathematical Modelling of Thin Layer Drying of Salak", Journal of Applied Sciences., vol. 9, no. 17, 2009, pp. 3048-3054
- [4] J. Juansah, I.W. Budiastra, K. Dahlan, and K.B. Seminaar, "Electrical Behavior of Garut Citrus Fruits During Ripening Changes in Resistance and Capacitance Models of Internal Fruits", International Journal of Engineering & Technology IJET-IJENS, vol. 12, no. 04, 2012, pp. 1-8
- [5] T. Samanta, "Physico-chemical Aspects of Some Biological Membranes. II Studies on The Conductances of The Inner Membrane of Chicken Egg in Presence of Alkali Chlorides and The Application of Absolute Reaction Rate Theory", Colloids and Surfaces B: Biointerfaces, vol 27, 2002, pp. 95-101.