

Digital Repository Universitas Jember



UNIVERSITAS AIRLANGGA



AnMicro

ASEAN Network
on Microbial Utilization



**The 23rd Seminar and 12nd Congress of Indonesian Society
for Biochemistry and Molecular Biology
in Conjunction with the 2nd International Conference**

“Collaboration Seminar of Chemistry and Industry (CoSCI)”

IOP Conference Series
Proceedings services for science

Surabaya – East Java, Indonesia




PAPER • OPEN ACCESS

The 12th Congress of Indonesian Society for Biochemistry and Molecular Biology in Conjunction With The 2nd International Conference “Collaboration Seminar of Chemistry and Industry (CoSCI)” and AnMicro Workshop

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011001

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Preface

Second International Conference “Collaboration Seminar of Chemistry and Industry (CoSCI 2018) in conjunction with 23rd Indonesian Society for Biochemistry and Molecular Biology (ISBMB) seminar was taken place at Universitas Airlangga, Surabaya-Indonesia on October 11-12, 2018. The event presented a theme” Recent Development of Omics Technology For Human Prosperity”.

"Omics is a general term for a broad discipline of science and engineering for analyzing the interactions of biological information objects in organism. The main focus is on: 1) mapping information objects such as genes, proteins, and ligands; 2) finding interaction relationships among the objects; 3) engineering the networks and objects to understand and manipulate the regulatory mechanisms; and 4) integrating various omes and omics subfields.

The event was held to facilitate for the scientists, scholars, engineers and students from universities, research institutes and industries to present ongoing research activities, especially in Biochemistry, Biology, Chemistry, Medicine and other in related fields. It was also to encourage future collaboration among all participants.

The conference of CoSCI and ISBMB created proceedings from the papers that were submitted by participants, after they were reviewed by committee members and international reviewers. This volume intends to provide readers with the recent advances in the Omics Technology such as Chemistry, Biochemistry and Molecular Biology, and Medicine field.

We would like to thank to all authors who contributed to the proceedings and also to the organizing committee, reviewers, speakers, sponsor, and all the conference participants for their supporting in the conference of CoSCI 2018 and 23rd Seminar of ISBMB.

Dr. Purkan, M.Si

Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga,
Surabaya, Indonesia, Nov. 10, 2018



PAPER • OPEN ACCESS

Committee

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011002

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

COMMITTEE

The Chairman

1. Dr. Purkan, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia.
2. Dr. Sri Sumarsih, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia

Steering Committee

1. Dr. Abdulloh, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. Dr. Alfinda Novi Kristanti, DEA (Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
3. Dr. Ni'matuzarroh, M.Si, Biology Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
4. Satya Candra Wibawa Sakti, M.Sc, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
5. Harsasi Setyawati, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
6. Ahmadi Jaya Permana, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
7. Kautsar UI Haq, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
8. Dr. Rico Ramadhan, M.Sc, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
9. Dr. Indeswati Diyatri, drg., M.S, Faculty of Dentistry, Universitas Airlangga
10. Dr. Dian Mulawarmanti, drg., M.S, Faculty of Dentistry, Hang Tuah University, Surabaya Indonesia

Scientific Committee

Editors:

1. Ali Rohman, M.Si, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. M. Zakki Fahmi, M.Si, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
3. Dr. Verawat Champreda, M.Sc, National Center for Genetic Engineering and Biotechnology (BIOTEC), Thailand
4. Dr. Sehanat Prasongsuk, Department of Botany, Chulangkorn Universty, Thailand



Reviewers :

1. Prof. Ni Nyoman Tri Puspaningsih, M.Si, Biochemistry Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. Prof. dr. Soetjipto, MS., Ph.D, Biochemistry Department, Faculty of Medicine, Universitas Airlangga
3. Prof. Seung Wook Kim, Dept of Chemical and Biological Eng, Korea University
4. Prof. Rendy Kartika, Chemistry Department, Louisiana State University, Baton Rouge, LA
5. Prof. Hunsa Punnapayak, Department of Botany, Chulalongkorn University, Thailand
6. Prof. Antonius Suwanto, Division of Microbiology, Institut Pertanian Bogor, Indonesia
7. Prof. Dessy Natalia, Biochemistry Research Division, Faculty of Mathematics and Natural Sciences, Institut Teknologi Bandung, Indonesia
8. Prof. Dr, Suraini Abdul Aziz, Faculty of Biotechnology and Biomolecular Sciences, University Putra Malaysia
9. Prof. Afaf Baktir, Biochemistry Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
10. Dr. Mulyadi Tanjung, M.Si, Natural Product Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
11. Tjitjik Srie Tjahjandarie, Ph.D, Natural Product Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
12. Prof Fedik Rantam, Stem Cell Research Division, Faculty of Veterinary, Universitas Airlangga, Indonesia
13. Prof. Yosephine Sri Wulan Manuhara, Biology Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
14. Dr. Hery Suwito, M.Si (Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
15. Dr. Nanik Siti Aminah, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
16. Dr. rer. nat. Ganden Supriyanto, M.,Sc, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
17. Dr. Pratiwi Pudjiastuti, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia

PAPER • OPEN ACCESS

Peer review statement

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011004

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the [collection](#) - download the first chapter of every title for free.

Peer review statement

All papers published in this volume of *IOP Conference Series: Earth and Environmental Science* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.



PAPER • OPEN ACCESS

Revealing the important role of allosteric property in sucrose phosphate synthase from sugarcane with N-terminal domain deletion

To cite this article: W D Sawitri and B Sugiharto 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 012043

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the [collection](#) - download the first chapter of every title for free.

Revealing the important role of allosteric property in sucrose phosphate synthase from sugarcane with N-terminal domain deletion

W D Sawitri^{1,2*}, B Sugiharto^{1,2}

¹Prodi Magister Bioteknologi, Program Pascasarjana, Universitas Jember, Jl. Kalimantan 37, Jember 68121, Jawa Timur, Indonesia

²Laboratorium Bioteknologi dan Biologi Molekul, UPT Laboratorium Terpadu dan Sentra Inovasi CDAST, Universitas Jember, Jl. Kalimantan 37, Jember 68121, Jawa Timur, Indonesia

*widhi.pasca@unej.ac.id

Abstract. Sucrose occupies many essential roles to control regulation of carbon partitioning in plants, including prokaryotic cells. Sucrose phosphate synthase (SPS; EC 2.4.1.14) is a key enzyme to catalyze the form of sucrose in primary sucrose synthesis pathway. Plants SPS has a molecular size around 120 kDa, which consists of N-terminal domain, C-terminal domain, and central domain. We produced the recombinant sugarcane SPS (*SoSPS1*) in *Escherichia coli*, however, the expression often appears to be a shorter form with retained enzyme activity. In our result, we reported that the shorter form is suggested to have a truncated N-terminal 20-kDa region. The truncated form of *SoSPS1* (Δ N-SPS) tends to enhance the specific activity 10-fold compared to full-length *SoSPS1*. The full-length *SoSPS1* showed a remarkable allosteric activation by glucose-6-phosphate (G6P), while none of the N-terminal truncated form had such a characteristic. By kinetic analysis of full-length *SoSPS1*, a higher substrate affinity was shown in the presence of G6P. Conversely, the Δ N-SPS showed a similar substrate affinity whether G6P was added or not. Based on these results, we revealed that N-terminal region of *SoSPS1* has essential role for allosteric regulation by G6P and may function like a suppressor domain for the enzyme activity.

Keywords: sucrose phosphate synthase, N-terminus deletion, recombinant enzyme, sugarcane

1. Introduction

In most plants, sucrose is the major products of photosynthesis and common form of carbohydrate transported to various growing tissues and storage organs for allocation of carbon resources. Sucrose phosphate synthase (SPS) plays a crucial role to catalyze the formation of sucrose-6-phosphate (S6P) from fructose-6-phosphate (F6P) and uridine diphosphate-glucose (UDP-G) as substrates [1; 2; 3]. The activity of SPS is allosterically controlled by metabolites, such as glucose-6-phosphate (G6P) and inorganic phosphate [4]. However, the precise allosteric mechanism of SPS is still unclear. Such molecular characteristics of SPS are believed to be crucial for playing the physiological role of regulating photosynthetic carbon flux into sucrose [18; 20].

Plants SPS has a molecular size around 120 kDa, which consists of N-terminal domain, C-terminal domain, and central domain. The central region contains glycosyltransferase domain responsible for the catalytic function of SPS [5]. The C-terminal region resembles sucrose phosphate phosphatase and the N-terminal region has no clear similarity with any other proteins [17].

It has previously cloned cDNAs from sugarcane leaves, *SoSPS1* and *SoSPS2*, encoding two isoforms of SPS. *SoSPS1* is abundantly expressed in sugarcane leaves and is considered to be a representative of the isoenzyme responsible for photosynthetic carbon allocation with the regulatory function [6]. In this study, cDNA of *SoSPS1* was expressed in *E. coli* cells and recombinant enzymes with various lengths of the N-terminal deletion were produced. We report here that the N-terminal region of sugarcane SPS is not essential for the catalytic reaction itself, but plays a role in the allosteric regulation.

2. Experimental Method

A series of mutant SPSs with the N-terminal truncation was constructed and generated by PCR. The PCR product of the N-terminal truncated forms designated as His Δ N1, His Δ N2, His Δ N3, His Δ N4,



His Δ N5 and His Δ N6 were subcloned into pGEM-T vector (Promega) at the insertion site and then cut with SbfI and EcoRI. The resulting DNA fragments encoding the N-terminal truncated SPSs were inserted into PstI and EcoRI sites of pTrcHisA. All the SPS genes thus constructed were verified by DNA sequencing.

The constructed plasmids were introduced into *E. coli* BL21 (DE3) competent cells (Novagen) to express the full length SPS and various N-terminal truncated forms. The transformed bacterial cells were expressed at 20°C, then 0.05mM IPTG was added and continued overnight. The bacterial cells were suspended in extraction buffer [50mM Tris-HCl (pH 7.5), 150mM NaCl, 1mM MgCl₂, 1mM EDTA and 1mM PMSF], and disrupted by sonication on ice. Cell homogenate was centrifuged at 15,000 rpm for 15 min at 4°C and the resulting supernatant was mixed gently with DE52 anion exchange cellulose (Whatman). The filtrate was directly loaded onto a column of cComplete His-Tag Purification Resin (Roche) equilibrated with 50 mM Tris-HCl (pH 7.5)/150mM NaCl. The column was washed with the same buffer containing 30mM imidazole and 10% glycerol (w/v) and the resin binding proteins were eluted by increasing concentration of imidazole to 120mM. The eluted sample from the His-tag resin column was concentrated and applied to size exclusion chromatography on a Superdex 200 column (GE Healthcare Life Sciences) and eluted with 50mM Tris-HCl (pH 7.5)/150mM NaCl.

Proteins in fractions obtained during purification were separated by sodium dodecyl sulphate (SDS) polyacrylamide-gel (10%) electrophoresis (PAGE) as described by Apriasti [7]. The gels were directly stained with Coomassie Brilliant Blue. The quantification of protein bands stained with Coomassie Brilliant Blue and visualized by the colour development was performed by densitometry using ImageJ software (<http://imagej.nih.gov/ij/>).

SPS activity was assayed as described previously [8] with an appropriate modification. The assay mixture (50 ml) contained 50mM Hepes-NaOH (pH 7.5), 20mM MgCl₂, 20mM F6P and 20mM UDP-G. The mixture was incubated at room temperature for 10 min and reaction was stopped by an addition of 35 ml of 1M NaOH, followed by incubation at 95°C for 10 min to decompose unreacted F6P. To determine sucrose formed by the enzyme reaction, 125 ml of 0.1% (w/v) resorcinol in 95% ethanol and 30% (w/v) HCl was added and heated at 85°C for 8 min. The developed color of sucrose derivative was measured at absorbance 520nm with using a microtiter plate reader.

3. Results and Discussion

Activity of the truncation forms of His Δ N2 to His Δ N6, using at least partially purified preparation (Fig. 1A), were measured in the presence of the saturating concentrations of F6P and UDP-G, together with that of the full length SPS. As shown in Fig. 1B, the truncated forms showed an increasing tendency of the specific activity and this was the most remarkable for His Δ N6 that showed an increase in the activity higher by 10-fold than that of the full length SPS. These results were in good agreement with previous report [9] that SPS activity in the *E. coli* cells became higher when accumulation level of Form B was increased. Form B was a proteolytic product of recombinant full-length SPS when it was produced from *E. coli* expression system.




Figure 1 Comparison of activity of the N-terminal truncated and full length. (A) SDS-PAGE analysis of various forms of SPS used for activity measurement. (B) Specific activity of the N-terminal truncated and full-length forms of SPS




Figure 2 Kinetic properties of the N-terminal truncated and full-length. (A) Activity of various forms of SPS was measured in the assay mixture containing 20mM UDP-G and increasing concentrations of F6P from 0 to 10mM in the presence or absence of 6mM G6P. (B) Activity of various forms of SPS was measured in the assay mixture containing 10mM F6P and increasing concentrations of UDP-G from 0 to 20mM in the presence or absence of 6mM G6P.




Figure 3 An allosteric effect of G6P on the activity of recombinant SPSs and authentic SPS from plant leaves (maize leaf)

By kinetic analysis of the full length and truncated forms of sugarcane SPS thus prepared, three important results were obtained: (i) the full length enzyme showed a remarkable allosteric activation by G6P, while none of the truncated enzymes tested here had such characteristics (**Fig. 3**), (ii) specific activity of the full length and truncated SPSs showed a tendency that longer truncation has higher activity (**Fig. 1**) and (iii) whether G6P was added or not, the truncated forms showed similar substrate affinity with F6P and UDP-G, while the full length enzyme showed a higher substrate affinity when G6P was present (**Fig. 2A** and **2B**).

Plant SPSs consist of three domains, glucosyltransferase domain (GTD) (55 kDa), the C-terminal phosphohydrolase domain (30 kDa) and the N-terminal domain (20 kDa). The N-terminal domain has no apparent homology with known motifs [10]. Notably, cyanobacteria SPSs [11] and *H. orenii* SPS (HoSPS) [12], which lack the N-terminal domain of plant SPSs, are not allosterically regulated, implying an involvement of the N-terminal domain in the allosteric regulation. Our present data give a clear piece of evidence that the N-terminal region composed of at least 171 residues of sugarcane SPS is crucial for its allosteric sensitivity to G6P and that the N-terminal truncation as short as 37 residues leads the mutant enzyme to be constitutively active with the allosteric sensitivity lost. The N-terminal region of plant SPS was postulated to be intrinsically disordered by bioinformatics analysis. It has been proposed that polypeptide regions with intrinsic disorders play a role for modulation of allostery [13; 15; 16], such as optimizing

high or low affinity binding and controlling the ability to interact the protein surface [14]. Therefore, plant SPS might be one of examples for such allosteric modulation.

4. Conclusions

Eventually, present biochemical studies has clarified the functional roles of the N-terminal region of sugarcane SPS. It would be advantageous to analysis both the full length and the N-terminal truncated forms for elucidation of the precise model of the allostery. In addition, dynamic simulations analysis may useful for mapping the energy landscape of allosteric process. These combined studies would provide an insight into molecular mechanism of SPS for controlling carbon partitioning in plant sucrose biosynthesis [19].

References

- [1] Salerno, G. L., & Curatti, L. 2003 Origin of sucrose metabolism in plants: when, how, and why?. *Trends Plant Sci.* **8** 63-69
- [2] Lunn, J. E., & MacRae, E. 2003 New complexities in the synthesis of sucrose. *Curr. Opin. Plant Biol.* **6** 208-214
- [3] Sugiharto, B. 2018 Chapter 8: Biotechnology of Drought-Tolerant Sugarcane, *Sugarcane-Technology and Research*, IntechOpen 139-165
- [4] Chen, W. L., Yang, C. C., Lee, P. D. 2007 Cloning and Expression of Sweet Potato Tuber Sucrose Phosphate Synthase Gene in Escherichia coli *Taiwanese J. Agric. Chem Food Sci.* **45(2)** 91-100
- [5] Sawitri, W. D., Afidah, S. N., Nakagawa, A., Hase, T., & Sugiharto, B. 2018 Identification of UDP-glucose binding site in glycosyltransferase domain of sucrose phosphate synthase from sugarcane (*Saccharum officinarum*) by structure-based site-directed mutagenesis. *Biophys. Rev.* **10** 293-298
- [6] Castleden C. K., Aoki, N., Gillespie, V. J., MacRae, E. A., Quick, W. P., Buchner, P., Foyer, C. H., Furbank, R. T., & Lunn, J. E. 2004 Evolution and Function of the Sucrose-Phosphate Synthase Gene Families in Wheat and Other Grasses. *Plant Physiol.* **135** 1753-1764
- [7] Apriasti, R., Widyaningrum, S., Hidayati, W. N., Sawitri, W. D., Darsono, N., Hase, T., & Sugiharto, B. 2018 Full sequence of the coat protein gene is required for the induction of pathogen-derived resistance against sugarcane mosaic virus in transgenic sugarcane. *Mol. Biol. Rep.* doi: 10.1007/s11033-018-4326-1
- [8] Takahashi, S., Ono, K., Ugaki, M., Ishimaru, K., Aoki, N., & Ohsugi, R. 2000 Ser 162-Dependent Inactivation of Overproduced Sucrose-Phosphate Synthase Protein of Maize Leaf in Transgenic Rice Plants. *Plant Cell Physiol.* **41(8)** 977-981
- [9] Sawitri, W.D., Narita, H., Ishizaka-Ikeda, E., Sugiharto, B., Hase, T., & Nakagawa, A. 2016 Purification and characterization of recombinant sugarcane sucrose phosphate synthase expressed in E. coli and insect Sf9 cells: an importance of the N-terminal domain for an allosteric regulatory property. *J. Biochem.* **159(6)** 599-607.
- [10] Cumino, A., Curatti, L., Giarrocco, L., & Salerno, G.L. 2002 Sucrose metabolism: Anabaena sucrosephosphate synthase and sucrose-phosphate phosphatase define minimal functional domains shuffled during evolution. *FEBS Lett.* **517** 19-23
- [11] Lunn, J. E., Gillespie V. J., & Furbank, R. T. 2003 Expression of cyanobacterial sucrose-phosphate synthase from Synechocystis sp. PCC 6803 in transgenic plants. *J. Exp. Bot.* **381** 223-237
- [12] Chua, T.K., Bujnicki, J.M., Tan, T.C., Huynh, F., & Patel, B.K. 2008 The structure of sucrose phosphate synthase from Halothermothrix orenii reveals its mechanism of action and binding mode. *Plant Cell.* **20** 1059-1072
- [13] Ferreon, A.C.M., Ferreom, J.C., Wright, P.E., & Deniz, A.A. 2013 Modulation of allostery by protein intrinsic disorder. *Nature* **498** 308-310
- [14] Hilser, V.J. and Thompson, E.B. 2007 Intrinsic disorder as a mechanism to optimize allosteric coupling in proteins. *Proc. Natl. Acad. USA* **104** 8311-8315
- [15] Baskaran, S., Roach, P. J., Roach, D., & Hurley, T. D. 2010 Structural basis for glucose-6-phosphate activation of glycogen synthase. *Proc. Natl. Acad. USA* **107** 17563-17568
- [16] Pederson, B. A., Cheng C., Wilson, W. A., & Roach, P. J. 2000 Regulation of glycogen synthase. *J. Biol. Chem.* **275** 27753-27761
- [17] Haighler, C.H., Singh, B., Zhang, D., Hwang, S., Wu, C., Cai, W.X., Hozain, M., Kang, W.,

- Kiedaisch, B., Strauss, R.E., Hequet, E.F., Wyatt, B.G., Jividen, G.M., & Holaday, A.S. 2007 Transgenic cotton over-producing spinach sucrose phosphate synthase enhanced leaf sucrose synthesis and improved fiber quality under controlled environmental conditions. *Plant Mol. Bio.* **63** 815-832
- [18] Maloney, V. J., Park, J. Y., Unda, F., & Mansfield, S. D. 2015 Sucrose phosphate synthase and sucrose phosphate phosphatase interact *in planta* and promote plant growth and biomass accumulation. *J. Exp. Bot.* **66(14)** 4383-4394
- [19] Sharkey, T.D., Laporte, M.M., dan Kruger, E.L. 2000 "Will increased photosynthetic efficiency lead to increased yield in rice?", *Redesigning Rice Photosynthesis to Increase Yield*. Elsevier Science B.V. 73
- [20] Bab, V. M., & Haigler, C. H. 2001. Sucrose phosphate synthase activity rises in correlation with high-rate cellulose synthesis in three heterotropic systems. *Plant physiol.* **127(3)** 1234-1242

Acknowledgement

We acknowledged Hibah KERIS Universitas Jember and RISTEK DIKTI (Grant: Penelitian Unggulan Strategis Nasional (PUSNAS)).

