

## Society

### INTERNATIONAL UNION FOR PURE AND APPLIED BIOPHYSICS



The International Union for Pure and Applied Biophysics (IUPAB) is a member of the ICSU (International Council for Science) family. Affiliated to it are the national adhering bodies of 50 countries. Its function is to support research and teaching in biophysics. Its principal regular activity is the triennial International Congresses and General Assemblies.

IUPAB and Springer launched the journal "Biophysical Reviews" in 2009.

- [INTERNATIONAL UNION FOR PURE AND APPLIED BIOPHYSICS](#)
- **Ethics & Disclosures**
- The journal is committed to maintaining the highest level of integrity in the content published.
- This journal has a Conflict of Interest policy in place and complies with international, national and institutional standards on research involving Human Participants and Animals and Informed Consent.



- 
- The journal is a member of the Committee on Publication Ethics (COPE) and subscribes to its principles on how to deal with acts of misconduct thereby committing to investigate allegations of misconduct in order to ensure the integrity of research.
- Content published in this journal is peer reviewed.



## **Editor-in-Chief**

**Cristobal G. dos Remedios**

Discipline of Anatomy & Histology  
Bosch Institute, Anderson Stuart Building (F13)  
University of Sydney, Sydney 2006, Australia  
Email: crisdos@anatomy.usyd.edu.au

## **Deputy Editor**

**Damien Hall**

Australian National University  
Research School of Chemistry  
Canberra, ACT, Australia

## **Editors Special Issues**

**Antonio Benedetto** (*Ionic liquids*)

University College Dublin  
Dublin, Ireland

**Elisabeth Ehler** (*Non-myofibrillar causes of human heart failure*)

University of London King's College  
Biophysics  
London, UK

**Amy Li** (*Titin*)

University of Vermont  
Burlington, VT, USA

## **Executive Editors**

**N. R. Jagannathan**

All India Institute of Medical Sciences  
Department of NMR & MRI Facility  
New Delhi 110029, India

**H. Nakamura**

Laboratory of Protein Informatics  
Research Center for Structural and Functional Proteomics  
Institute for Protein Research  
Osaka University  
3-2 Yamadaoka, Suita  
Osaka 565-0871, Japan

**W. Olson**

Department of Chemistry & Chemical Biology  
Rutgers – School of Arts and Sciences

10 Taylor Road  
Piscataway, NJ 08854, USA

## **Rosangela Itri**

Instituto de Física  
Universidade de São Paulo  
São Paulo, Brazil

## **Editorial Board**

T.W. Allen, Royal Melbourne Institute of Technology, Victoria, Australia  
M. Auger, Université Laval, Québec, Canada  
J. Baenziger, University of Ottawa, Canada  
E. Baulieu, Institut National de la Santé et de la Recherche Médicale, Le Kremlin-Bicêtre, France  
A. Benedetto, University College Dublin, Ireland  
L. Blanchoin, Institut de Recherches en Technologies et Sciences pour le Vivant, Grenoble, France  
P. Ciancaglini, Universidade de São Paulo, Ribeirao Preto - SP, Brazil  
P. J. Cozzone, Université de la Méditerranée, Marseille, France  
D. Crossman, Auckland, New Zealand  
E. M. De La Cruz, Yale University, New Haven, CT, USA  
E. Ehler, University of London King's College, London, UK  
L. Finzi, Emory University, Atlanta, Georgia, USA  
H.J. Galla, University of Muenster, Germany  
K. Gehmlich, Cardiovascular Medicine, University of Oxford, UK  
J. C. Gómez-Fernández, University of Murcia, Murcia, Spain  
J.W.K. Ho, Victor Chang Cardiac Research Institute, Darlinghurst, NSW, Australia  
R. Itri, Instituto de Física, Universidade de São Paulo, São Paulo, Brazil  
J. Kozelka, University Paris Descartes, France & Masaryk University, Brno, Czech Republic  
A. Li, University of Vermont, Burlington, VT, USA  
B. Maggio, Universidad Nacional de Córdoba, Córdoba, Argentina  
M. M. Morales, Cidade Universitária, Rio de Janeiro, Brazil  
K. Nagayama, National Institute for Physiological Sciences, Okazaki, Japan  
H. Noji, The University of Tokyo, Tokyo, Japan  
G. U. Nienhaus, Ulm University, Ulm, Germany  
P. Pohl, University Linz, Linz, Austria  
G.C. K. Roberts, University of Leicester, Leicester, U.K.  
C. Royer, Rensselaer Polytechnic Institute, Troy, NY, USA  
F. Separovic, University of Melbourne, Melbourne, Australia  
T. Sewell, University of Cape Town, Cape Town, South Africa  
C. M. Soares, Universidade Nove de Lisboa, Oeiras, Portugal  
M. Steenman, INSERM, Nantes, France  
D. Szczesna-Cordary, University of Miami, FL, USA  
J. Takagi, Institute for Protein Research, Osaka University, Japan  
A. Watts, University of Oxford, Oxford, UK



G.J.L. Wuite, VU University Amsterdam, The Netherlands  
K. Yasuda, Waseda University, Tokyo, Japan

## **Editor Emeritus**

**J. Garnier**, International Council for Science (ICSU), Paris, France

Biophysical Reviews publishes critical and timely reviews from key figures in the field of biophysics. Interested authors are encouraged to discuss the possibility of contributing a review with the Editor-in-Chief. Biophysical Reviews covers the entire field of biophysics, generally defined as the science of describing and defining biological phenomenon using the concepts and the techniques of physics. This includes but is not limited by such areas as bioinformatics, biophysical methods and instrumentation, medical biophysics, biosystems, cell biophysics and organisation, macromolecules: dynamics, structures and interactions, and membrane biophysics, channels and transportation.

Biophysical Reviews mainly targets basic research scientist, but as a number of reviews also have a strong link to medicine, the material is also likely to be of relevant to clinicians with an interest in the biophysical basis of diseases, treatments and interventions. Students of biophysics and the biomedical sciences will also find the reviews useful as they present in depth analyses of important topics.

**Related subjects** » [Biochemistry & Biophysics](#) - [Biophysics & Biological Physics](#) - [Cell Biology](#) - [Nanotechnology](#)

## **Abstracted/Indexed in**

PubMedCentral, SCOPUS, INSPEC, EMBASE, Chemical Abstracts Service (CAS), Google Scholar, Academic OneFile, CNKI, Gale, INIS Atomindex, OCLC, ProQuest Advanced Technologies & Aerospace Database, ProQuest Biological Science Database, ProQuest Natural Science Collection, ProQuest SciTech Premium Collection, ProQuest Technology Collection, Reaxys, Summon by ProQuest

## **Subscription information**

2018. Volume 10 (6 issues each). 66,39 € suggested net price\* plus shipping charges. If an online version exists, subscription includes both the print edition and online access via SpringerLink (Basic SpringerLink License).

1. Editorial  
**Foreword to ‘Multiscale structural biology: biophysical principles and mechanisms underlying the action of bio-nanomachines’, a special issue in Honour of Fumio Arisaka’s 70th birthday**  
Damien Hall, Junichi Takagi, Haruki Nakamura Pages 105-129  
[Download PDF \(3694KB\)](#) [View Article](#)
2. Letter to the Editor  
**Forty years of research on the assembly and infection process of bacteriophage**  
Fumio Arisaka Pages 131-136  
[Download PDF \(166KB\)](#) [View Article](#)
3. Letter to the Editor  
**Congratulations to Dr. Fumio Arisaka on his 70th birthday**  
Peter Schuck Page 137  
[Download PDF \(279KB\)](#) [View Article](#)
4. Brief Communication  
**My happy years of collaboration with Fumio Arisaka**  
Michael G. Rossmann Pages 139-140  
[Download PDF \(768KB\)](#) [View Article](#)
5. Brief Communication  
**Most of it started with T4 phage and was then taken over**  
Shigeki Takeda Pages 141-144  
[Download PDF \(446KB\)](#) [View Article](#)
6. Brief Communication  
**Statistical description of the denatured structure of a single protein, staphylococcal nuclease, by FRET analysis**  
Mariko Yamaguchi, Emi Ohta, Takuya Muto, Takayoshi Watanabe... Pages 145-152  
[Download PDF \(1925KB\)](#) [View Article](#)
7. Review  
**Molecular evolution of an oligomeric biocatalyst functioning in lysine biosynthesis**  
Tatiana P. Soares da Costa, Belinda M. Abbott, Anthony R. Gendall... Pages 153-162  
[Download PDF \(2983KB\)](#) [View Article](#)
8. Review  
**Unified understanding of folding and binding mechanisms of globular and intrinsically disordered proteins**  
Munehito Arai Pages 163-181  
[Download PDF \(1605KB\)](#) [View Article](#)
9. Review  
**Anaerobic crystallization of proteins**  
Miki Senda, Toshiya Senda Pages 183-189  
[Download PDF \(1312KB\)](#) [View Article](#)
10. Review  
**Molluscan hemocyanin: structure, evolution, and physiology**  
Sanae Kato, Takashi Matsui, Christos Gatsogiannis, Yoshikazu Tanaka Pages 191-202  
[Download PDF \(4524KB\)](#) [View Article](#)
11. Review  
**Protein-solvent interaction**

- Tsutomu Arakawa Pages 203-208  
[Download PDF \(523KB\)](#) [View Article](#)
12. Review  
**Serial femtosecond crystallography at the SACLA: breakthrough to dynamic structural biology**  
Eiichi Mizohata, Takanori Nakane, Yohta Fukuda, Eriko Nango... Pages 209-218  
[Download PDF \(1130KB\)](#) [View Article](#)
13. Review  
**“Just a spoonful of sugar...”: import of sialic acid across bacterial cell membranes**  
Rachel A. North, Christopher R. Horne, James S. Davies... Pages 219-227  
[Download PDF \(649KB\)](#) [View Article](#)
14. Review  
**Analytical ultracentrifugation in structural biology**  
Satoru Unzai Pages 229-233  
[Download PDF \(821KB\)](#) [View Article](#)
15. Review  
**Ion–protein interactions of a potassium ion channel studied by attenuated total reflection Fourier transform infrared spectroscopy**  
Yuji Furutani Pages 235-239  
[Download PDF \(1626KB\)](#) [View Article](#)
16. Review  
**Toward an understanding of biochemical equilibria within living cells**  
Germán Rivas, Allen P. Minton Pages 241-253  
[Download PDF \(1234KB\)](#) [View Article](#)
17. Review  
**Heat denaturation of the antibody, a multi-domain protein**  
Yoko Akazawa-Ogawa, Hidenori Nagai, Yoshihisa Hagihara Pages 255-258  
[Download PDF \(435KB\)](#) [View Article](#)
18. Review  
**Sedimentation velocity analytical ultracentrifugation for characterization of therapeutic antibodies**  
Susumu Uchiyama, Masanori Noda, Elena Krayukhina Pages 259-269  
[Download PDF \(2309KB\)](#) [View Article](#)
19. Review  
**A role of heme side-chains of human hemoglobin in its function revealed by circular dichroism and resonance Raman spectroscopy**  
Masako Nagai, Naoki Mizusawa, Teizo Kitagawa, Shigenori Nagatomo Pages 271-284  
[Download PDF \(2146KB\)](#) [View Article](#)
20. Review  
**High-speed atomic force microscopy and its future prospects**  
Toshio Ando Pages 285-292  
[Download PDF \(3091KB\)](#) [View Article](#)

21. Review  
**Identification of UDP-glucose binding site in glycosyltransferase domain of sucrose phosphate synthase from sugarcane (*Saccharum officinarum*) by structure-based site-directed mutagenesis**  
[Widhi Dyah Sawitri, Siti Nurul Afidah, Atsushi Nakagawa...](#) Pages 293-298  
[Download PDF \(1419KB\)](#) [View Article](#)
22. Review  
**Multiple analyses of protein dynamics in solution**  
[Tadayuki Ogawa, Nobutaka Hirokawa](#) Pages 299-306  
[Download PDF \(1847KB\)](#) [View Article](#)
23. Review  
**Lipid environment of membrane proteins in cryo-EM based structural analysis**  
[Kazuhiro Mio, Chikara Sato](#) Pages 307-316  
[Download PDF \(1287KB\)](#) [View Article](#)
24. Review  
**Single-molecule fluorescence-based analysis of protein conformation, interaction, and oligomerization in cellular systems**  
[Kenji Okamoto, Michio Hiroshima, Yasushi Sako](#) Pages 317-326  
[Download PDF \(963KB\)](#) [View Article](#)
25. Review  
**Light-induced chromophore and protein responses and mechanical signal transduction of BLUF proteins**  
[Tomotsumi Fujisawa, Shinji Masuda](#) Pages 327-337  
[Download PDF \(2120KB\)](#) [View Article](#)
26. Review  
**Prefoldin, a jellyfish-like molecular chaperone: functional cooperation with a group II chaperonin and beyond**  
[Muhamad Sahlan, Tamotsu Zako, Masafumi Yohda](#) Pages 339-345  
[Download PDF \(1128KB\)](#) [View Article](#)
27. Review  
**Antimicrobial resistance (AMR) nanomachines—mechanisms for fluoroquinolone and glycopeptide recognition, efflux and/or deactivation**  
[Mary K. Phillips-Jones, Stephen E. Harding](#) Pages 347-362  
[Download PDF \(1392KB\)](#) [View Article](#)
28. Review  
**Hypothesis: structural heterogeneity of the unfolded proteins originating from the coupling of the local clusters and the long-range distance distribution**  
[Satoshi Takahashi, Aya Yoshida, Hiroyuki Oikawa](#) Pages 363-373  
[Download PDF \(535KB\)](#) [View Article](#)
29. Review  
**Relaxation mode analysis for molecular dynamics simulations of proteins**  
[Ayori Mitsutake, Hiroshi Takano](#) Pages 375-389  
[Download PDF \(1203KB\)](#) [View Article](#)
30. Review  
**Hierarchical design of artificial proteins and complexes toward synthetic structural biology**



- Ryoichi Arai Pages 391-410  
[Download PDF \(6032KB\)](#) [View Article](#)
31. Review  
**Creation of artificial protein–protein interactions using  $\alpha$ -helices as interfaces**  
Sota Yagi, Satoshi Akanuma, Akihiko Yamagishi Pages 411-420  
[Download PDF \(1860KB\)](#) [View Article](#)
32. Review  
**“Multiple partial recognitions in dynamic equilibrium” in the binding sites of proteins form the molecular basis of promiscuous recognition of structurally diverse ligands**  
Daisuke Kohda Pages 421-433  
[Download PDF \(3750KB\)](#) [View Article](#)
33. Review  
**Gliadins from wheat grain: an overview, from primary structure to nanostructures of aggregates**  
Reiko Urade, Nobuhiro Sato, Masaaki Sugiyama Pages 435-443  
[Download PDF \(2099KB\)](#) [View Article](#)
34. Review  
**Assembly and remodeling of viral DNA and RNA replicons regulated by cellular molecular chaperones**  
Takeshi Sekiya, Yifan Hu, Kohsuke Kato, Mitsuru Okuwaki... Pages 445-452  
[Download PDF \(965KB\)](#) [View Article](#)
35. Review  
**Membrane-mediated amyloid deposition of human islet amyloid polypeptide**  
Kenji Sasahara Pages 453-462  
[Download PDF \(1023KB\)](#) [View Article](#)
36. Review  
**Bacteriophage T4 long tail fiber domains**  
Paul Hyman, Mark van Raaij Pages 463-471  
[Download PDF \(1260KB\)](#) [View Article](#)
37. Review  
**Biophysical studies of protein solubility and amorphous aggregation by systematic mutational analysis and a helical polymerization model**  
Yutaka Kuroda Pages 473-480  
[Download PDF \(1265KB\)](#) [View Article](#)
38. Review  
**How multi-scale structural biology elucidated context-dependent variability in ectodomain conformation along with the ligand capture and release cycle for LDLR family members**  
Terukazu Nogi Pages 481-492  
[Download PDF \(1670KB\)](#) [View Article](#)
39. Review  
**Salt-induced formations of partially folded intermediates and amyloid fibrils suggests a common underlying mechanism**  
Yuji Goto, Masayuki Adachi, Hiroya Muta, Masatomo So Pages 493-502  
[Download PDF \(1736KB\)](#) [View Article](#)

40. Review  
**Environment-transformable sequence–structure relationship: a general mechanism for proteotoxicity**  
[Jianxing Song](#) Pages 503-516  
[Download PDF \(2626KB\)](#) [View Article](#)
41. Review  
**Formation and properties of amyloid fibrils of prion protein**  
[Kei-ichi Yamaguchi](#), [Kazuo Kuwata](#) Pages 517-525  
[Download PDF \(1000KB\)](#) [View Article](#)
42. Review  
**Recent progress on understanding the mechanisms of amyloid nucleation**  
[Eri Chatani](#), [Naoki Yamamoto](#) Pages 527-534  
[Download PDF \(1098KB\)](#) [View Article](#)
43. Review  
**Review of the nature, diversity and structure of bacteriophage receptor binding proteins that target Gram-positive bacteria**  
[Ahmed S. A. Dowah](#), [Martha R. J. Clokie](#) Pages 535-542  
[Download PDF \(1170KB\)](#) [View Article](#)
44. Review  
**Reconstitution of membrane tethering mediated by Rab-family small GTPases**  
[Joji Mima](#) Pages 543-549  
[Download PDF \(1791KB\)](#) [View Article](#)
45. Review  
**Structure and assembly mechanism of virus-associated pyramids**  
[Tessa E. F. Quax](#), [Bertram Daum](#) Pages 551-557  
[Download PDF \(4739KB\)](#) [View Article](#)
46. Review  
**Bacterial flagellar axial structure and its construction**  
[Katsumi Imada](#) Pages 559-570  
[Download PDF \(3428KB\)](#) [View Article](#)
47. Review  
**Overview of the mechanism of cytoskeletal motors based on structure**  
[Yusuke Kato](#), [Takuya Miyakawa](#), [Masaru Tanokura](#) Pages 571-581  
[Download PDF \(3195KB\)](#) [View Article](#)
48. Review  
**States of phage T3/T7 capsids: buoyant density centrifugation and cryo-EM**  
[Philip Serwer](#), [Elena T. Wright](#), [Borries Demeler](#), [Wen Jiang](#) Pages 583-596  
[Download PDF \(3712KB\)](#) [View Article](#)
49. Review  
**Structural insights on the dynamics of proteasome formation**  
[Koichi Kato](#), [Tadashi Satoh](#) Pages 597-604  
[Download PDF \(2953KB\)](#) [View Article](#)
50. Review  
**Multiscale molecular dynamics simulations of rotary motor proteins**  
[Toru Ekimoto](#), [Mitsunori Ikeguchi](#) Pages 605-615  
[Download PDF \(1799KB\)](#) [View Article](#)



# Identification of UDP-glucose binding site in glycosyltransferase domain of sucrose phosphate synthase from sugarcane (*Saccharum officinarum*) by structure-based site-directed mutagenesis

Widhi Dyah Sawitri<sup>1,2</sup> · Siti Nurul Afidah<sup>3</sup> · Atsushi Nakagawa<sup>4</sup> · Toshiharu Hase<sup>5</sup> · Bambang Sugiharto<sup>1,2,3</sup> 

Received: 13 October 2017 / Accepted: 16 November 2017 / Published online: 8 December 2017

© International Union for Pure and Applied Biophysics (IUPAB) and Springer-Verlag GmbH Germany, part of Springer Nature 2017

## Abstract

Sucrose phosphate synthase (SPS) is believed to be the key enzyme for controlling the biosynthesis of sucrose. SPSs consist of a functional glycosyltransferase domain that shares conserved residues with the glycosyltransferase domain of sucrose biosynthesis-related protein. The formation of sucrose-6-phosphate is catalyzed by SPS with the transfer of a glycosyl group of uridine diphosphate glucose (UDP-G) as an activated donor sugar to a fructose-6-phosphate as a sugar acceptor. However, understanding of the mechanism of catalytic and substrate binding in SPS is very limited. Based on amino acid sequence alignments with several enzymes that belong to the glycosyltransferase family, the UDP-G binding sites that might be critical for catalytic mechanism were identified. Here, we report that single point mutation of R496, D498, and V570 located in the proposed UDP-G binding site led to less active or complete loss of enzyme activity. Through structure-based site-directed mutagenesis and biochemical studies, the results indicated that these residues contribute to the catalytic activity of plant SPS. Moreover, understanding of the UDP-G binding site provides an insight into new strategies for enzyme engineering and redesigning a catalytic mechanism for UDP.

**Keywords** Sucrose phosphate synthase · Glycosyltransferase · Uridine diphosphate glucose · Sugarcane · Site-directed mutagenesis

## Abbreviations

F6P Fructose-6-phosphate  
G6P Glucose-6-phosphate

SPS Sucrose phosphate synthase  
UDP-G Uridine diphosphate glucose;

This article is part of a Special Issue on ‘Biomolecules to Bio-nanomachines - Fumio Arisaka 70th Birthday’ edited by Damien Hall, Junichi Takagi and Haruki Nakamura.

✉ Bambang Sugiharto  
sugiharto.fmipa@unej.ac.id

- <sup>1</sup> Center for Development of Advanced Science and Technology (CDAST), University of Jember, Jl. Kalimantan 37, Jember 68121, Indonesia
- <sup>2</sup> Postgraduate Program in Biotechnology, Jember University, Jl. Kalimantan 37, Jember 68121, Indonesia
- <sup>3</sup> Department of Biology, Faculty of Mathematics and Natural Sciences, University of Jember, Jl. Kalimantan 37, Jember 68121, Indonesia
- <sup>4</sup> Research Center for State-of-the-art Functional Protein Analysis, Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka 565-0871, Japan
- <sup>5</sup> Division of Protein Chemistry, Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka 565-0871, Japan

## Introduction

Sucrose is synthesized in cytosol and is the main form of transporting photosynthesis products throughout plant cells (Gifford et al. 1984). Sucrose occupies many significant roles for plant cell metabolism, including cyanobacteria and unicellular algae. The biochemical and molecular studies reported on sucrose biosynthesis in prokaryotic offer new insights into the origin of sucrose metabolism (Lunn and MacRae 2003; Salerno and Curatti 2003).

One of the key enzymes to catalyze the first step of the sucrose synthesis pathway is sucrose phosphate synthase (SPS; EC 2.4.1.14). Plant SPS has a molecular size of approximately 120 kDa which consists of three domains, including a glycosyltransferase domain that regulates the catalytic function of SPS (Castleden et al. 2004). In contrast, recombinant sugarcane SPS (*SoSPS1*) expression in *Escherichia coli* mostly appears to be a shorter form (100 kDa) with retained

enzyme activity, and the shorter form was predicted to have a truncated N-terminal 20-kDa region. It has recently been revealed that the expression of N-terminal truncated form ( $\Delta$ N-SPS) tends to increase the specific activity 10-fold compared to full-length SPS. In our previous results, we reported that the regulation of the N-terminal domain might function as a suppressor domain for plant SPS activity (Sawitri et al. 2016). The crystal structure of non-photosynthetic bacterial SPS has been solved with different characteristics compared to plant SPS. The lack of N-terminal and C-terminal domains of plant SPS are shown in *Halothermothrix orenii* SPS. Therefore, the structure of bacterial SPS contains only a glycosyltransferase domain for substrate binding and catalytic reaction (Salerno and Curatti 2003; Chua et al. 2008). However, the precise regulatory function of SPS in plants still remains unclear since there is no crystal structure of plant SPS.

In this study, we report the identification of substrate-binding sites for uridine diphosphate glucose (UDP-G) as a sugar donor at the glycosyltransferase domain of recombinant *So*SPS1. As noted above, *So*SPS1 has been used in a previous study for elucidating SPS regulation in sugarcane. The expression of *So*SPS1 is actively regulated in the photosynthetic tissue and representative for the sucrose biosynthesis enzyme (Sugiharto et al. 1997). The knowledge of substrate binding sites would provide an important insight into the catalytic mechanism of SPS in plants. Moreover, it is interesting to determine the regulation of plant SPS involved in carbon partitioning which is a critical process in distributing chemical energy converted through photosynthesis.

## Diphosphate moiety binding of UDP to *So*SPS1

The enzymes responsible for regulation in sucrose metabolism are grouped into sucrose-biosynthesis-related proteins (SBRPs), and the conserved domain of glycosyltransferase has been determined in SPS and sucrose synthase (SuSy) (Pontis 1978; Cumino et al. 2002; Salerno and Curatti 2003). Other enzymes that are involved in the synthesis of polysaccharides are likewise members of glycosyltransferase, such as glycogen synthase (GS) and starch synthase (Coutinho et al. 2003; Sheng et al. 2009). The glycosyltransferases are generally utilizing sugar moiety from activated nucleotide sugar as a donor to promote enzymatic reaction (Breton and Imberty 1999; Tarbouriech et al. 2001).

SPS catalyzes the synthesis of sucrose-6-phosphate (S6P) from UDP-G and fructose-6-phosphate (F6P) (Leloir and Cardini 1955; Amir and Preiss 1982; Huber and Huber 1996). The specific substrate for plant SPS is UDP-G, whereas the substrate for bacterial SPS is not specific for UDP-G. Unlike plant SPS, bacterial SPS is able to bind other nucleoside

diphosphate glucose (NDP-G), such as adenosine diphosphate glucose (ADP-G) (Lunn et al. 1999; Chua et al. 2008). In spite of the fact that many glycosyltransferase enzymes recognize similar substrates, the amino acid sequences between different families have diverged (Ünlügil and Rini 2000; Imberty et al. 2006). Thus, identifying the binding sites for UDP-G is necessary to elucidate the mechanism of enzyme regulation in plant SPS.


It has been reported that SPS, SuSy, and GS share a close similarity with the glycosyltransferase domain and the stereochemistry of their substrates, such as UDP-G (Chua et al. 2008; Baskaran et al. 2010; Zheng et al. 2011; Wu et al. 2015). However, the amino acid sequence alignment with conserved residues among SPS, SuSy, and GS has not been well reported (Diricks et al. 2015).

The deduced sequences of *So*SPS1 and other representative enzymes of glycosyltransferases were aligned using ClustalW2 and compared to identify the predicted NDP binding pocket. The sequences were: sugarcane SPS (*So*SPS1, GenBank ID: BAA19242.1), *H. orenii* SPS (*Ho*SPS, GenBank ID: ACB11221.1), *Arabidopsis thaliana* SuSy (*At*SuSy, GenBank ID: NP\_197583.1), *Nitrosomonas europaea* SuSy (*Ne*SuSy, GenBank ID: CAD85125.1), and *Escherichia coli* GS (*Ec*GS, GenBank ID: EDU64330.1). The comparison of the *So*SPS1 sequence alignment with its homologs is shown in Fig. 1a. There are three critical residues, R496, K501, and E591, in *So*SPS1 (corresponding to R580, K585, and E675 in *At*SuSy; R567, K572, and E663 in *Ne*SuSy; R300, K305, and E377 in GS) (Fig. 1a). In previous studies, it has been reported that these critical residues may lead the conformational changes of enzyme architecture. The structure of *N. europaea* sucrose synthase (PDB ID: 4RBN), has proven that these three residues, Arg, Lys, and Glu, have a large effect on its catalytic and conformational change (Wu et al. 2015).

The above-mentioned highly conserved glycosyltransferases have been identified at R496 in *So*SPS1. These conserved Arg residues are essential for interaction between enzyme and substrate UDP-G, notably the diphosphate moiety. The multiple sequence alignment in Fig. 1a shows that plant SuSy is significantly more similar to bacterial SuSy and SPS compared to glycogen synthase. To date, the crystal structure of sucrose synthase from *A. thaliana* (PDB ID: 3S28) in complex with product UDP-G has been solved and the diphosphate group of UDP-G showed to interact with residue R580 (Zheng et al. 2011). Interestingly, the residue D582 in *At*SuSy exhibits a salt bridge interaction with R580 and K585 (Fig. 1b). Asp residue (D582 in *At*SuSy) is not directly involved in the catalytic process, but D582 appears to stabilize R580 and K585 for binding its substrate. Therefore, R496, K501, and D498 in *So*SPS1 might have a crucial contribution to interact with the pyrophosphate group of UDP-G.

In this present work, structure-based site-directed mutagenesis is demonstrated to confirm experimentally the involvement of any residues in UDP-G binding sites of plant SPS. The





**Fig. 1** Comparison of residues in glycosyltransferases group. **a** Multiple sequence alignment of SPS in sugarcane (*So*SPS1), SPS in non-photosynthetic bacterial *H. orenii* (*Ho*SPS), sucrose synthase in *A. thaliana* (*At*SuSy), sucrose synthase in bacteria *N. europaea* (*Ne*SuSy), and glycogen synthase from *E. coli* (*Ec*GS). The three highly conserved and critical residues that consist of arginine, lysine, and glutamic acid are shown in red. The residues that are predicted to be involved in nucleotide


sugar interaction is shown in blue. **b** The UDP binding site residues at positions corresponding to Arg-580, Asp-582, Lys-585, and Glu-675 for diphosphate moiety binding and Gln-648 and Asn-654 for uridine moiety binding of *A. thaliana* sucrose synthase (PDB ID: 3S28). **c** Predicted interaction hydrogen bonding between Ser-269 and Arg-270 in bacterial SPS (PDB ID: 2R60)

point mutation was specified as S495A, R496A, P497A, D498A, R496K and D498E for the diphosphate moiety and V570A for the uridine moiety. The QuikChange Lightning Site-Directed Mutagenesis Kit (Agilent Technologies, USA) was utilized to introduce specific point mutations in the wild-type  $\Delta$ N-*So*SPS1 as previously described (Sawitri et al. 2016). In order to obtain biochemical characteristics, the wild-type and mutants were expressed into *E. coli* BL21 (DE3) using expression vector pTrcHisA (Life Technologies, Invitrogen, USA) and the expressions of wild-type and mutants were monitored by immunoblotting analysis using polyclonal antibody against the SPS protein. As shown in Fig. 2a, the bands of wild-type and mutants with apparent molecular size around 100 kDa were detected as expected. The yield of recombinant production was similar for all expressed proteins, indicating that wild-type and mutants were expressed in *E. coli* BL21 (DE3) cells. The sequential purification of wild-type and mutated proteins were carried out by DE52 anion exchange cellulose (Whatman, UK) and affinity system Complete His-Tag Purification Resin (Roche, Switzerland). The protein in the fraction obtained during the purification steps was monitored using sodium dodecyl sulfate polyacrylamide-gel electrophoresis (SDS-PAGE) as described by Laemmli (1970) and

Coomassie Brilliant Blue staining as shown in Fig. 2b. Afterwards, the eluted samples were concentrated in 10 K Amicon Ultra Centrifugal Filter (Merck Millipore, Ireland) and the purified samples were further assayed for biochemical analysis to characterize the identified residues.

In this report, alanine-scanning mutagenesis has been applied to investigate the predicted substrate-binding site in *So*SPS1 by comparing the enzyme activity of wild-type and mutants (S495A, R496A, P497A, and D498A). SPS activity as previously described (Huber et al. 1985; Sawitri et al. 2016) was measured in the presence of substrate UDP-G and F6P at saturating concentrations (20 mM). As shown in Fig. 2c, enzyme activity of mutant P497A was comparable with wild-type. As expected, residue Pro at 497 is not involved in catalytic activity and does not induce conformational changes in SPS structure. Conversely, single mutation of S495 significantly altered activity of the enzyme. The residue Ser495–Arg496, which corresponds to Ser269–Arg270 in *Ho*SPS, has been suggested to contribute a hydrogen-bonding interaction between Ser and Arg (Fig. 1c). If Ser at S495 is replaced with Ala, it adopts the sequence conserved in SuSy and the SPS activity was remarkably increased compared to wild-type  $\Delta$ N-*So*SPS1. The reason for this phenomenon was not clear, but we





**Fig. 2** Site-directed mutagenesis and expression of recombinant *SoSPS1* in *E. coli*. **a** The resulting SPS mutation at glycosyltransferase domain was transformed into *E. coli* overnight and the total cell proteins were detected by western blotting using antibodies against polypeptide of SPS. **b** The affinity purification of His-tagged  $\Delta$ N-*SoSPS1* as monitored by SDS-PAGE and Coomassie blue staining. *Lane 1* The eluted sample from DE52 anion exchange cellulose. *Lanes 2–5* Washing steps of affinity His-tag purification with increasing concentration of imidazole

0 and 20 mM. *Lanes 6–10* The fraction of samples eluted successively with the buffer containing 100 mM imidazole. *Lanes 11–12* The purified SPSs were concentrated and exchanged to buffer composed of 50 mM Tris-HCl, pH 7.5 and 150 mM NaCl. *Lane 13*: Protein molecular marker (Biorad-Dual Color Standards). **c** Activity of SPS mutants (S495A, R496A, P497A, and D498A) compared to wild-type  $\Delta$ N-*SoSPS1* and vector pTrcHis as a control. Arrows represent the targeted proteins

predicted that this effect might be attributed to the loss of the intramolecular hydrogen bond. Replacement of Ala at R496 results in complete loss of enzyme activity (Fig. 2c). These results were in good agreement with the previously mentioned mutagenic replacement at position R496 (corresponding to R580 in *AtSuSy*; R567 in *NeSuSy*; R300 in GS) emphasizing the critical role of catalytic regulation in plant SPS. Moreover, mutation at D498 displayed a tendency to be less active compared to the wild-type enzyme (Fig. 2c). The obtained result provides evidence that D498 stabilizes R496 to interact with UDP-G through hydrogen bond interaction. On the other hand, we conducted further mutational analysis on these two charged residues to examine another possible role of R496 and D498 by replacing each with Lys and Glu, respectively. We previously presumed that the salt bridge interaction between Arg as positively charged residue and Asp as negatively charged

residue could be substituted to other potential salt bridge partners such as Lys and Glu. However, single mutation of R496K and D498E reveals an unexpected influence of these mutations without retaining the enzymatic activity (data not shown). The reason of these results is not yet clearly understood, but previous reports have suggested that these side chains play important roles to establish the structure of the enzymatic transition state and cannot be substituted to other residues by mutagenesis (Lairson et al. 2008; Wu et al. 2015).

The similarity of conserved motifs among SPS and *SuSy* suggest a possible regulation for the UDP binding mechanism. The interplay between R580 and D582 in *AtSuSy* is stronger than the interaction of Lys-Asp or Arg-Glu. The role of Asp is to maintain the mode of the UDP binding region through salt bridge interaction with Arg and to be considered as capable of binding substrate UDP-G. Thus, the predicted interaction

between residues in *SoSPS1* and UDP-G notably with the diphosphate moiety has been experimentally proven (Fig. 2c). However, our present work does not report the kinetic properties comparison between wild-type and each mutant due to the fact that some mutated SPSs become inactive.

### Substrate preference for the uridine moiety of UDP in *SoSPS1*

The characterization of residues that are responsible for binding substrate preference is apparently necessary. Nucleotide binding at the glycosyltransferase domain has been identified in sucrose synthase (Zheng et al. 2011; Wu et al. 2015) and glycogen synthase (Sheng et al. 2009) from X-ray crystal structure analysis. The crystal structure of bacterial SPS (*HoSPS*) with the substrate F6P and the product S6P have been solved. However, the *HoSPS* structure and its complex with the substrate UDP-G could not be obtained. Thus, the predicted residues which bind to UDP-G was generated by computational docking, and residue Leu at position 342 in *HoSPS* was reported as a nucleotide-binding residue (Chua et al. 2008). The structural analysis of glycogen synthase from *E. coli* revealed that the residues Y355 and H356 interact with the ADP moiety. The role of residue Y355 is to stack the adenine ring, and adenine accepts a hydrogen bond from H356 (Sheng et al. 2009). Based on the crystal structure of *AtSuSy* with the UDP-G complex, the overlap structure comparison of *AtSuSy* and *NeSuSy* could be observed. The residues that interact with the phosphate and ribose moieties are well conserved, but those interacting with nucleotide are not (Zheng et al. 2011; Wu et al. 2015; Diricks et al. 2015).

Kinetic parameters of sucrose synthase from *A. thaliana* and *N. europaea* displays different substrate preferences between UDP and ADP, respectively. Likewise, plant SPS shows substrate specificity to UDP-G, whereas the *HoSPS* shows a tendency to accept NDP-G, such as ADP-G (Chua et al. 2008). As shown in Fig. 1a, the sequence alignment implies that residues H565 and V570 in *SoSPS1* (corresponding to residues Q648 and N654 in *AtSuSy*; residues R636 and A642 in *NeSuSy*; and residues Y355 and H356 in *EcGS*) play a role in specific substrate preference. The previously reported SPS structure from *H. orenii* has shown us that residue L324, corresponding to V570 in *SoSPS1*, is predicted to interact with NDP-G, whereas the other residue corresponding to H565 in *SoSPS1* is still not clear. Although these two residues involved with nucleotide sugar are not well conserved, the sequences are considered to have a functional value which is in good agreement with other reports (Sheng et al. 2009; Zheng et al. 2011; Wu et al. 2015; Diricks et al. 2015).

In addition, subsequent site-directed mutagenesis was directed particularly at V570 to substitute Val to Ala, and our result demonstrated that V570A mutation leads to loss of SPS activity

(data not shown). This indicated that Val at 570 in *SoSPS1* is an important residue to drive the catalytic mechanism of plant SPS. The sequence alignment in previous report showed that Val residue is highly conserved in other plant SPS sequences and predicted substrate specificity for UDP-G (Chua et al. 2008). In this report, we did not compare the kinetic properties of mutants for substrate specificity between UDP-G and ADP-G. On the contrary, engineering altered substrate specificity to switch from ADP to UDP has recently been reported in bacterial sucrose synthase for industrial applications. The single, double, and triple mutations of recombinant bacterial *SuSy* were made at corresponding positions to H565 and V570 in *SoSPS1*. However, the experiment did not succeed and the substrate preference showed no change, still preferring ADP (Diricks et al. 2015).

### Perspective

Similar to sucrose synthase, plant SPS showed a strong preference for UDP-G as a substrate, as opposed to bacterial SPS, which are predicted to bind ADP-G. In plants, UDP-G and ADP-G are directed to the biosynthesis of sucrose, cellulose, and starch (Salerno and Curatti 2003; Baroja-Fernández et al. 2003). It is noteworthy that the enzymes involved in nucleotide sugar metabolism are essential to control the regulation of carbon partitioning in plants. This led to the hypothesis that one of the strategies to redesign the regulation of plant sugar metabolism is through substrate binding site engineering of plant SPS. This work is challenging since a crystal structure of plant SPS has not yet been obtained. Identifying the specific residues related to substrate nucleotide sugar by mutagenesis is proposed to overcome the lack of information about functional residues in *SoSPS1*. Following the analogy of glycosyltransferase group structures, our results have clarified that residues R496, D498, and V570 in *SoSPS1* are important for catalysis by plant SPS. Further mutational analysis is required to alter substrate preference and offer new prospects for structural arrangement accommodating UDP-G or ADP-G.

**Acknowledgements** This work was supported by Ministry of Research, Technology and Higher Education of the Republic of Indonesia (Penelitian Unggulan Strategis Nasional and Penelitian Dasar Unggulan Perguruan Tinggi) and by the International Collaborative Research Program of Institute for Protein Research, Osaka University, ICR-14-03 and 05, Japan.

### Compliance with ethical standards

**Conflict of interest** Widhi Dyah Sawitri declares that she has no conflicts of interest. Siti Nurul Afidah declares that she has no conflicts of interest. Atsushi Nakagawa declares that he has no conflicts of interest. Toshiharu Hase declares that he has no conflicts of interest. Bambang Sugiharto declares that he has no conflicts of interest.

**Ethical approval** This article does not contain any studies with human participants or animals performed by any of the authors.

## References

- Amir J, Preiss J (1982) Kinetic characterization of spinach leaf sucrose-phosphate synthase. *Plant Physiol* 69:1027–1030
- Baroja-Fernández E, Muñoz FJ, Saikusa T, Rodríguez-López M, Akazawa T, Pozueta-Romero J (2003) Sucrose synthase catalyzes the de novo production of ADPglucose linked to starch biosynthesis in heterotrophic tissues of plants. *Plant Cell Physiol* 44(5):500–509
- Baskaran S, Roach PJ, Depaoli-Roach AA, Hurley TD (2010) Structural basis for glucose-6-phosphate activation of glycogen synthase. *Proc Natl Acad Sci U S A* 107(41):17563–17568
- Breton C, Imbert A (1999) Structure/function studies of glycosyltransferases. *Curr Opin Struct Biol* 9:563–571
- Castleden CK, Aoki N, Gillespie VJ, MacRae EA, Quick WP, Buchner P, Foyer CH, Furbank RT, Lunn JE (2004) Evolution and function of the sucrose phosphate synthase gene families in wheat and other grasses. *Plant Physiol* 135:1753–1764
- Chua TK, Bujnicki JM, Tan TC, Hyunh F, Patel BK, Sivaraman J (2008) The structure of sucrose phosphate synthase from *Halothermothrix orenii* reveals its mechanism of action and binding mode. *Plant Cell* 20:1059–1072
- Coutinho PM, Deleury E, Davies GJ, Henrissat B (2003) An evolving hierarchical family classification for glycosyltransferases. *J Mol Biol* 328:307–317
- Cumino A, Curatti L, Giarrocco L, Salerno GL (2002) Sucrose metabolism: anabaena sucrose-phosphate synthase and sucrose-phosphate phosphatase define minimal functional domains shuffled during evolution. *FEBS Lett* 517:19–23
- Diricks M, Bruyn FD, Daele PV, Walmagh M, Desmet T (2015) Identification of sucrose synthase in nonphotosynthetic bacteria and characterization of the recombinant enzymes. *Appl Microbiol Biotechnol* 99:8465–8474
- Gifford RM, Thome JH, Hitz WD, Giaquinta RT (1984) Crop productivity and photoassimilate partitioning. *Science* 225:801–808
- Huber SC, Huber JL (1996) Role and regulation of sucrose-phosphate synthase in higher plants. *Annu Rev Plant Physiol Plant Mol Biol* 47:431–444
- Huber SC, Kerr PS, Ruffy TW (1985) Diurnal changes in sucrose phosphate synthase activity in leaves. *Plant Physiol* 64:81–87
- Imbert A, Wimmerova M, Koca J, Breton C (2006) Molecular modeling of glycosyltransferases. In: Brockhausen I (ed) *Methods in molecular biology*, 347: Glycobiology protocols. Humana, Totowa, pp 145–156
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
- Lairson LL, Henrissat B, Davies GJ, Withers SG (2008) Glycosyltransferases: structures, functions, and mechanisms. *Annu Rev Biochem* 77:521–555
- Leloir LF, Cardini CE (1955) The biosynthesis of sucrose phosphate. *J Biol Chem* 214:157–165
- Lunn JE, MacRae E (2003) New complexities in the synthesis of sucrose. *Curr Opin Plant Biol* 6:208–214
- Lunn JE, Price GD, Furbank RT (1999) Cloning and expression of a prokaryotic sucrose-phosphate synthase gene from the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol Biol* 40:297–305
- Pontis HG (1978) On the scent of the riddle of sucrose. *Trends Biochem Sci* 3:137–139
- Salerno GL, Curatti L (2003) Origin of sucrose metabolism in higher plants: when, how and why? *Trends Plant Sci* 8:63–69
- Sawitri WD, Narita H, Ishizaka-Ikeda E, Sugiharto B, Hase T, Nakagawa A (2016) Purification and characterization of recombinant sugarcane sucrose phosphate synthase expressed in *E. coli* and insect Sf9 cells: an importance of the N-terminal domain for an allosteric regulatory property. *J Biochem* 159(6):599–607
- Sheng F, Jia X, Yep A, Preiss J, Geiger JH (2009) The crystal structures of the open and catalytically competent closed conformation of *Escherichia coli* glycogen synthase. *J Biol Chem* 284(26):17796–17807
- Sugiharto B, Sakakibara H, Sumadi, Sugiyama T (1997) Differential expression of two genes for sucrose-phosphate synthase in sugarcane: molecular cloning of the cDNAs and comparative analysis of gene expression. *Plant Cell Physiol* 38(8):961–965
- Tarbouriech N, Chamock SJ, Davies GJ (2001) Three-dimensional structures of the Mn and mg dTDP complexes of the family GT-2 glycosyltransferase spsA: a comparison with related NDP-sugar glycosyltransferases. *J Mol Biol* 314:655–661
- Ünligil UM, Rini JM (2000) Glycosyltransferase structure and mechanism. *Curr Opin Struct Biol* 10:510–517
- Wu R, Diez MDA, Figueroa CM, Matchey M, Iglesias AA, Ballicora MA, Liu D (2015) The crystal structure of *Nitrosomonas europaea* sucrose synthase reveals critical conformational changes and insights into sucrose metabolism in prokaryotes. *J Bacteriol* 197(17):2734–2746
- Zheng Y, Anderson S, Zhang Y, Garavito RM (2011) The structure of sucrose synthase-1 from *Arabidopsis thaliana* and its functional implications. *J Biol Chem* 286(41):36108–36118

51. Review

**Molecular dynamics simulation of bacterial flagella**

Akio Kitao, Hiroaki Hata Pages 617-629

[Download PDF \(3402KB\)](#) [View Article](#)

52. Review

**The versatile mutational “repertoire” of *Escherichia coli* GroEL, a multidomain chaperonin nanomachine**

Tomohiro Mizobata, Yasushi Kawata Pages 631-640

[Download PDF \(2246KB\)](#) [View Article](#)

53. Review

**Artificial bio-nanomachines based on protein needles derived from bacteriophage T4**

Hiroshi Inaba, Takafumi Ueno Pages 641-658

[Download PDF \(6670KB\)](#) [View Article](#)

54. Review

**Hierarchical structure assembly model of rice dwarf virus particle formation**

Atsushi Nakagawa, Naoyuki Miyazaki, Akifumi Higashiura Pages 659-665

[Download PDF \(3903KB\)](#) [View Article](#)

55. Review

**Mechanism of evolution by genetic assimilation**

Ken Nishikawa, Akira R. Kinjo Pages 667-676

[Download PDF \(774KB\)](#) [View Article](#)

56. Review

**Structural atlas of dynein motors at atomic resolution**

Akiyuki Toda, Hideaki Tanaka, Genji Kurisu Pages 677-686

[Download PDF \(2903KB\)](#) [View Article](#)

