

## Jurnal Kedokteran Brawijaya

[Home](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [EDITORIAL BOARD](#)

[Home](#) > [Vol 31, No 1 \(2020\)](#)

### Jurnal Kedokteran Brawijaya

Jurnal Kedokteran Brawijaya, ISSN (online): 2338-0772, is a communication media and a scientific publication in the field of medical science published by Faculty of Medicine, Universitas Brawijaya.

Jurnal Kedokteran Brawijaya is a B accredited national journal (2014-2019) based on the Decree of the Directorate General of Strengthening Research and Development of the Ministry of Research, Technology, and Higher Education Republic of Indonesia No. 040/P/2014 dated February 14, 2014.

The accreditation status has been renewed base on the Decree of Director General on Strengthening Research and Development of the Ministry of Research, Technology, and Higher Education Republic of Indonesia 23/E/KPT/2019, August 8, 2019, accredited as SINTA 2.

Jurnal Kedokteran Brawijaya contains articles from research focused on basic medicine, clinical medicine, epidemiology, and preventive medicine (social medicine).

Jurnal Kedokteran Brawijaya is regularly published twice in one year (at the end of February and at the end of August). Jurnal Kedokteran Brawijaya was first published in April 1984.

Brawijaya Medical Journal has been indexed in [Google Scholar](#), [Portal Garuda / Indonesian Publication Index \(IPI\)](#), [SINTA Science and Technology Index](#), and [CossReff](#).

Citation Profile at [Google Scholar](#) as of January 2020:

- Number of citations: 1720
- h-index: 15
- index i10: 37

### Announcements

#### News: Changes in JKB

Become an International Language Journal

Posted: 2019-05-05

[More...](#)

[More Announcements...](#)


### Vol 31, No 1 (2020)


### Table of Contents

#### Research Article

##### **The Effects of Thyme on IL-6 Level and the Bacterial Count in the Brain and Spleen of Mice Infected by Methicillin-Resistant Staphylococcus aureus (MRSA)**

 **Lisayani Prasetyowati, Muchlis Achsan Adji Sofro, Helmia Farida, Galuh Yulieta**


 Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.1-6


 Viewed : 151 times

 PDF

##### **Analog Rice Reduces Weight and Total Cholesterol Levels in Overweight and Hypercholesterolemic Rats**

 **Septiarina Putri Dewantari, Hairrudin Hairrudin, Elly Nurus Sakinah**


 Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.7-12


 Viewed : 97 times

 PDF

##### **The Effect of Extra Virgin Olive Oil (EVOO) on Fetal Birth Weight in Preeclampsia Rat Model**

 **Yulia Silvani, Afniari Maharani, Agnestia Naning Dian Lovita**

 Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.13-18

 Viewed : 117 times

 PDF

##### **The Effect of Vitamin E Administration on Histopathological Description of Mice (Mus musculus) Liver Exposed to Mercuric Chloride**

#### About JKB

#### Aim and Scope

#### Publication Ethics

#### Visitor Statistic

#### Contact Us

#### User

Username

Password

Remember me

#### Information for Reviewer

#### Peer Review Process

#### Information for Author

#### General Information

#### Online Submissions

#### Author Guidelines

#### The Authors Data

#### Ethics Statement

#### Template

#### Tools



- [Mendeley User Guide](#)
- [Insert Citation using Mendeley](#)
- [CSL JKB](#)

#### Index



Acivrida Mega Charisma, Intan Febiola Arianing  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.19-22  
Viewed : 71 times

Untitled PDF

Analysis of Cell Block and Cytology Specimen Preservation from Lung Aspiration Biopsy

Adinda Sandya Poernomo, Willy Sandhika, Vicky Sumarki Budipramana  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.23-27  
Viewed : 75 times

PDF

Analysis of Lamina Papyracea Area and Ethmoid Sinus Volume in Indonesian Patients Undergoing Paranasal Sinus CT Scan

Wahyuni Wahyuni, Nurlaily Idris, Mirna Muis, Andi Alfian Zainuddin, Muhammad Fadjar Perkasa, Muhammad Ilyas  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.28-32  
Viewed : 108 times

PDF

Correlation between Ultrasound Findings of Uric Acid Precipitate in MTP I and Acute Gout Flare in Gouty Arthritis

Elys SUSANTI, Muhammad Ilyas, Nurlaily Idris, Andi Alfian Zainuddin, Bachtiar Murtala, Faridin HP  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.33-38  
Viewed : 77 times

PDF

Effect of Javanese Turmeric (Curcuma xanthorrhiza) Extract on Hepatitis Model of Alcohol-Induced Mice

Agnes Ilene Suprpto Puteri, Willy Sandhika, Nurina Hasanatuludhhiyah  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.39-42  
Viewed : 134 times

PDF

Relationship of TTF-1 and EGFR on Lung Adenocarcinoma at Dr. Saiful Anwar General Hospital Malang

andy lumban gaol, Suryanti Dwi Pratiwi, Ngakan Putu Putra, Eviana Norahmawati, Harun Al Rasyid  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.43-48  
Viewed : 67 times

PDF

Tulsi (Ocimum sanctum) Leaf Ethanol Extract Reduces Inflammatory Cell Infiltration in Aspirin-Induced Gastritis Rats

Festi Artika Sari, Willy Sandhika, Tri Hartini Yuliawati  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.49-52  
Viewed : 212 times

PDF

Non-pharmacological Randomised Control Trial: Green Coconut (Cocos nucifera L.) Water to Reduce Dysmenorrhea Pain

Fajar Ari Nugroho, Oktaviana Manda Putri, Yuseva Sariati  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.53-57  
Viewed : 80 times

PDF

Iron Status of Newborns and Umbilical Cord Blood Hcpidin Levels in Gender Differences

Qodri Santosa, AG Soemantri, Soeharyo Hadisaputra, Ariawan Soejoenoes  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.58-65  
Viewed : 88 times

PDF

Effects of Pulmonary Vascular Resistance Index on Oxygen Saturation in Patients with Atrial Septal Defect

Supomo Supomo  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.66-68  
Viewed : 88 times

PDF

Plasma Glucagon-Like Peptide-1 and Cholecystokinin Responses to Fast Food in Healthy-Weight and Obese Men

Dian Handayani, Dianita Setya Pradita Putri, Hidayat Sujuti, Sri Andarini, Widya Rahmawati, Inggita Kusumastuty, Leny Budhi Harti, Nindy Sabrina  
Jurnal Kedokteran Brawijaya, Vol 31, No 1 (2020), pp.69-75  
Viewed : 90 times

PDF

Notifications

- View
- Subscribe

Journal Content

Search

Search Scope

All

Search

Browse

- By Issue
- By Author
- By Title

Information

- For Readers
- For Authors
- For Librarians

Visitor Counter

ID	156755	US	18195
SG	433	IN	430
MY	405	GB	153
NL	113	TR	104
EU	88	GR	83
Newest:	NZ You:		ID
Today:			9
Month:			370
Total:	178048		
Supercounters.com			

Keywords

Aging Apoptosis Aterosklerosis BDNF Candida albicans Diabetes MDA Salmonella typhi  
TNF-α apoptosis cedera otak  
traumatik diabetes melitus inflammatory cells  
kehamilan kolesterol malondialdehid obesitas  
pengetahuan polimorfisme gen ACTN3  
streptozotocin tes kepekaan antibiotik

Current Issue

ATOM 1.0  
RSS 2.0  
RSS 1.0



## Research Article

### **Analog Rice Reduces Weight and Total Cholesterol Levels in Overweight and Hypercholesterolemic Rats**

#### **Beras Analog Menurunkan Berat Badan dan Kadar Kolesterol Total pada Tikus *Overweight* dan Hiperkolesterolemia**

Septiarina Putri D<sup>1</sup>, Hairrudin<sup>2</sup>, Elly Nurus S<sup>3</sup>

<sup>1</sup>Medical Study Program Faculty of Medicine Universitas Jember Jember

<sup>2</sup>Department of Biochemistry Faculty of Medicine Universitas Jember Jember

<sup>3</sup>Department of Pharmacology Faculty of Medicine Universitas Jember Jember

#### **ABSTRACT**

*Overweight, whose prevalence is increasing, is caused by unbalanced consumption patterns such as high consumption of fat and can trigger hypercholesterolemia. Analog rice (AR), which has better nutritional content than ordinary rice, is thought to have beneficial effects on health. This study aimed to determine the potential of analog rice in reducing body weight and total cholesterol levels. A total of 24 male Wistar rats were divided into four groups, namely: I) control, II) ordinary rice treatment, III) analog rice 1 treatment (PBA1) and IV) analog rice 2 treatment (PBA2). The control group was given standard feed for 40 days ad libitum. Groups II, III, and IV were given a combination induction of high-fat diet (HFD) for 40 days ad libitum and intraperitoneal injection of streptozotocin (STZ) 35 mg/kg single dose on day 33. All groups were then given the treatment of diet for three weeks according to their respective groups, namely the standard, ordinary rice, analog rice 1, and analog rice 2 feed. Body weight and total cholesterol levels were measured in the pre-induction, post-induction/pre-treatment (pre-test), and post-treatment (post-test). The results of the study showed that the induction of the HFD-STZ combination carried out resulted in a significant increase in body weight and total cholesterol levels. Measurements made after dietary therapy showed that body weight and total cholesterol levels in the I and II groups did not change significantly, whereas those in the III and IV groups decreased significantly ( $p < 0.05$ ). This study concluded that the administration of analog rice could reduce body weight and total cholesterol levels in overweight and hypercholesterolemia rats.*

**Keywords:** Analog rice, high-fat diet, hypercholesterolemia, overweight, streptozotocin

#### **ABSTRAK**

*Overweight yang prevalensinya semakin meningkat diakibatkan oleh pola konsumsi tidak seimbang, seperti konsumsi tinggi lemak, dapat memicu hiperkolesterol. Beras analog (BA) yang memiliki kandungan gizi lebih baik dari beras biasa, diperkirakan memiliki efek yang menguntungkan bagi kesehatan. Penelitian ini bertujuan untuk mengetahui potensi BA dalam menurunkan berat badan (BB) dan kadar kolesterol total. Sebanyak 24 ekor tikus Wistar jantan dibagi menjadi 4 kelompok, yaitu: I) kontrol, II) perlakuan beras biasa (PBB), III) perlakuan beras analog 1 (PBA1), dan IV) perlakuan beras analog 2 (PBA2). Kelompok kontrol diberi pakan standar selama 40 hari ad libitum. Kelompok II, III, dan IV diberi induksi kombinasi berupa high-fat diet (HFD) selama 40 hari ad libitum dan injeksi streptozotocin (STZ) intraperitoneal 35 mg/kgBB dosis tunggal pada hari ke-33. Semua tikus selanjutnya diberi perlakuan pemberian pakan/diet selama 3 minggu sesuai dengan kelompoknya masing-masing, berturut-turut yaitu pakan standar, beras biasa, BA1 dan BA2. Berat badan dan kadar kolesterol total diukur pra-induksi, pasca-induksi/ pra-perlakuan (pre-test), dan pasca-perlakuan (post-test). Hasil penelitian menunjukkan induksi kombinasi HFD-STZ yang dilakukan mengakibatkan peningkatan BB dan kadar kolesterol total yang bermakna. Pengukuran yang dilakukan pasca terapi diet menunjukkan BB dan kadar kolesterol total kelompok I dan II tidak mengalami perubahan bermakna, sedangkan kelompok III dan IV mengalami penurunan yang bermakna ( $p < 0,05$ ). Kesimpulan pada penelitian adalah pemberian BA mampu menurunkan BB dan kadar kolesterol total pada pada tikus yang mengalami overweight dan hiperkolesterolemia.*

**Kata Kunci:** Beras analog, high-fat diet, hiperkolesterolemia, overweight, streptozotocin

---

Correspondence: Hairrudin. Department of Biochemistry Faculty of Medicine Universitas Jember, Jl. Kalimantan No. 37, Jember Tel. 08123200364 Email: [hairrudin.fk@unej.ac.id](mailto:hairrudin.fk@unej.ac.id)

DOI: <http://dx.doi.org/10.21776/ub.jkb.2020.031.01.2>

## INTRODUCTION

Overweight is a condition of body weight that exceeds the normal limit. The prevalence is increasing both in developed and developing countries such as Indonesia. In 2016, more than 1.9 billion adults aged 18 and over suffered from overweight. Unhealthy eating patterns, such as the tendency to consume foods high in fat, high in calories, low in fiber, are one of the triggers for increased overweight and obesity among teenagers and young adults. Overweight can develop into obesity and trigger hypercholesterolemia (1-3). The prevalence of hypercholesterolemia among Indonesian inhabitants over 15 years is 35.9% (4).

Analog rice is a rice-grain-like processed product that can be made from partially or wholly non-rice ingredients, and the nutrition can be modified (5,6). The content of food fiber (7,8) and Resistant starch (RS) (8,9) of AR (analog rice) is generally higher than ordinary rice. Food fiber, which can be grouped into soluble fiber and insoluble fiber, is a residue from plant cell walls composed of carbohydrates, consumable, and resistant to the process of digestion and absorption in the human small intestine. Food rich in fiber can reduce body weight and total cholesterol levels (2), while RS is starch, and its degradation products cannot be absorbed in the small intestine (10). High RS levels in the diet can increase fatty acid oxidation, and trigger satiety, so consumption of a diet rich in RS can be used to treat obesity (11).

Several studies regarding the potential of AR in several medical conditions have been carried out. Corn analog rice (Smart Rice) is able to provide hypoglycemic effects in diabetes mellitus patients with an average of 24.2 percent after a three-week consumption (7). Analog Rice with main compositions of sago starch, fresh cassava, coconut pulp, and rice bran given to Sprague Dawley rats for four weeks was able to reduce total cholesterol and Low-Density Lipoprotein (LDL) levels and increase High-Density Lipoprotein (HDL) levels compared to the group control (12). Administration of AR made from a mixture of sago starch and red bean flour for four weeks caused a reduction in total cholesterol in prediabetes patients (13). This study used AR containing the main raw materials of MOCAF flour, corn flour, rice flour, and sodium alginate, with two variations in the composition of raw materials (BA1 and BA2). The calorie content in both types of AR was made equal to ordinary rice by regulating the composition of raw materials.

## METHOD

### Experimental Animals

This study was a pure experiment using a pre-test post-test control group design. The study used 24 white male rats (*Rattus norvegicus*), Wistar strain,  $\pm$  three months old, and 150-200 grams body weight. Rats were obtained from the Biochemical Laboratory of the Faculty of Medicine, Airlangga University. Ethical Clearance research was obtained from the Ethics Commission of the Faculty of Medicine, University of Jember through a decree number 1176/H25.1.11/KE/2018. After the rats were adapted for seven days, the rats were randomly divided into four groups (each group consisted of 6), namely: I) control group, II) ordinary rice treatment (PBB), III) analog rice treatment 1 (PBA1), and IV) analog rice treatment 2

(PBA2). Every rat was placed in a cage with the same shape, size, and material. During the adaptation period, mice got standard feed and distilled water ad libitum.

### Overweight and Hypercholesterolemia Induction

The induction of overweight and hypercholesterolemia was carried out for the treatment groups (II, III, and IV) with an induction combination of HFD and STZ injection (14-16). The high-fat diet provided contained 22.8% pork fat (table 1) and was administered for 40 days, while STZ was given intraperitoneally once on the 33<sup>rd</sup> day calculated from the first day of HFD administration. The STZ dose used was a low dose, which was 35 mg/kg BW (8,14). Induction of a combination of low-dose HFD and STZ has been proven to cause mice to experience hyperglycemia, weight gain (14-16), hypercholesterolemia, and hypertriglyceridemia (14,15). STZ was dissolved in citric buffer with a concentration of 0.05 M and pH 4.5 (15,16). Confirmation of overweight and hypercholesterolemia was done after induction, after the rats were first fasted for 12 hours. Cholesterol levels were measured through blood obtained from the tails of the rats and were measured using a digital cholesterol level meter (15). Rats were classified as hypercholesterolemia if they have total cholesterol levels >111 mg/dl (11). Rats that gained weight over the mean of +2SD of the control group were considered overweight. Overweight and hypercholesterolemia rats obtained were used as subjects in this study by providing dietary therapy for three weeks.

Table 1. Composition of pellets per 100 grams

Component	Standard pellet	HFD pellet	Rice pellet	BA1 pellet	BA2 pellet
Standard pellet <sup>1</sup>	100	77,2	60	60	60
Lard	0	22,8	0	0	0
Ordinary rice	0	0	40	0	0
BA1	0	0	0	40	0
BA2	0	0	0	0	40
Total	100	100	100	100	100

Note: 1=standard pellet containing 60% carbohydrate, 23% fish flour, 6% soybeans and other additives, HFD=High fat diet, BA1=analog rice 1, BA2=analog rice 2

### Analog Rice and Pellet Preparation

Analog rice in this study was artificial rice made utilizing a hot extruder using Subagio Technique (7) with the main raw materials of MOCAF, corn flour, rice flour, and an addition of sodium alginate with two different composition variations. The composition was listed based on a formula from one of the Smart Rice variants modified by Hairrudin et al. (Table 2). Analog rice 1 had less MOCAF content but more rice flour, while analog rice 2 had more MOCAF content but less rice flour content (8).

Table 2. Calorie content and analog rice composition per 100g

Ingredients	Ordinary rice	BA1	BA2
MOCAF	0	27	54
Rice flour	100	36	9
Corn flour	0	27	27
Soy Protein	0	6,5	6,5
Palm Oil	0	2,3	3,5



**Table 2. Calorie content and analog rice composition per 100g (Continued)**

Ingredients	Ordinary rice	BA1	BA2
Sodium alginate	0	1,2	0
Total	100	100	100

**Note:** MOCAF = Modified cassava flour, ordinary rice, BA1 = analog rice 1, BA2 = analog rice 2. Sources: (8)

The calorie content, both BA1 and BA2, was made equal to ordinary rice. Analog rice had a fiber content of 1.77% in analog rice formula 1, 0.86% in analog rice formula 2, and 0.41% in ordinary rice. The fiber and RS content of ordinary rice, BA1, and BA2 can be seen in Table 3. Regular rice and AR were then used as pellets by mixing them with standard pellets with a ratio of 60% ordinary rice or AR and 40% standard pellets.

**Table 3. Nutrient content in rice, BA1, and Ba2**

Group	Fiber	RS
Ordinary rice	0,41	0,01
BA1	1,77	16,83
BA2	0,86	11,46

**Note:** RS = Resistant starch, BA1 = analog rice 1, BA2 = analog rice 2

#### Measurement of Body Weight and Total Cholesterol Levels

Weight measurements were carried out three times that were during the pre-induction (initial), post-induction/pre-treatment diet (pre-test), and post 3-week treatment diet (post-test). Weight measurements were carried out using a digital scale (Camry). Measurement of total cholesterol levels was done with an electrode-based biosensor method using a digital cholesterol gauge or cholesterol control check (Easy Touch) and blood cholesterol test strip. Measurement of total cholesterol levels was carried out twice, post-induction/pre-treatment diet (pre-test) and post 3-week treatment diet (post-test). Before the blood was obtained, the rats were fasted overnight (12 hours). Blood collection was done through the tail cut  $\pm 0.5$  cm. Blood drops were inserted into the cholesterol test strip that was attached to a digital cholesterol gauge to calculate the total cholesterol levels.

#### Statistical Analysis

The data obtained were summarized and presented in a tabular form, and the statistical analysis was carried out using IBM SPSS version 21 with a confidence level of 95% ( $\alpha = 0.05$ ). Data normality distribution was assessed using the Shapiro-Wilk, while data homogeneity was assessed using the Levene test. The One-Way ANOVA test with Bonferroni posthoc was carried out on initial body weight and pre-test as well as pre-test total cholesterol levels to determine the success of the combination induction. The Paired T-test was carried out on body weight and total cholesterol levels of pre-test and post-test to see the difference between before and after the 3-week diet treatment to know the success of treatment with analog rice.

## RESULTS

This study aimed to determine the potential of analog rice

with the main components of MOCAF flour, corn flour, rice flour, and sodium alginate in reducing body weight and total cholesterol levels in overweight rats with hypercholesterolemia. The rats were obtained by induction of HFD-STZ combination. After the induction, the body weight and total cholesterol levels of the rats increased. The next stage was giving intervention to the rats in the form of diet therapy for three weeks. A summary of the research results can be seen in Table 4 and 5.

#### Body Weight

The results of animal body weight measurements during the study can be seen in Table 4. Body weight was measured three times that were in pre-induction (initial), pre-test, and post-test. The mean body weight of the treatment rats (group II, III, and IV combined) at the beginning of the study was  $160.28 \pm 11.51$  gram. After the HFD-STZ induction, the mean body weight of the rats increased significantly to  $208.00 \pm 26.03$  gram, while the weight of the control group did not increase significantly. The One-Way ANOVA test results on all body weight groups after induction (pre-test) obtained  $p = 0.002$ , which means that there were significant differences in mean body weight between groups after HFD-STZ induction. Body weight differences among the groups were tested using the Bonferroni posthoc test. The Bonferroni posthoc test results for BW after induction among the induced groups (II, III, and IV) showed no significant differences, whereas among all the induced groups and group I (control) there were significant differences (I vs. II  $p = 0.008$ ; I vs. III  $p = 0.007$ ; I vs. IV  $p = 0.013$ ). The results of the analysis showed that body weight between the control group and all HFD-STZ-induced groups experienced significant differences, whereas between the treatment groups, there were no significant differences. These facts prove that the induction of the HFD-STZ combination carried out increased the body weight.

**Table 4. Mean of the weight measurement result**

Group	Initial BW (Mean $\pm$ SD)	Pre-Test BW (Mean $\pm$ SD)	Post-Test BW (Mean $\pm$ SD)
I (Control)	161,33 $\pm$ 11,15	152,83 $\pm$ 20,13	154,17 $\pm$ 19,82
II (PBB)	159,00 $\pm$ 10,16	208,83 $\pm$ 30,71*	192,00 $\pm$ 34,92
III (PBA1)	161,17 $\pm$ 11,81	209,83 $\pm$ 15,99**	190,17 $\pm$ 09,83*
IV (PBA2)	160,67 $\pm$ 14,31	205,33 $\pm$ 33,03*	186,67 $\pm$ 41,70**

**Note:** BW=body weight, SD=standard deviation, PBB=ordinary rice treatment, PBA1=analog rice treatment 1, PBA2=analog rice treatment 2, \*=significant difference between initial BW and pre-Test, \*\*=significant difference between BW pre-Test and pre-Test

After the diet treatment for three weeks, group II, III, and IV experienced a significant decrease in body weight, while group I (control) experienced an increase but not significant. The changes in body weight from each group before and after the diet were analyzed using the Paired T-test. The test results showed a significant decrease in weight in group III ( $p = 0.003$ ) and group IV ( $p = 0.007$ ), but not significant in group II ( $p = 0.350$ ), while group I actually experienced an increase in weight but not significant ( $p = 0.401$ ). This showed that the provision of analog rice could significantly reduce body weight while the provision of ordinary rice did not reduce body weight in overweight rats. The difference was due to differences in fiber content and RS. The greater the fiber and RS content in the feed

consumed, the greater the effectiveness of weight loss.

#### Total Cholesterol Levels

Total cholesterol levels in normal Wistar rats is  $103.8 \pm 7.2$  mg/dl (17). This fact shows that the highest normal total cholesterol levels is 111 mg/dl. In this study, the normal value of total cholesterol levels used the average cholesterol level of the control group, which was 116 mg/dl. Rats in the treatment groups (II, III, and IV) had hypercholesterolemia since they had total cholesterol levels higher than 116 mg/dl. The results of measurements of total cholesterol in rats during the study can be seen in Table 5. Total cholesterol levels were measured twice before and after being given the diet. Differences in cholesterol levels per group before and after diet (pre-test and post-test) were analyzed using the Paired T-test. The test results showed that total cholesterol levels in group III and group IV had a significant decrease with the same p-value (0.045). The level of total cholesterol in group II increased but was not significant ( $p = 0.622$ ), while the total cholesterol levels in group I did not experience a significant change ( $p = 0.849$ ). This showed that the provision of analog rice could significantly reduce total cholesterol levels. The fact is because AR has high fiber and RS content, the greater the fiber and RS content in the diet consumed, the greater the effectiveness of decreasing total cholesterol levels.

**Table 5. Mean of the total cholesterol measurement result**

Group	Pre-Test cholesterol (Mean $\pm$ SD)	Post-Test cholesterol (Mean $\pm$ SD)
I (Control)	116,00 $\pm$ 08,63	116,83 $\pm$ 06,24
II (PBB)	145,17 $\pm$ 28,74*	151,67 $\pm$ 12,71
III (PBA1)	176,33 $\pm$ 30,81*	142,33 $\pm$ 28,48**
IV PBA2)	152,00 $\pm$ 24,20*	120,83 $\pm$ 07,14**

**Note:** SD=standard deviation, PBB=ordinary rice treatment, PBA1=analog rice treatment 1, PBA2=analog rice treatment 2, \*=significant difference between initial and pre-Test BW, \*\*=significant difference between pre-Test and pre-test BW

## DISCUSSION

In this study, the induction of HFD-STZ combination performed was proven to significantly increase the body weight and total cholesterol levels in white rats (*Rattus norvegicus*). Overweight conditions in rats are caused by excess fat accumulation in the body due to high-fat consumption in HFD (15,17). This study used HFD that contained 22.8% lard. The administration of HFD results in increased absorption of triacylglycerol (TG) so that the levels in lipoproteins increase. This condition results in TG hydrolysis by increased lipoprotein lipases in the capillaries. The hydrolysis releases free fatty acids so that the levels in the blood increase (17,18). Some of the free fatty acids are re-formed into TG and stored in adipocytes (17). Prolonged administration of HFD results in more TG deposits resulting in overweight (18). Overweight can lead to obesity and is one of the triggers of insulin resistance that can trigger hypercholesterolemia. The combination of intraperitoneal HFD and low dose STZ injection has been shown to increase insulin resistance (8), thus causing glucose metabolic disorders marked by hyperglycemia (14-16), overweight (14,16), and hypercholesterolemia (15-16). Repeated administration of HFD results in

increased accumulation of fat in adipose tissue and increased levels of free fatty acids and fatty acid oxidation. Fatty acid oxidation increases the production of acetyl Co-A (18). It stimulates endogenous cholesterol synthesis or triggers excessive increases in total cholesterol in the blood resulting in hypercholesterolemia (15-16). One way to handle overweight and hypercholesterolemia is to consume a diet that contains high fiber (2,12,19,20) and RS.

The results of this study prove that diet intervention in the form of analog rice for three weeks can reduce body weight and total cholesterol levels in male Wistar strain rats that are overweight and hypercholesterolemia due to the induction of the HFD-STZ combination. After the 3-week diet, group III (PBA1) experienced body weight decrease by 19.67 grams, and group IV (PBA2) by 18.67 grams. Group II (PBB) did not experience a significant decrease in weight, while group I (control) experienced an increase in weight but not significant, which was as much as 1.33 grams. The mean total cholesterol levels of group III (PBA1) experienced a significant decrease of 34 mg/dl and group IV (PBA2) of 31.17 mg/dl. The mean total cholesterol levels in group II (PBB) increased but not significant as much as 6.5 mg/dl, while group I (control) experienced a decrease in total cholesterol but not significant as much as 0.83 mg/dl. The decrease in body weight and total cholesterol is inversely proportional to the fiber content contained in the diet. This shows that the higher the fiber content in the feed given, the greater the effect on weight loss and total cholesterol.

The food fiber content in analog rice plays an important role. c flour and MOCAF flour are alternative carbohydrate sources that have a higher fiber content compared to ordinary rice. Sodium alginate added to BA1 plays an important role in increasing its fiber content. MOCAF has a high crude fiber content of 3.4% to 12% (21,22). Corn flour is one source of food fiber in the form of soluble and water-insoluble fiber (23), while sodium alginate contains water-soluble fiber (24).

Water-soluble fiber triggers gel formation in the digestive tract that functions to slow gastric emptying and maintain satiety. Fiber-rich foods can support the peptide release in the form of cholecystokinin that is involved in the regulation of gastric emptying and stimulate the nucleus of fullness in the hypothalamus. Satiety induced by dietary fiber shows the potential for sustainable weight loss without hyperphagia and weight re-bound in obesity cases (2,25,26).

Fiber can bind with fatty acid salts in the small intestine and is released and become a food source of microflora in the colon through the fermentation process. The microflora produces fermented products from fibers in the form of Short Chain Fatty Acids (SCFA), such as propionic acid (8). Propionic acid has activity as an inhibitor of the enzyme 3-Hydroxy-3-methylglutaryl CoA (HMG CoA) reductase, thus inhibiting cholesterol synthesis. Insoluble fiber also has the effect of increasing the expression of Glucose Transporter Type 4 (GLUT-4), resulting in improved insulin sensitivity or reduced insulin resistance. Decreased insulin resistance results in the inhibition of cholesterol synthesis so that the hypercholesterolemia condition can be prevented (2).

The RS content in AR also has an important role in reducing body weight and total cholesterol levels. The RS content in AR is quite high, while rice contains almost no RS (Table 3). Hot-extrusion technology used in the AR manufacturing



process resulted in the starch in AR undergoing gelatinization and retrogradation (7). Retrogradation occurs because the technology goes through the stages of heating and cooling which facilitate amylose retrogradation. Retrogradation produces RS so that the level increase (10,27). Sodium alginate contains water-soluble fiber (24) which triggers gel formation in the gastrointestinal tract, thereby increasing RS production. This results in the RS level in BA1 becomes higher than BA2 because sodium alginate is added in BA1 composition.

High RS level causes inhibited digestion and absorption of food in the intestine (11), resulting in the synthesis of fatty acids, lipogenesis, free fatty acids, and decreased energy storage pathways which in turn causes decreasing energy storage in the form of TG in adipocytes, so body weight also decreases. The reduction in free fatty acids subsequently results in decreased fatty acid oxidation and reduced acetyl-CoA production. Acetyl-CoA is the raw material for cholesterol synthesis in the body. A decrease in acetyl-CoA production results in a decrease in total cholesterol levels (11,18).

This study concludes that the provision of analog rice 1 and analog rice 2 can significantly reduce body weight and total cholesterol levels in overweight and hypercholesterolemic rats. Analog rice 1 has a higher fiber and RS content than analog rice 2, so it gives a better effect on weight loss and total cholesterol levels. This shows the potential of analog rice made from MOCAF, corn flour, rice flour, and the addition of sodium alginate as an alternative solution for dietary therapy that can be utilized in the health sector, especially in overweight and hypercholesterolemia.

#### ACKNOWLEDGMENT

Our gratitude goes to all parties who have assisted in the compilation of this research article. Thank you to Prof. Ir. Achmad Subagio, M.Agr., Ph.D., who has assisted the process of analog rice production and the Integrated Laboratory of FTP University of Jember that has assisted the analysis process. We would also like to thank the members of research team, Fadiah Ulfa Khairina, Monika Roosyidah, Prajesiaji P., Azka Darajat, and Khanif Muflikhatun who contributed to the research.

#### REFERENCES

- World Health Organization. *Obesity and Overweight*. (Online) 2018. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> [accessed 23 Januari 2019].
- Fairudz A and Nisa K. *Pengaruh Serat Pangan terhadap Kadar Kolesterol Penderita Overweight*. Majority. 2015;4(8): 121-126.
- Septiana P, Nugroho FA, and Wilujeng CS. *Konsumsi Junk Food dan Serat Pada Remaja Putri Overweight dan Obesitas Yang Indekos*. Jurnal Kedokteran Brawijaya. 2018; 30(1): 61-67.
- Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia. *Riset Kesehatan Dasar*. Jakarta: Departemen Kesehatan RI; 2013.
- Budijanto S and Yuliyanti. *Study of Preparation Sorghum Flour and Application for Analogues Rice Production*. Jurnal Teknologi Pertanian. 2012; 13(3): 177-186.
- Yuwono SS and Zulfiah AA. *Formulasi Beras Analog Berbasis Tepung Mocaf dan Maizena dengan Penambahan CMC dan Tepung Ampas Tahu*. Jurnal Pangan dan Agroindustri. 2015;3(4):1465-1472.
- Subagio A, Witono Y, Hermanuadi D, Nafi A, and Windrati WS. *Pengembangan "Beras Cerdas" Sebagai Pangan Pokok Alternatif Berbahan Baku Mocaf*. Prosiding InSINas. Jember, November 29-30, 2012: 157-160.
- Hairrudin, Soetjipto, and Handajani R. *Hypoglycemic Effect of High-Resistant Starch Analog Rice Through GLP-1 and Insulin or High-Resistant Starch Analog Rice Attenuates Blood Glucose Level Through Enhancement of GLP-1 and Insulin*. Asian Journal Pharmaceutical and Clinical Reserch. 2019; 12(8): 172-175
- Noviasari S, Kusnandar F, Setiyono A, and Budijanto S. *Beras Analog sebagai Pangan Fungsional dengan Indeks Glikemik Rendah*. Jurnal Gizi Pangan. 2015; 10(3): 225-232.
- Zaman SA and Sarbini SR. *The Potential of Resistant Starch as a Prebiotic*. Critical Reviews in Biotechnology. 2016; 36(3): 578-584.
- Gentile CL, Ward E, Holst JJ, et al. *Resistant Starch and Protein Intake Enhances Fat Oxidation and Feelings of Fullness in Lean and Overweight/Obese Women*. Nutrition Journal. 2015; 14(113): 1-10.
- Kharisma T. *Studi Hipokolesterolemik Beras Analog Secara In Vivo pada Tikus Sprague Dawley (SD)*. [Tesis]. Institut Pertanian Bogor, Bogor. 2015.
- Hariyanto B, Cahyana PT, Putranto AT, Wahyuningsih SB, and Marsono Y. *Penggunaan Beras Sagu Untuk Penderita Pradiabetes*. Jurnal Pangan. 2017; 26(2): 1-10.
- Srinivasan K, Viswanad B, Asrat L, Kaul CL, and Ramarao P. *Combination of High-Fat Diet-Fed and Low-Dose Streptozotocin-Treated Rat: A Model for Type 2 Diabetes and Pharmacological Screening*. Pharmacological Research. 2005; 52(4): 313-320.
- Arumugam S and Natesan S. *Hypoglycemic Effects Of Barleria Noctiflora Fractions On High Fat Fed With Low Dose Streptozotocin Induced Type-2 Diabetes In Rats*. International Journal of Pharmacy and Pharmaceutical Sciences. 2016; 8(2): 193-200.
- Lu Y, Liu Y, Li H, Wang X, Wu W, and Gao L. *Effect and Mechanisms of Zinc Supplementation in Protecting Against Diabetic Cardiomyopathy in a Rat Model of Type 2 Diabetes*. Bosnian Journal of Basic Medical Sciences. 2015; 15(1): 14-20.
- Marques C, Meireles M, Norberto S, et al. *High-fat Diet-induced Obesity Rat model: a comparison between Wistar and Sprague-Dawley Rat*. Adipocyte. 2016; 5(1): 11-21.
- Rodwell VW, Bender DA, Botham KM, Kennelly PJ, and Weil PA. *Harper's Illustrated Biochemistry*. 30th ed, New York: McGraw-Hill Education; 2015; 168-

- 185, 223-226.
19. Harsa IMS. *Efek Pemberian Diet Tinggi Lemak terhadap Profil Lemak Darah Tikus Putih (Rattus norvegicus)*. Jurnal "Ilmiah Kedokteran". 2014; 3(1): 21-28.
  20. Wiardani NK, Sugiani PPS, and Gumala NMY. *Konsumsi Lemak Total, Lemak Jenuh, dan Kolesterol sebagai Faktor Resiko Sindroma Metabolik pada Masyarakat Perkotaan di Denpasar*. Jurnal Gizi Klinik Indonesia. 2011; 7(3): 107-114.
  21. Sunarsi S, Sugeng M, Wahyuni S, and Ratnaningsih W. *Memanfaatkan Singkong Menjadi Tepung Mocaf Untuk Pemberdayaan Masyarakat Sumberejo*. Sukoharjo; LPPM Univet Bantara Sukoharjo; 2011.
  22. Lopulalan CGC, Mailoa M, and Pelu H. *Analisa Sifat Kimia dan Fisik Modified Cassava Flour (Mocaf) (Varietas Lokal Sangkola) Asal Desa Waai, Maluku Tengah*. AGRITEKNO: Jurnal Teknologi Pertanian. 2018; 5(1): 7-12.
  23. Suarni and Yasin M. *Jagung sebagai Sumber Pangan Fungsional*. Iptek Tanaman Pangan. 2011; 6(1): 41-56.
  24. Salleh SN, Fairus AAH, Zahary M, Bhaskar Raj N, and Mhd Jalil AM. *Unravelling The Effects of Soluble Dietary Fibre Supplementation on Energy Intake and Perceived Satiety in Healthy Adults: Evidence from Systematic Review and Meta-Analysis of Randomised-Controlled Trials*. Foods. 2019; 8(1): 15.
  25. Bozzetto L, Costabile G, Della Pepa G, et al. *Dietary Fibre as a Unifying Remedy for the Whole Spectrum of Obesity-Associated Cardiovascular Risk*. Nutrients. 2018; 10(7): 943-975.
  26. Adam CL, Thomson LM, Williams PA, and Ross AW. *Soluble Fermentable Dietary Fibre (Pectin) Decreases Caloric Intake, Adiposity and Lipidaemia in High-Fat Diet-Induced Obese Rats*. PLoS ONE. 2015; 10(10): 1-14.
  27. Firdaus J, Sulistyarningsih E, Subagio A. *Resistant Starch Modified Cassava Flour (MOCAF) Improves Insulin Resistance*. Asian Journal of Clinical Nutrition. 2018; 10(1): 32-36.

