



NURSE & HEALTH

NURSE AND HEALTH: JURNAL KEPERAWATAN

- THE EFFECT OF HEALTH PROMOTION ABOUT DISASTER MANAGEMENT TOWARDS THE LEVEL OF KNOWLEDGE OF YOUTH RED CROSS MEMBERS
- HOME VISIT FOR APPLYING FAMILY NURSING: A PERSPECTIVE
- EFFECTIVENESS OF BACK EFFLUERAGE MASSAGE THERAPY USING CLOVE OIL ON UNDER BACK PAIN IN POSTPARTUM MOTHER
- THE EFFECTIVENESS OF INTRADIALYSIS STATIC BIKE EXERCISE TOWARD FUNCTIONAL CAPACITY IN HEMODIALYSIS PATIENTS
- THE EFFECT OF WALKING EXERCISE ON FATIGUE IN CANCER PATIENTS WHO WALK IN CHEMOTHERAPY AT RSUP Dr. M. DJAMIL PADANG
- THE INFLUENCE OF KNOWLEDGE AND DISASTER WARNING AND THE PREPARATION OF HIGH SCHOOL STUDENTS IN FACING EARTHQUAKE AND TSUNAMI
- COUNTERPRESSURE FOR DYSMENORRHEA PAIN IN TEENAGERS
- THE EFFECT OF CLINICAL TEACHING METHOD AND LEARNING STYLE ON CLINICAL NURSING LEARNING OUTCOMES
- THE EFFECT OF HEALTH EDUCATION THROUGH SESSION-HEALTH APPLICATION MEDIA ON BEHAVIOR MANAGEMENT BEHAVIOR OF DIABETES MELLITUS IN BENGKULU CITY
- EFFECT OF ALOE VERA EXTRACT AND DISCHARGE PLANNING TO ACCELERATE WOUND HEALING OF OLDER ADULT HERNIA SURGERY PATIENTS
- THE EFFECT OF INFRARED RAY AND COUNSELING ON DIABETIC FOOT ULCER HEALING PROCESS
- THE RELATIONSHIP OF HEALTHY LIFESTYLE INDEX (HLI) TO THE OCCURRENCE OF HYPERTENSION IN MOUNTAINS, COASTAL AND URBAN COMMUNITIES
- THE EFFECT OF ASSERTIVE COMMUNITY TREATMENT TOWARDS ANXIETY AND DEPRESSION ON BREAST CANCER CLIENT



LPPM AKADEMI KEPERAWATAN KERTA CENDEKIA SIDOARJO

P-ISSN : 2088-9909



NURSE & HEALTH

NURSE AND HEALTH: JURNAL KEPERAWATAN

Vol. 8 No. 2, July-December 2019

Editorial Team

Editor-in-Chief

Agus Sulistyowati, S.Kep., M.Kes, Maternity Nursing Department, Kerta Condokia Nursing Academy

Managing Editor

Kusuma Wijaya Ridi Putra, S.Kep.Ns., MNS, Community Nursing Department, Kerta Condokia Nursing Academy | WoS Researcher ID X-1011-2019

Receiving Editor

Faida Annisa, S.Kep.Ns., MNS, Psychiatric and Mental Health Nursing Department, Kerta Condokia Nursing Academy

Associate Editor, Indonesia

Moli Diana, S.Kep.Ns., M.Kes, Medical Surgical Nursing Department, Kerta Condokia Nursing Academy

Riesmiyatiningdyah, S.Kep.Ns., M.Kes, Pediatric Nursing Department, Kerta Condokia Nursing Academy

Editorial Board Member

Elok Triostuning, S.Psi., M.Si, Kerta Condokia Nursing Academy

Dini Prasetyo Wijayanil, S.Kep.Ns., M.Kep, Emergency Nursing Department, Kerta Condokia Nursing Academy | Scopus ID 57201402405 | h-index 1

Marlita Dewi Lestari, S.Kep.Ns., M.Kes, Nursing Management Department, Kerta Condokia Nursing Academy

Journal Manager

Kusuma Wijaya Ridi Putra, S.Kep.Ns., MNS, Community Nursing Department, Kerta Condokia Nursing Academy | WoS Researcher ID X-1911-2019

Reviewer

Assoc. Prof. Dr. Chintana Watcharasin, Pediatric Nursing Department, Burapha University, Thailand

Assist. Prof. Dr. Pornchai Jullamate, Gerontology and Nursing Sciences Department, Burapha University, Thailand | Scopus ID 11640574200 | h-index 5

Assist. Prof. Dr. Yupin Tanatwanitch, Adult Nursing Department, Burapha University, Thailand

Assist. Prof. Dr. Jinjuta Chalsona Dallas, Psychiatric and Mental Health Nursing Department, Burapha University, Thailand | Associate Dean for International Affairs

Assist. Prof. Dr. Chanandchidadussadee Toonsiri, Community Nursing Department, Faculty of Nursing, Burapha University

Hem Kumar Nepal, RN, MNS, Clinical Nurse Quality Management Service in Mongar Regional Referral Hospital

Suveksha Shrestha, RN, MNS, Clinical Practice Manager in Gurkha Welfare Trust Nepal

Dr. Pipit Festi Wilyanarti, S.Kep.Ns., M.Kes, Faculty of Health Sciences, Muhammadiyah University, Surabaya | Scopus ID 57106955475 | h-index 1

Candra Panji Asmoro, S.Kep.Ns., M.Kep (Dr. Candidate), Faculty of Nursing, Airlangga University | Scopus ID 57204562356 | h-index 1

Rista Fauziningtyas, S.Kep.Ns., M.Kep, Faculty of Nursing, Airlangga University | Scopus ID 57204880986

Feri Ekapraselia, S.Kep.Ns., M.Kep, STIKES dr. Soebandi Jember | Emergency Nursing | Scopus ID 57203873645

Ika Nur Pratiwi, S.Kep.Ns., M.Kep, Faculty of Nursing, Airlangga University | Scopus ID 57202753063

Alamat Redaksi (Editorial Address):

Kampus Akademi Keperawatan Kerta Condokia Sidoarjo

Jalan Lingkar Timur, Rangkah Kidul, Sidoarjo, 61234

Telp. (031) 8961496, Fax. (031) 8961497

E-mail: jurnal.nurcandhealth.2017@gmail.com

Focus and Scope

"Nurse and Health: Jurnal Keperawatan" is a Scientific Journal managed by Community Service and Research Division of Kerta Cendekia Nursing Academy. The scope of this journal is articles in the fields of health, public health, and nursing (basic nursing, medical surgical nursing, pediatric nursing, maternity nursing, community health nursing, family nursing, emergency nursing, psychiatric and mental health nursing, gerontological nursing, and also nursing management). This journal is published in print and online every 6 months, namely June and December.

Section Policies

Editorial

Open Submissions Indexed Peer Reviewed

Guest Editorial

Open Submissions Indexed Peer Reviewed

Letter to the Editor

Open Submissions Indexed Peer Reviewed

Concept Analysis

Open Submissions Indexed Peer Reviewed

Review Article

Open Submissions Indexed Peer Reviewed

Original Research: Research Methodology Paper

Open Submissions Indexed Peer Reviewed

Original Research Article

Open Submissions Indexed Peer Reviewed

Perspective

Open Submissions Indexed Peer Reviewed

Case Study

Open Submissions

Indexed

Peer Reviewed

Public

Nurse and Health: Jurnal Keperawatan, a research journal in the field of nursing.

To maintain the journal's quality, all articles are peer-reviewed.

The journal is published by the Faculty of Nursing, Universitas Jember.

The journal is published twice a year.

Editor

The journal is published by the Faculty of Nursing, Universitas Jember.

The journal is published by the Faculty of Nursing, Universitas Jember.

The journal is published by the Faculty of Nursing, Universitas Jember.

The journal is published by the Faculty of Nursing, Universitas Jember.

R

P

H

J

K

E

R

.

.

.

.

.

.

.

.

.

.

.

Peer Review Process

All papers submitted to Nurse and Health: Jurnal Keperawatan undergo double blind and external peer review. In the double-blind review process neither the reviewers nor the authors know the identity of each other. This enables unbiased and accurate review of the manuscript. In all cases, reviewers will be asked to declare any conflict of interest based on the contents of the manuscript.

After a manuscript is submitted, it is reviewed by a member of the Editorial Board. If the manuscript passes the editorial review, it is sent to external reviewers for double-blind peer review.

The reviewers are selected on the basis of their area of expertise and interests, if the reviewer accepts the offer to review the manuscript, they will be sent the complete manuscript and a Manuscript Review Form.

In all cases, reviewers will be asked to declare any conflict of interest based on the contents of the manuscript. If a conflict of interest exists, the reviewers are requested to decline to review the manuscript.

Please provide an objective critical assessment of the manuscript about the concept of the study, relevance in relation to current scientific knowledge, scientific content, language, and grammar. You will be asked to make a recommendation for publishing the manuscript. Please provide reasons for your recommendations.

If you believe that the manuscript needs changes for improvement before it is accepted for publication, please make the suggestions on how to improve it. If the comments are negative please help the authors in improving their manuscript by explaining weaknesses in scientific content or language. Any offensive language in the comments cannot be tolerated.

Based on the reviewer's comments the assigned editor will take a decision about the acceptance or rejection of the manuscript. The editor may decide to: 1) accept the manuscript without revisions, 2) invite authors to resubmit the manuscript after minor or major revisions while the final decision is kept pending, or 3) reject the manuscript.

Publication Frequency

Nurse and Health: Jurnal Keperawatan publishes one volume per year with two issues per volume. It is on June and December.

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

Archiving

This journal utilizes the LOCKSS system to create a distributed archiving system among participating libraries and permits those libraries to create permanent archives of the journal for purposes of preservation and restoration. [More](#)

Publication Ethics

Nurse and Health : Jurnal Keperawatan is a peer-reviewed journal to publish articles in the field of nursing science, research, and management. *Nurse and Health : Jurnal Keperawatan* committed to the ethical standards set for all entities involved in the publication of article.

To maintain high ethical standards of publication of quality science the publisher strives to work closely at all times with journal editors, authors and peer-reviewers. The ethics statement for *Nurse and Health : Jurnal Keperawatan* is based on those by the Committee on Publication Ethics (COPE) Code of Conduct guidelines available at www.publicationethics.org.

The work described in your article must have been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans and EC Directive 86/609/EEC for animal experiment.

The essentials of *Nurse and Health : Jurnal Keperawatan* publishing ethics for all groups involved in the publishing process are as follows:

Editors' Responsibilities

- The editor of a journal holds a vital position taking important editorial decisions on all peer-reviewed submitted for publication.
- The editor should maintain the transparency of the academic research & record, preclude professional needs from cooperating ethical standards, and always be willing to publish retractions, rectifications, and erratum when required.
- The editor should assess manuscripts for their scientific quality and intellectual content, free from any sort of biased decisions based on discrimination of race, gender, geographical origin, or religion of the author(s). The editor should evaluate manuscripts objectively based on their academic merit free of any commercial or self-interests.
- The editor should not disclose any information on submitted manuscripts before publication of the manuscript.
- Promoting research rectitude must be preserved. If at any stage the publisher suspects any kind of misconduct in research, it should be investigated promptly in detail with suitable authority; and if any suspicious act of misconduct is observed in the peer review, it should be resolved with diligence.

Reviewers' Responsibilities

- Providing a detailed, constructive, and unbiased evaluation in a timely manner on the scientific content of the work.
- Indicating whether the writing is relevant, concise & clear and evaluating the originality and scientific accuracy.
- Maintaining the confidentiality of the complete review process.
- Notifying the journal editor about any financial or personal conflict of interest and declining to review the manuscript when a possibility of such a conflict exists.
- Notifying the journal editor of any ethical concerns in their evaluation of submitted manuscripts; such as any violation of ethical treatment of animal or human subjects or any considerable similarity between the previously published article and any reviewed manuscript.

Authors' Responsibilities

- All the work reported in the manuscript must be original and free from any kind of plagiarism.
- The work should not have been published elsewhere or submitted to any other journal(s) at the same time.
- Any potential conflict of interest must be clearly acknowledged.
- Proper acknowledgements to other work reported (individual/company/institution) must be given. Permission must be obtained from any content used from other sources.
- Only those who have made any substantial contribution to the interpretation or composition of the submitted work, should be listed as 'Authors'. While other contributors should be mentioned as 'co-authors'.

Publishers' Responsibilities

- *Nurse and Health : Jurnal Keperawatan* is committed to work with journal editors, defining clearly their relevant roles, in order to ensure appropriate decisions regarding publication procedures and maintaining the transparency of editorial decisions.

- *Nurse and Health : Jurnal Keperawatan* ensures the integrity autonomy and originality of each published article with respect to:
 - publication and research fundings
 - publication ethics and recludite
 - conflicts of interest
 - confidentiality
 - authorship
 - article modifications
 - timely publication of content.

A

NI
CO

PL

TR
PT
E
ar
b

E

T
P

Copyright Form

Download [Copyright Form](#)

Journal Title: _____

Manuscript Title: _____

I/we certify that I/we have participated sufficiently in the intellectual content, conception and design of this work or the analysis and interpretation of the data (when applicable), as well as the writing of the manuscript, to take public responsibility for it and have agreed to have my/our name listed as a contributor. I/we believe the manuscript represents valid work. Each author confirms they meet the criteria for authorship as established by the *Nurse and Health : Jurnal Keperawatan*. Neither this manuscript nor one with substantially similar content under my/our authorship has been published or is being considered for publication elsewhere, except as described in the covering letter. I/we certify that all the data collected during the study is presented in this manuscript and no data from the study has been or will be published separately. I/we attest that, if requested by the editors, I/we will provide the data/information or will cooperate fully in obtaining and providing the data/information on which the manuscript is based, for examination by the editors or their assignees. Financial interests, direct or indirect, that exist or may be perceived to exist for individual contributors in connection with the content of this paper have been disclosed in the cover letter. Sources of outside support of the project are named in the cover letter.

I/We hereby transfer(s), assign(s), or otherwise convey(s) all copyright ownership, including any and all rights incidental thereto, exclusively to the Journal, in the event that such work is published by the Journal. The Journal shall own the work, including 1) copyright; 2) the right to grant permission to republish the article in whole or in part, with or without fee; 3) the right to produce preprints or reprints and translate into languages other than English for sale or free distribution; and 4) the right to republish the work in a collection of articles in any other mechanical or electronic format.

We give the rights to the corresponding author to make necessary changes as per the request of the journal, do the rest of the correspondence on our behalf and he/she will act as the guarantor for the manuscript on our behalf.

All persons who have made substantial contributions to the work reported in the manuscript, but who are not contributors, are named in the Acknowledgment and have given me/us their written permission to be named. If I/we do not include an Acknowledgment that means I/we have not received substantial contributions from non-contributors and no contributor has been omitted.

Name	Signature	Date signed
1 _____	_____	_____
2 _____	_____	_____
3 _____	_____	_____
4 _____	_____	_____
5 _____	_____	_____

6 _____ (Additional signatures may be added provided the authors meet the *Nurse and Health : Jurnal Keperawatan* criteria stated above)



NURSE AND HEALTH: JURNAL KEPERAWATAN

TABLE OF CONTENTS
DAFTAR ISI

ARTIKEL (ARTICLES)	HALAMAN (PAGES)
1. THE EFFECT OF HEALTH PROMOTION ABOUT DISASTER MANAGEMENT TOWARDS THE LEVEL OF KNOWLEDGE OF YOUTH RED CROSS MEMBERS Loren Juksen, Ida Rahmawati, Qrestha Utami	62-68
2. HOME VISIT FOR APPLYING FAMILY NURSING: A PERSPECTIVE Kusuma Wijaya Ridi Putra	69-71
3. EFFECTIVENESS OF BACK EFFLUERAGE MASSAGE THERAPY USING CLOVE OIL ON UNDER BACK PAIN IN POSTPARTUM MOTHER Rini Rahmayanti, Yola Yolanda	72-77
4. THE EFFECTIVENESS OF INTRADIALYSIS STATIC BIKE EXERCISE TOWARD FUNCTIONAL CAPACITY IN HEMODIALYSIS PATIENTS Ria Desnita, Vivi Syofia Sapardi	78-82
5. THE EFFECT OF WALKING EXERCISE ON FATIGUE IN CANCER PATIENTS WHO WALK IN CHEMOTHERAPY AT RSUP Dr. M. DJAMIL PADANG Weny Amelia, Mira Andika	83-90
6. THE INFLUENCE OF KNOWLEDGE AND DISASTER WARNING AND THE PREPARATION OF HIGH SCHOOL STUDENTS IN FACING EARTHQUAKE AND TSUNAMI Ruri Maiseptya Sari, Hanifah Hanifah, Vike Pebri Giena	91-100

- | | | |
|--|----------------|---|
| <p>7. COUNTERPRESSURE FOR DYSMENORRHEA PAIN IN TEENAGERS
Retty Nirmala Santiasari, Siska Christianingsih</p> | <p>101-107</p> | <p>DOI: http://jnsn.ujember.ac.id
December
Accepted
http://ejournal.ujember.ac.id</p> |
| <p>8. THE EFFECT OF CLINICAL TEACHING METHOD AND LEARNING STYLE ON CLINICAL NURSING LEARNING OUTCOMES
Ida Zuhroidah, Nurul Huda, Bagus Dwi Cahyono</p> | <p>108-114</p> | <p>© 2019
<u>ORIGINAL</u></p> |
| <p>9. THE EFFECT OF HEALTH EDUCATION THROUGH SESSION-HEALTH APPLICATION MEDIA ON BEHAVIOR MANAGEMENT BEHAVIOR OF DIABETES MELITUS IN BENGKULU CITY
Hanifah Hanifah, Vike Pebri Giena, Ruri Maisetya Sari</p> | <p>115-123</p> | <p></p> |
| <p>10. THE EFFECT OF ALOE VERA EXTRACT AND DISCHARGE PLANNING TO ACCELERATE WOUND HEALING OF OLDER ADULT HERNIA SURGERY PATIENTS
Mukhammad Toha, Mokh Sujarwadi, Ida Zuhroidah</p> | <p>124-133</p> | <p></p> |
| <p>11. THE EFFECT OF INFRARED RAY AND COUNSELING ON DIABETIC FOOT ULCER HEALING PROCESS
Mokh Sujarwadi, Mukhammad Toha, Nurul Huda</p> | <p>134-139</p> | <p></p> |
| <p>12. THE RELATIONSHIP OF HEALTHY LIFESTYLE INDEX (HLI) TO THE OCCURRENCE OF HYPERTENSION IN MOUNTAINS, COASTAL, AND URBAN COMMUNITIES
Iis Noventi, Yanis Kartini</p> | <p>140-152</p> | <p></p> |
| <p>13. THE EFFECT OF ASSERTIVE COMMUNITY TREATMENT TOWARDS ANXIETY AND DEPRESSION ON BREAST CANCER CLIENT
Andikawati Fitriyasari, Ahmad Yusuf, Ririn Probowati</p> | <p>153-159</p> | <p></p> |

DOI: <https://doi.org/10.36720/nhjk.v8i2.121>

Toha, M., Sujarwadi, M., & Zuhroidah, I. Nurse and Health: Jurnal Keperawatan. 2019 July-December; 8 (2): 124-133

Accepted: December 3, 2019

<http://ejournal-kertacendekia.id/index.php/nhjk/>

© 2019 Nurse and Health: Jurnal Keperawatan

ORIGINAL RESEARCH

E-ISSN: 2623-2448 | P-ISSN: 2088-9909

THE EFFECT OF ALOE VERA EXTRACT AND DISCHARGE PLANNING TO ACCELERATE WOUND HEALING OF OLDER ADULT HERNIA SURGERY PATIENTS

Mukhammad Toha ^{1*}, Mokh Sujarwadi ¹, Ida Zuhroidah ¹

¹ Jember University, Pasuruan Campus

***Correspondence:**

Mukhammad Toha

Email: toha.akper@unej.ac.id

Address: Jember University, Pasuruan Campus, Jl. KH. Mansyur No.207, Tembokrejo, Kec. Purworejo, Pasuruan City, East Java Province 67118

ABSTRACT

Background: Older adult patients who underwent hernia surgery commonly get obstacles in the process of wound healing as a result of decline of body functions. Aloe vera which contains of anti-inflammatory, anti-microbial and skin fibroblasts stimulator, can be used as adjunctive therapy in treating wounds.

Objective: This study aimed to identify the effect of aloe vera extract and discharge planning to accelerate wound healing of older adult hernia surgery patients at dr. R. Soedarsono General Hospital, Pasuruan.

Method: This study uses Quasi-Experimental with Static-group comparison approach. The population is all post-operative hernia patients aged 55-65 years at the operating room of dr. R. Soedarsono General Hospital, Pasuruan. The sample consists of 20 respondents taken according to the inclusion criteria. The independent variable is the treatment of wounds using aloe vera extract accompanied by discharge planning, while the dependent variable was the acceleration of wound healing especially inflammatory and proliferation phases. The data were collected through observation of macroscopic current wound care. Data were analyzed using nonparametric Mann Whitney test with significance level of ≤ 0.05 .

Result: The results showed the effect of the treatment using aloe vera extract and discharge planning to accelerate wound healing of older adult hernia surgery patients, the inflammatory phase of healing occurs on the sixth day for inflammation, exudation, and edema ($p = 0.022$, $p = 0.028$, and $p = 0.029$, respectively). The proliferative phase occurs on day three with $p = 0.015$.

Conclusion: This research indicates further research on the benefits of aloe vera and discharge planning for wound healing especially in older adult patients is required. Thus, a stronger basis for recommending aloe vera extract as complementary therapy for wound care is available.

Key words: Aloe Vera, Discharge Planning, Surgical Wound, Older Adults Age.

INTRODUCTION

Wound is an injury experienced by humans. Wound is defined as losing epithelial integrity from skin (Schwartz et al, 2010). Smeltzer dan Bare (2010) state that wound can be distinguished by contamination consisting of clean wound, contaminated clean wound, and dirty wound. Clean wound generally comes from surgery

process, which has passed incision procedure prepared aseptically. Its healing takes fast because there is no infection.

At Dr. R. Soedarsono General Hospital, Pasuruan, wound care procedures performed using a solution of NaCl 0.9%. However, older adults who had hernia surgery takes days four wound healing. At the tenth day after surgery, the wound is

still moist. The wound edges are not fully connected. It is painful and inflamed. Those conditions are not suitable with a theory stated that at the 7-10 day, the surgical suture can be released and the wound has been in good condition. (Mansjoer, 2014).

The unhealed wound will trigger some risks, such as bleeding, infection, dehiscence, and eviscerations (Mansjoer, 2014). The problem in wound treatment after hernia surgery of older adults can be special attention in nursing. In caring patients, nursing profession is demanded to develop the independent and safe nursing intervention. Exploring natural stuffs containing chemical substance as antiseptic and micronutrient is developed since it is able to help wound healing process. It is included to complementary nursing. This effort is proper to Regulations of Nursing Practical Implementation stating that nursing practice is aimed for individual, family, group, and society through promotive preventive activities, social empowerment, and complementary nursing actions (Depkes RI, 2010).

Aloe vera (*Aloe vera linn*) contains many substances which are useful to precipitate wound healing by (1) serving essential micro-nutrient; (2) anti-inflammatory effect; (3) anti-microbial effect; and (4) stimulating skin fibroblasts. Aloe vera also gives positive effects for collagen balance which has an important role in wound healing. Nevertheless, the effects of aloe vera for hernia wound healing process of older adults.

Wound healing process in hernia surgery of older adults needs more intensive care. It is not only at the hospital, but also at home. Thus, health promotion needs to be prepared, both for clients and family who will take care of the patient. Discharge planning is a process of preparing patients before they go back to home. The process identifies the needs and plans for facilitating the continuity of treatment either at home or another health service unit

(Kozier, 2008). Discharge planning for patients after surgery is not implemented well in dr. R. Soedarsono General Hospital. The information given is only controlling time and instruction of taking medicine. Most discharge planning is used in resume of patients.

Data from Operating Rooms of dr. R. Soedarsono General Hospital reveal that in 2014 hernia was in the first position (209 cases). In 2015, that case increased to 330 cases, while in 2016 it decreased into 255 cases. For case distribution based on age, there were 138 (54%) patients at the age of 55 years and more. 117 patients (46%) were at the age of less than 55 years old. The data proved that most of its patients were older adults. Based on the medical records from Surgery Unit of dr. R. Soedarsono General Hospital in 2015, the surgical suture in less than 55 years old patients were released at the tenth day. Meanwhile, for older patients, the surgical sutures were put off at the fifteenth day. The releasing process of surgical suture is executed when the wound has been dry and its edges united, so that surgical suture is not necessary.

Wound healing process is divided into three main phases, these are: 1) hemostasis and inflammation, 2) proliferation, 3) maturity and remodelling. These phases are overlapping and they happen since the wound comes out until healing process. (Wilksman *et al*, 2007). Inflammation phase is the first phase of wound healing process. At this condition, the wound is reddish and inflamed. The body increases its temperature and the pain comes. This step takes 3-4 days after injury. This healing process means cleaning process of the rests. Inflammation response causes the artery seeps and releases plasma and PMN (polymorphonucleocytes) around the tissue. Neutrophils along with local cells phagocytose the rests and microorganisms.

Proliferation phase starts at the fourth day after injury and takes until the 24th day. Clinically, this phase is marked by red tissue which replaces dermal tissue. After

that, fibroblasts secrete the collagen for regenerating cells along with angiogenesis and epithelization. In the last process of epithelization, contracture happens where keratinocytes differentiate to form outside protective layer or stratum korneum. After the basic structures completed, finishing interior starts. Dermal tissue increases its power by fibroblasts. Maturation phase or remodelling takes one year or more after injury (Aminuddin, 2009).

There are many technics for treating wounds, one of them is applying antiseptic substances which are often used for treating wounds. These antiseptic stuffs have antimicrobial substances which can fight microorganism in wounds. However, it also has its side effect, these are irritated skin, skin color change, and scar tissue because scar tissue is due to foreign object during the wound healing process (Kurniawan, 2007). Because of the side effects from chemical substances, the wound treatment uses normal saline solvent. This solvent is easy to absorb by the body, thus it will not create disadvantages.

Aloe vera is plantable and easy to find everywhere. In some countries, people use aloe vera as the first aid of treating wound. Aloe vera contains active substances which are useful for precipitating in healing wound. It contains glucomanan, acemannan saponin, lignin, vitamin A, vitamin C, enzymes and amino acid for regenerating cells. Aloe vera stimulates epidermic, increase fibroblasts function, and form the new blood vessels (Furnawanthi, 2006).

This study aimed to identify the effect of aloe vera extract and discharge planning to accelerate wound healing of older adult hernia surgery patients at dr. R. Soedarsono General Hospital, Pasuruan.

METHODS

Study Design

This research uses Quasi Experimental design, with static-group comparison design approach.

Setting

This research was conducted at the dr. R. Soedarsono General Hospital, Pasuruan on February to March 2017.

Research Subject

The population for objects is entire patients of hernia surgery at the age of 55-65 years old in surgery room of dr. R. Soedarsono General Hospital, Pasuruan. The samples consist of 20 respondents with inclusion criteria, namely male gender, had albumin levels of 3.5-4.5 g%, Hb levels of 10-14 g%, GDA 85-125 mg%, using consecutive sampling method. The independent variable in the treatment group here is wound treating process using topical aloe vera extract after cleansing with Na Cl 0.9% along with discharge planning, while in the control group the surgical wound treating without topical aloe vera extract was given. Moreover, the dependent variable is precipitation of wound healing after hernia surgery in inflammation and proliferation phases.

Instruments

The instruments used are observation reports about wound healing after hernia surgery. These are inflammatory phase including inflammation, edema, and exudation. Meanwhile, proliferation phase includes measuring uniting wound edges. The measurement of wound healing is done macroscopically in the third and sixth day after surgery without pretest.

Data Analysis

The collected data are analyzed using Mann Whitney test with ≤ 0.05 for meaning level compared to controlling and measuring group in each measuring session.

Ethical Consideration

This research has gone obtained permission from the director of the dr. R. Soedarsono General Hospital, Pasuruan

with number license of Permit Letter: 800/734/423.212/2017.

RESULTS

Examination of the Effect of Aloe Vera Extract and Discharge Planning to Accelerate Wound Healing of Older Adult Hernia Surgery Patients at dr. R. Soedarsono General Hospital, Pasuruan

Table 1. Examination of the Effect of Aloe Vera Extract and Discharge Planning to Accelerate Wound Healing (Inflammatory Phase) of Older Adult Hernia Surgery Patients on Sixth Days After Surgery at dr. R. Soedarsono General Hospital, Pasuruan by using Mann Whitney Test (n = 20).

Component of Inflammatory	Group				Asymp. Sig.
	Control		Treatment		
	N	%	N	%	
Inflammation					0.022
> 2 cm	0	0	0	0	
0.6-2 cm	6	60	1	10	
No	4	40	9	90	
Total	10	100	10	100	
Edema					0.029
> 2 cm	0	0	0	0	
0.6-2 cm	4	40	0	0	
No	6	60	10	100	
Total	10	100	10	100	
Exudation					0.028
Unclear	0	0	0	0	
Clear	8	80	3	30	
No	2	20	7	70	
Total	10	100	10	100	

Source: Primary data of questionnaire, 2017

Based on table 1, It was found that the use of Aloe Vera Extract significantly had an impact on the inflammatory process, both inflammation, edema, and exudation ($p = 0.022$, $p = 0.029$, $p = 0.028$, respectively).

Table 2. Examination of the Effect of Aloe Vera Extract and Discharge Planning to Accelerate Wound Healing (Proliferation Phase) of Older Adult Hernia Surgery Patients on Third Days After Surgery at dr. R. Soedarsono General Hospital, Pasuruan by using Mann Whitney Test (n = 20).

Component of Proliferation Phase	Group				Asymp. Sig.
	Control		Treatment		
	N	%	N	%	
Uniting Wound Healing					0.015
Not Uniting	4	40	0	0	
Some	6	60	8	80	
Entire	0	0	2	20	
Total	10	100	10	100	

Source: Primary data of questionnaire, 2017

The results on the table 2 found that the use of Aloe Vera Extract on day 3rd had a significant impact on the proliferation phase (uniting wound healing) ($p = 0.015$).

DISCUSSION

The results of this study found that the use of Aloe Vera Extract significantly had an impact on the inflammatory process, both inflammation, edema, and exudation ($p = 0.022$, $p = 0.029$, $p = 0.028$, respectively). Inflammation is a response toward lesion in living tissue which has vascularization (Kumar, 2008). Vascular and cellular response happen when the tissue is cut or injured. Vasoconstriction of blood vessel happens and clotted fibro platelet is formed to control the bleeding. This reaction takes five until ten minutes and is followed by vasodilation of venular. Microcirculation loses its vasoconstriction ability because norepinephrine is ruined by intracellular enzyme. Histamine is released to increase permeability of capillary (Smeltzer dan Bare, 2010).

When microcirculation breaks, blood elements such as antibody, protein plasma, electrolyte, complement, and water come into vascular spatial in two until three days. It creates edema, increased temperature,

redness, and pain. The first thing happening is inflammation. When inflammatory reaction appears, arteriolar enlarges. Hence, much more blood flows into local microcirculation. The stretched capillary is filled by the blood. This condition is hyperemia or congestion triggering redness in severe inflammation. The change of local pH or local concentration of certain ions can stimulate tip of nerves. The release histamine or other bioactive substances can stimulate nerves. Some of inflammation is caused by hyperemia, and most of it is triggered by liquid supply and cells from blood circulation into interstitial tissue.

Exudation is the extra vascular inflammatory liquid, with high density (more than 1.020). It contains 2-4 mg% protein with emigrating leukocyte. This liquid is heaped up as the impact of vascular permeability increase, more intravascular hydrostatics as the impact of increased local blood circulation, and any other complicated things causing emigrating leukocyte (Kumar, 2008).

The inflammation healing phase using aloe vera extract and discharge planning on the third day does not result well compared to controlling method. It is related to the age factor, that is elderly will take more time for healing process. In elderly, Langerhans cells reduce and macrophage becomes less active, hence they reduce immune activity within the skin. Moreover, the less perspiration process makes the skin dry, so the healing process takes more time. It is proved when in the third day, the wound is still 60% reddish and has 30% edema. Besides, 70% patients still have moist wound. According to Gaylene and Patricia (2000), one of the factors obscuring wound healing is about age.

Extension of inflammation phase, basically occurs in two groups. Nevertheless, the group which uses aloe vera and discharge planning for treating wound takes shorter period in healing the wound. Probably it is due to saponin content in aloe vera. Saponin is able to be

antiseptic and cleanser to fight against germs and bacteria. According to Cowser (2010) and Kathuria (2011), ace Annan has anti-inflammatory effect by reducing bradykinin and histamine activities. Meanwhile, enzymes bradykinase and peptidase are able to construct bradykinins as anti-inflammation without toxicity effect. Furthermore, using Na Cl 0.9% as cleansing in wound treatment keeps the wound moist. The moist condition only keeps the situation optimal for granulation growth, hence redness, inflammation, and exudation continue.

Discharge planning is a method to help patient reaching optimal health level by promoting health to patient and family. One of the promotion methods is informing of how important early mobilization and its steps. The comparison of patient independence level for treatment group and controlling group shows that there is no significant difference ($p=0.137$). However, descriptively it shows that independence level of treatment group is higher. The early mobilization for patients after surgery can help the blood circulation runs well. Therefore, inflammatory phase can be minimized.

The effect of aloe vera in inflammation healing process is supported by a research by Akhoondinasab et.al (2014). They said that applying 300 mg/kg *mannose-6-phosphate* for mouse is able to accelerate wound healing process compared to giving sodium chloride 0.9% by controlling group. On the other hand, Hunter et.al (2006) stated that chromones 8-*C-glucosyl-(2'-O-cinnamoyl)-7-O-methylaloeol* A given 200 micrograms to mouse can help equally with giving 200 micrograms hydrocortisone and does not reduce thymus gland size. Anti-inflammation is also resulted from brady kinase, carboxypeptidase enzyme and salicylic acid contained in aloe vera.

The research result for proliferation phase shows differences between wound treatment using aloe vera along with

discharge planning, and controlling group. Level of wound edges uniting on the third, day after surgery shows significance value $p < 0.05$ ($p = 0.015$). It is the proof that treating wound using aloe vera along with discharge planning significantly affects toward wound edges uniting since the third day.

Proliferation phase is the continuing of inflammation phase. If inflammation phase is successful, it will be followed by proliferation phase. Fibroblasts multiply themselves and form the tissues for migrating cells. Epithelia cells form bud on the wound edges. The bud will bloom into capillary which is nutrition source for the new granule tissue. In this phase, fiber is formed and destroyed to adapt with wound stretching. (Sjamsuhidajat dan Jong, 2011).

Surgical wound treatment using NaCl 0.9% is aimed to keep skin's humidity that the wound will be healed soon. However, for elderly, they have low ability of absorption, thus the mass of liquid reduces 2.5% especially in dermal layer. Fibroblasts producing collagen and elastin decrease. The less Langerhans cells and less active macrophage reduce dermal immune activity. The dermal blood vessel becomes thicker and less permeable. The adipose fat tissue loses, so the migrating process of cells to surface takes longer period.

Aloe vera contains lignin which can increase skin absorbing ability. The research from Dal'Beló et.al (2006) about effect of aloe vera for skin hydration showed that aloe vera is a natural stuff which can hydrate skin effectively. According to Laursen and Nielson (2000) in Ramamorty (2011), in wound healing process, glucomannan completes by lectin way where mannose sugar component stimulates secretion of mannose binding lectin protein (MBL). It is tied with carbohydrate of bacteria wall. This bond acitifies 3b complement functioned as chemotaxis and opsonization, hence macrophage becomes more active in eradicating the germs. Macrophage will

release *Epidermal Growth Factor* (EGF), *Transforming Growth Factor β 1* (TGF), *Fibroblast Growth Factor* (FGF) dan *Interleukins* (IL-1) besides other cytokines. (Perdanakusuma, 2007). Ramamorty (2011) and Cowsert (2010) stated that Acemannan actifies macrophage and increase nitric oxide synthase in transcription level followed by increasing of *TGF- β* , *EGF*, *FGF*. They stimulate proliferation, cellular motion, contractility, differentiation, and angiogenesis which are crucial in wound healing process. The hypothesis is supported by a research from Atiba, et al (2011) stating that aloe vera can increase wound healing process. Aloe vera works through junctional intercellular stimulating communication and proliferation fibroblasts. Minerals and vitamins, especially vitamin C in aloe vera help fibroblast in forcing collagen.

Type I and III collagen are the primary dermal component which are important in proliferation. In wound healing process after hernia surgery, granulation tissue is not visible and scar tissue forming is minimal. It happens because the wound is tight, sterile, and minimum for tissue disorder. This wound healing process runs well and fast. Basically, the wound healing process in inflammation and proliferation phase occur in two groups. However, the treatment using aloe vera extract along with discharge planning runs more fast compared to the group using Na Cl 0.9%. Consequently, the wound healing process happens primarily. The result of this research proves that aloe vera along with discharge planning work well in wound healing process after hernia surgery in older adults.

CONCLUSION

The wound healing process after hernia surgery in older adults takes shorter period by using aloe vera extract along with discharge planning. It takes more fast compared to controlling group. Healing process of inflammation happens seriously

on the sixth day, while proliferation phase occurs on the third day.

SUGGESTIONS

Applying 25% aloe vera extract on the wound after hernia surgery is recommended because it significantly speeds up the wound healing process, both in inflammation and proliferation phase. *Discharge planning* can affect significantly toward level of depression after surgery, so that it can speed up wound healing process. The next researcher should modify nursing service in which respondent can stay and get treatment every day. Aloe vera extract is applied daily hence the phytochemical effect will be more optimal. The next researcher should study the effect of aloe vera for pain, inconvenience, psychological acceptance and financial matters.

ACKNOWLEDGMENT

The authors would like to thanks to Director of Pasuruan Nursing Academy as permit of this research, and to dr. R. Soedarsono General Hospital, Pasuruan as a research site.

DECLARATION OF CONFLICTING INTEREST

None.

FUNDING

The funds used in this study came from the researchers' personal funds.

AUTHOR CONTRIBUTION

Mukhammad Toha: Conduct research as the research leader and compile manuscripts.

Mokh Sujarwadi: Conduct research.

Ida Zuhroidah: Conduct research.

ORCID

Mukhammad Toha

<https://orcid.org/0000-0002-2648-0626>

Mokh Sujarwadi

<https://orcid.org/0000-0002-3818-0671>

Ida Zuhroidah

<https://orcid.org/0000-0002-4544-0099>

REFERENCES

- Afandi, M.F. (2011). *Kemampuan Penyembuhan Luka dengan Lidah Buaya*. Retrieved from <http://kesehatan.kompasiana.com/alter-natif/2011/01/01/kemampuan-penyembuhan-luka-pada-aloevera>.
- Akhoondinasab, M.R., Akhoondinasab, M., Saberi, M. (2014). Comparison of healing effect of aloe vera extract and silver sulfadiazine in burn injuries in experimental rat model. *World journal of plastic surgery*, 3(1), 29.
- Aminuddin. (2009). Prinsip Dasar Penyembuhan Luka. Retrieved from <http://aminetn.wordpress.com/2009/07/26/prinsip-dasar-penyembuhan-luka/>
- Atiba, A., Nishimura, M., Kakinuma, S., Hiraoka, T., Goryo, M., Shimada, Y., Uzuka, Y. (2011). Aloe vera oral administration accelerates acute radiation-delayed wound healing by stimulating transforming growth factor- β and fibroblast growth factor production. *The American Journal of Surgery*, 201(6), 809-18.
- Atik, N. dan Iwan J. (2009). *Perbedaan Efek Pemberian Topikal Gel Lidah Buaya (Aloe vera L.) dengan Solusio Povidone Iodine terhadap Penyembuhan Luka Sayat pada Kulit Mencit (Mus musculus)*. MKB. 2009:8:2:87-93.
- Banu, A., Sathyanarayana, B., & Chattannavar, G. (2012). Efficacy of fresh Aloe vera gel against multi-drug resistant bacteria in infected leg ulcers.

- The Australasian medical journal*, 5(6), 305.
- Chithra, P., Sajithlal, G.B., & Chandrakasan, G. (2008). Influence of Aloe Vera on Collagen Characteristics in healing Dermal Wounds in Rats. *Mol-Cell-Biochem*, 181(1-2), 71-76.
- Cowsert, Lex, M. (2010). Biological Activities of Acemannan. Retrieved from <http://www.symmetrydirect.com/pdf/Bioactivofacemannan.pdf>.
- Choonhakarn, C., Busaracome, P., Sripanidkulchai, B., Sarakam, P. (2010). A Prospective, randomized clinical trial comparing topical aloe vera with 0.1% triamcinolone acetonide in mild to moderate plaque psoriasis. *Journal Eur Acad Dermatol Venereol*, 24(2), 168-72
- Dal' Belo, S.E., Gaspar, L.R., Maia, Campos, P.M. (2006). Moisturizing effect of cosmetic formulations containing Aloe vera extract in different concentrations assessed by skin bioengineering techniques. *Skin Research Technol*, 12(4), 2441-6.
- Das, S., Mishra, B., Gill, K., Ashraf, M.S., Singh, A.K., Sinha, M., Singh, T.P. (2011). Isolation and characterization of novel protein with anti-fungal and anti-inflammatory properties from Aloe vera leaf gel. *International Journal of Biological Macromolecules*, 48(1), 38-43.
- Departemen Kesehatan RI. (2010). *Izin dan Penyelenggaraan Praktik Perawat*, Permenkes republic Indonesia No HK.02.02/Menkes/148/2010. Pasal 8.
- Egesie, U., Chima, K., & Galam, N. (2011). Anti-inflammatory and analgesic effects of aqueous extract of Aloe Vera (Aloe barbadensis) in rats. *African Journal of Biomedical Research*, 14(3), 209-12.
- Eshghi, F., Hosseinimehr, S.J., Rahmani, N., Khademloo, M., Norozi, M.S., & Hojati, O. (2010). Effects of Aloe vera cream on posthemorrhoidectomy pain and wound healing: results of a randomized, blind, placebo-control study. *The Journal of Alternative and Complementary Medicine*, 16(6), 647-50.
- Furnawanthi, I. (2006). *Khasiat & Manfaat Lidah Buaya Si Tanaman Ajaib*. Purworejo: Agro Media Pustaka.
- Hosseinimehr, S.J., Khorasani, G., Azadbakht, M., Zamani, P., Ghasemi, M., & Ahmadi, A. (2010). Effect of aloe cream versus silver sulfadiazine for healing burn wounds in rats. *Acta Dermatovenerologica Croatica*, 18(1), 0-0.
- Hutter, J.A., Salman, M., Stavinota, W.B., Satsangi, N., Williams, R.F., Streeper, R.T., & Weintraub, S.T. (2006). Antiinflammatory C-glucosyl Chromone from Aloe Barbadensis. *J-Nat-Pro*, 59(5), 541-543.
- Irshad, S., Butt, M., & Younus, H. (2011). In-vitro antibacterial activity of Aloe barbadensis Miller (Aloe vera). *International research journal of pharmaceuticals*, 1(2), 59-64.
- Ismail. (2007). *Luka dan Perawatannya*. Retrieved from <http://images.mailmkes.multiply.com/attachment/0/RDd@AoKCEMAADk5LMI1/Merawat%20luka.pdf?nmid=88915450> on November 20th, 2011.
- Junquiera, L.C., Carneiro J. (2007). *Basic Histology: Text and Atlas*. (Edisi ke-11). New York: McGraw-Hill.
- Khorasani, G., Hosseinimehr, S.J., Azadbakht, M., Zamani, A., & Mahdavi, M.R. (2009). Aloe versus silver sulfadiazine creams for second-degree burns: a randomized controlled study. *Surgery today*, 39(7), 587-91.
- Kumar, V., Abbas, A.K., Fausto, N., Aster, J.C., Robbins, and Cottrans. (2010). *Pathologic Basis of Disease*. (8th ed) . Philadelphia: Saunders Elsevier.
- Kathuria, N., Gupta, N., Manisha, R., Prasad, R., Nikita. (2011). Biologic

- Effects of Aloe Vera Gel. *The Internet Journal of Microbiology*, 9, 2.
- Kurniawan, H. (2007). Efek Daun Kelor (*Moringa Oleifera*) dalam Mempercepat Penurunan Tanda Inflamasi Eritema pada Luka Bersih pada Marmut (*Cavia Porcellus*). *The Indonesian Journal of Health Science*, 1(1), 23-31.
- Kozier, B. et al. (2008). *Fundamental of Nursing: Concepts, Process and Practice*. New Jersey. Pearson Education.
- Marriner-Tomey & Alligood. (2010). *Nursing Theorists and Their Works*. (7th ed.) St.Louis: Mosby Elsevier, Inc.
- Mansjoer Ali, Kuspuji T., Rakhmi S., Wahyu Ika, Wiwik S.W. (2014). *Kapita Selektta Kedokteran*. (edisi 4). Jakarta: Media Aesculapius.
- Mendonça, F.A.S., Passarini Junior, J.R., Esquisatto, M.A.M., Mendonça, J.S., Franchini, C.C., & Santos, G.M.Td. (2009). Effects of the application of Aloe vera (L.) and microcurrent on the healing of wounds surgically induced in Wistar rats. *Acta Cirúrgica Brasileira*, 24(2), 150-55.
- Nursalam. (2011). *Manajemen Keperawatan: Aplikasi dalam Praktik Keperawatan Profesional Edisi 3*. Jakarta: Salemba Medika.
- Nugroho, W. (2008). *Keperawatan Gerontik & Geriatrik*. (ed 3). Jakarta, EGC.
- Oryan, A., Naeini, A.T., Nikahval B., Gorjian, E. (2010). Effect of aqueous extract of Aloe vera on experimental cutaneous wound healing in rat. *Veterinarski arhiv*, 80(4), 509-22.
- Oyelami, O.A., Oneyami, A., Oyedeji, O.A., & Adeyemi, L.A. (2009) Preliminary Study of Effectiveness of Aloe Vera in Scabies Treatment. *Journal Phytotherapy Research*, 23(10), 144882-4.
- Pellizzoni, M., Ruzickova, G., Kalhotka, L., Lucini, L. (2012). Antimicrobial activity of different Aloe barbadensis Mill. and Aloe arborescens Mill. leaf fractions. *Journal of Medicinal Plants Research*, 6(10), 1975-81.
- Ramamoorthy, L., Kemp, M.C., Tizard, I.R., Ramamoorthy, L., Kemp, M.C., & Tizard, I.R. (2011). *Acemannan, beta-(1,4)-asetat Mannan, menginduksi produksi oksida nitrat dalam baris sel makrofag RAW 264,7*. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8863833> on April, 25th 2012.
- Setiabudi, A.W. (2008). *Lidah Buaya*. Retrieved from <http://www.artikel-lidah-buaya/pages.htm> on September 9th, 2011.
- Sjamsuhidajat, R., & Jong, W.D. (2011). *Buku Ajar Ilmu Bedah Edisi ke 3*. Jakarta: EGC.
- Smeltzer, S., & Bare, C. (2010). *Brunner & Suddarth's Textbook of Medical Surgical Nursing 12th Edition*. Philadelphia : Lippincott Williams & Wilkins.
- Sukmono, J. (2006). *Pedoman Terapi Herbal*. Jember: Sanggar Meditasi dan Tanaman Obat "Adem Ati".
- Schwartz, S.I., Brunicaardi, F.C., Anderson, D.K., Billiar, T.R., Dunn, D.L., Hunter, J.G., Mathews, J.B., & Pollock, R.E. (2010). *Schwartz's Principles of Surgery Ninth Edition*. New York: McGraw Hill Professional.
- Tamher, S., & Noorkasiani. (2009). *Kesehatan Usia Lanjut dengan Pendekatan Asuhan Keperawatan*. Jakarta.
- Widagdo, T.D. (2010). *Perbandingan Pemakaian Aloe Vera 25%, 40% dan Silver Sulfadiazine 1% Topikal pada Penyembuhan Luka Bakar Derajat II*. Retrieved from <http://eprints.undip.ac.id/21436/>
- Wilksman, L.B., Solomonik, I., Spira, R., Tennenbaum, T. (2007). Novel Insights into Wound Healing Sequence of Events. *Toxicologic Pathol*, 35, 767-79.
- Zakariya, M. (2008). *Efektifitas*

Penggunaan Madu dibandingkan Povidone Iodine 10% terhadap Penyembuhan Luka Insisi pada Marmut (Cavia Cobaya). Unpublished thesis by Faculty of Nursing, Airlangga University, Surabaya.

Cite This Article As: Toha, M., Sujarwadi, M., & Zuhroidah, I. The Effect of Aloe Vera Extract and Discharge Planning to Accelerate Wound Healing of Older Adult Hernia Surgery Patients. *Nurse and Health: Jurnal Keperawatan* 2019; 8(2): 124-133.

