



INDONESIAN JOURNAL OF
**NURSING
PRACTICES**

Improvement of Prolactin Hormone Levels on Postpartum Mothers Taken by The Oketani Massage and Pressure in GB-21 Point

Machmudah, Nikmatul Khayati, Sri Widodo, Elsi Dwi Hapsari, Fitri Haryanti

The Effect of Touch and Sound Therapy to The Level of Pain and Physiological Parameters in Infants

Rafika Dora Wijaya, Tita Husnitawati Madjid, Enny Harliany Alwi

Nurses' Attitude Toward End of Life Care in Emergency Departement and Intensive Care Unit In Rural Hospital

Muhammad Zulfatul A'la, Zumrotul Farikhah, Mulia Hakam

The Correlation between The Knowledge Level of Diabetes Management toward The Preprandial Glucose Levels

Fahni Haris, Linda Yuli Kristianti

Health Lecturers and Students Views' about Healthcare Workers' as Healthy Lifestye Role Models: a Qualitative Study

Agustina Arundina Triharja Tejoyuwono

The Effects of Hypnosis Relaxation Towards Fetus Heart Frequency and Vital Signs among Pregnant Women with Preeclampsia

Siti Khuzaiyah, Anies, Sri Wahyuni

Relationship Between Drug Consumption, Supervisors' Knowledge and Support, and Patients' Obedience to Take Tuberculosis Drugs

Ni Luh Widani, Sondang Sianturi

Vol 4, No 1

June 2020

VOLUME 4 NO 1 JUNE 2020

INDONESIAN JOURNAL OF
**NURSING
PRACTICES** 

Editorial board of IJNP (*Indonesian Journal of Nursing Practices*)

Editor in Chief:

Erna Rochmawati (Universitas Muhammadiyah Yogyakarta)

Managing Editor:

Yanuar Fahrizal (Universitas Muhammadiyah Yogyakarta)

Editor:

Herni Susanti (Universitas Indonesia)

Asniar (Universitas Syiah Kuala Aceh)

Shanti Wardaningsih (Universitas Muhammadiyah Yogyakarta)

Dewi Puspita (Universitas Muhammadiyah Yogyakarta)

Nur Chayati (Universitas Muhammadiyah Yogyakarta)

Danita Sugiyo (Universitas Muhammadiyah Yogyakarta)

Yusi Riwayatul Afsah (Universitas Muhammadiyah Yogyakarta)

Novita Kurnia Sari (Universitas Muhammadiyah Yogyakarta)

Nurul Hidayah (Universitas Muhammadiyah Yogyakarta)

Office:

School of Nursing, Faculty of Medicine and Health Science

Universitas Muhammadiyah Yogyakarta

Siti Walidah Building F3, 4th floor

Brawijaya Street, Tamantirto, Kasihan, Bantul, Yogyakarta

Telp +62 274 387656 ext 215

Email: IJNP@umy.ac.id

Table of Contents

- 1 **Improvement of Prolactin Hormone Levels on Postpartum Mothers Taken by The Oketani Massage and Pressure in GB-21 Point**
Machmudah, Nikmatul Khayati, Sri Widodo, Elsi Dwi Hapsari, Fitri Haryanti
- 7 **The Effect of Touch and Sound Therapy to The Level of Pain and Physiological Parameters in Infants**
Rafika Dora Wijaya, Tita Husnitawati Madjid, Enny Harliany Alwi
- 14 **Nurses' Attitude Toward End of Life Care in Emergency Departement and Intensive Care Unit In Rural Hospital**
Muhammad Zulfatul A'la, Zumrotul Farikhah, Mulia Hakam
- 21 **The Correlation between The Knowledge Level of Diabetes Management toward The Preprandial Glucose Levels**
Fahni Haris, Linda Yuli Kristianti
- 28 **Health Lecturers and Students Views' about Healthcare Workers' as Healthy Lifestye Role Models: a Qualitative Study**
Agustina Arundina Triharja Tejoyuwono
- 37 **The Effects of Hypnosis Relaxation Towards Fetus Heart Frequency and Vital Signs among Pregnant Women with Pre-eclampsia**
Siti Khuzaiyah, Anies, Sri Wahyuni
- 43 **Relationship Between Drug Consumption, Supervisors' Knowledge and Support, and Patients' Obedience to Take Tuberculosis Drugs**
Ni Luh Widani, Sondang Sianturi

**Muhammad Zulfatul A'la¹,
Zumrotul Farikhah¹, Mulia Hakam¹**
¹Universitas Jember

Corresponding Author: Muhammad Zulfatul A'la
Email: m.zulfatul@unej.ac.id

Nurses' Attitude Toward End of Life Care in Emergency Departement and Intensive Care Unit In Rural Hospital

Article Info

Online : <http://journal.ums.ac.id/index.php/ijnp>
ISSN : 2548 4249 (Print)
: 2548 592X (Online)
DOI : 10.18196/ijnp.41103

Abstract

Background: Death is a psychological and physical event that affects patients caring for, especially in Emergency Departement (ED) and the Intensive Care Unit (ICU). Patients and families need the end of life care from a health professional, including nurses. Nurses need to have knowledge, skill, attitude, and interpersonal competencies to provide end of life care.

Objective: This study aimed to explored differences of attitudes towards dying care between ED and ICU ward nurses in a rural hospital.

Method: Variable in this study was the nurse's attitude. This study used a quantitative comparative cross-sectional research design. The samples were 24 nurses from the emergency ward and 16 from the intensive care unit who were recruited by total sampling. Data were collected to use Frommelt Attitudes Towards the Care of the Dying Care Form B Indonesian version (FATCOD-BI).

Result: The results revealed that nurses' attitudes toward caring for dying patients in the ED were 101.42 ± 6.646 (30-150) on average, and in the intensive care unit was 106.44 ± 7.633 (30-150) on average. Nurses in the ICU had a more positive attitude than the ED ($p = 0.034$). There are differences between the proportion of respondents based on gender, level of education, length of time working in the emergency ward, and ICU. The result showed gender, level of education, and range of time working could influence nurses' attitudes towards caring for dying patients in the emergency ward and ICU.

Conclusion: There is a difference between nurses' attitudes towards EOLC in the ED and ICU room at rural hospitals. Moreover, the ICU nurses have a more positive attitude than ED nurses.

Keywords: Caring; Dying patients; Emergency department; ICU; Nurse's attitudes

INTRODUCTION

Patients generally experience deaths in the terminal phase, especially in the Emergency Departement (ED), Intensive Care Unit (ICU), and inpatient rooms (Beckstrand et al., 2017). Patients need end of life care such as reducing symptoms and family support when entering the process of grieving (Gama, Barbosa, Vieira, 2012). Death is a psychological and physical event that affects not only the patient who experiences it but also the people around them,

including health professionals (Gama, Barbosa, Vieira, 2012). Fear, anxiety, distress, and sadness can occur in families and health professionals when facing patients with death (Buglass, 2010). End of life care (EOLC) is an essential concept in of palliative care (National Consensus Project for Quality Palliative care, 2009). A nurse is involved in the process of dying directly (Gastmans, 2012). The care delivered is the management of symptoms related to death and the psychological, social, spiritual

support (Higgs, 2010). Nurses are essential health care providers that provided caring processes in ICU, emergency room, and home. Therefore, Nurses need knowledge and skills to offer effective EOLC. Also, attitude toward EOLC and interpersonal competencies need to be improved by nurses (Ali & Ayoub, 2010).

The ED and ICU have several similarities in characteristics of care. Firstly, ED and ICU received the patient who is in advanced illness. Moreover, Both of them require specialized health care providers (doctors, nurses, assistants, etc.). Therefore, nurses in the ED and ICU need special skills or expertise. Secondly, ED and ICU have aim to life-saving in the terminal patient, so the resuscitation procedure is mostly applied. Therefore, the death process occurs in those rooms regularly. Thirdly, ED and ICU have an inappropriate environment for patients and families because it is no particular room for the family, so it affects the quality of EOLC (Beckstrand et al., 2017).

EOLC in the ED is a challenge for nurses because the ED is the first place to receive patients in the hospital setting (Decker et al., 2015). Decker et al. (2015) stated that it is several barriers to conduct EOLC in the emergency room, such as lack of time, low priority, lack of privacy, and unsuitable environment. Otherwise, in ICU, it has different characteristics and obstacles for applying EOLC. ICU is a place for patients with complicated medical conditions (Baliza et al., 2013). The study in ICU revealed that ICU has several obstacles to the provision of EOLC, including the lack of nursing staff, lack of communication between patients, doctors, and family members (Beckstrand et al., 2017).

A nurse can implement the EOLC, including giving attention and direction related to appropriate treatment, providing social support for patients and families, delivering interventions to reduce pain response, discussing the death process with patients, and listening to the client's wishes for end of life (WHO, 2014). Moreover, the principles of dying care are delivering a patient to a peaceful death (Gloss & Hospitals, 2017).

Nursing Interventions in EOLC supposed to be based on a positive attitude of caring for dying patients. A positive attitude of nurses in the care process can influence the effectiveness of EOLC (Gallagher et al.,

2015). The results of the study stated that negative nurses' attitudes, such as apathy, fear, and anxiety, could reduce the level of quality of care in dying patients (Grubb & Arthur, 2016). The attitude of nurses towards EOLC care is essential to explore as a reference and evaluation of the EOLC process given to clients in the emergency room and ICU.

Moreover, the concept of EOLC in rural hospitals is an essential issue because rural hospitals have unique services and have limited access and infrastructure (Bakitas et al., 2015). However, according to Rainsford et al. (2017) illustrates that EOLC in the rural area is still lack of article that discussed this issue. Based on the background that has been stated, it is interesting to explore nurses 'attitudes towards the end of life and to compare nurses' attitudes in the ED and ICU in rural hospitals.

METHOD

A quantitative study is applied in this study and used a comparative descriptive survey method. This study utilized a cross-sectional approach (Polit & Beck, 2009). In this study, researchers used a comparative analysis (comparison) to distinguish or compare nurses' attitudes towards EOLC in ED and ICU.

The sampling technique used in this study is a non-probability sampling. The sampling method in this study is total sampling with a total number of 40 respondents consisting of 24 nurses in ED and 16 nurses in the ICU at two rural hospitals in Jember. The research data was taken from December 2018 to January 2019. Nurse attitude toward EOLC utilized the FATCOD-BI questionnaire (The Frommelt Attitudes Toward Care of The Dying Care Form B (FATCOD-B) Indonesian Version). The FATCOD-B questionnaire consisted of 30 statement items. The FATCOD-B instrument consists of favorable and unfavorable statements. The statement items included in the favorite statement are items number 1, 2, 4, 16, 18, 20, 21, 22, 23, 24, 25, 27, and 30 (A'la, 2016). The favorable statement has a score of 1 (strongly disagree) and a score of 5 (strongly agree) and the other items included in the unfavorable statement. The Likert scale started from 1 (strongly agree) until 5 (strongly disagree) is used. The total score is between 30 and 150. The higher score described that is more positive the attitude toward the end of life care (Tait et al., 2015). The Frommelt Attitudes Towards Care of The Dying Care Form B

(FATCOD-B) Indonesian version has been tested for reliability by A'la (2016) using Cronbach's alpha. The results of the reliability test were 0.86. The validity range of each statement item is 0.651-0.713. FATCOD-BI used two domains in its discussion, namely Domain I (Positive attitude towards caring for dying patients domain) and Domain II (Perception of patient care based on family and patient domains). FATCOD-BI used two domains in its discussion, namely Domain I (Positive attitude towards caring for dying patients domain) and Domain II (Perception of patient care based on family and patient domains). Domain I of FATCOD-BI consists of item number 1, 2, 3, 5, 6, 7, 8, 9, 11, 13, 14, 15, 17, 26, 29, 30. Domain II of FATCOD-BI consists of item number 4, 10, 12, 16, 18, 19, 20, 21, 22, 23, 24, 25, 27, and 28 (Henoeh et al., 2013).

The Ethical Committee of Faculty of Dentistry, University of Jember, has approved this research with numbers 237 / UN25.8 / KEPK / DL / 2019. Before filling out the questionnaire, the respondent received informed consent. Moreover, patients have the autonomy to choose to participate in the research process.

Data analysis using SPSS 20. The T-test is used to analyze the comparison of nurses' end of life attitude in ED and ICU because the data are normally distributed. Furthermore, to see the difference in the proportion of respondents' characteristics in ED and ICU nurses, researchers used the Chi-square test.

RESULTS

Table 1 explained that the average respondent's age working in ED was 32.25 years. Otherwise, in the ICU, the average was 34.88 years. The independent t-test results found that there was no difference in the average age (p-value > 0.05). Moreover, The proportion of male nurses in ED (n = 15, 62.5%) was more than women (n = 9, 37.5%). However, in ICU, the proportion of females is more than male. Chi-square test showed that there was a difference in the proportion of gender in ED and ICU (p-value <0.05). Also, in education level, it showed that it was a difference in the proportion in ED and ICU with a p-value of 0.046 (p-value <0.05). Nevertheless, it was no homogeneity in gender and level of education proportion (p-value >0.05).

Table 1. Description of characteristics of respondents in Nurses (n=40).

Characteristic of Respondent	ED (N=24) (n)	%	ICU (N=16) (n)	%	P-value
Ages (Mean±SD)	32,25 ±6,045		34,88 ±3,810		0,132 ^a
Gender					
Male	15	62,5	4	25,0	0,045 ^{b*}
Female	9	37,5	12	75,0	
Level of Education					
Nursing Diploma	13	54,2	10	62,5	0,046 ^{b*}
Nursing Bachelor	1	4,2	4	25,0	
Nurse Profession	10	41,7	2	12,5	
Work Experience					
<5 years	8	33,3	0	0	0,003 ^{b*}
5-10 years	13	54,2	7	43,8	
10-15 years	3	12,5	4	25,0	
>15 years	0	0	5	31,3	
Religion					
Muslim	24	100,0	16	100,0	-
Non-Muslim	0	0	0	0	
Marriage status					
No married	5	20,8	-	-	0,071 ^b
Married	19	79,2	16	100	
Experience in caring for dying patient					
Yes	21	87,5	16	100	0,262 ^b
No	3	12,5	-	-	
Training experience in caring for dying patient					
Yes	14	58,3	4	25,00	0,080 ^b
No	10	41,7	12	75,00	

^a Non-Paired t-Test ^b Chi Square *have a correlation

Nurses in ED who have less than five years experience amounted to 8 people, 5-10 years there were 13 people, and 10-15 years amounted to 3 people. While in the ICU room, there are no nurses who have work experience of fewer than five years. Chi-square test results between the length of work and workplace variables (emergency room and ICU) showed that there was a difference between the number of nurses working in the emergency room and ICU based on the length of work with p-value 0.003 (p-value <0.05). In the emergency room, some nurses are just starting to enter the workforce so that their experience is <5 years. While in the ICU room, there are no nurses who have work experience <5 years, and there are nurses who have been senior or have worked for >15 years.

All nurses are Muslim (100%), and there are five unmarried emergency nurses and 19 married nurses, while 16 nurses who worked in the ICU; all of them were married. Nurses who have experience dealing with dying patients are 37 people (92.5%),

and the number of nurses who have no experience dealing with dying patients is three people (7.5%). Three nurses who have never dealt with patients near death are working in the emergency room. All nurses who worked in the ICU (N = 16) had faced patients dying. The number of nurses who had attended the training of patient care before death 16 people consisting of 14 nurses in the emergency room and four nurses in the ICU room. Whereas, who had never received training in dying care patients consisted of 10 emergency nurses and 12 ICU nurses. Overall, more nurses have never attended training than those who have ever participated in the training.

Tabel 2. The difference in the nurse's attitude toward EOLC (N=40)

Variable	FATCOD-BI		Total	P-Value
	ED Mean (SD)	ICU Mean (SD)		
Domain I (Positive attitude towards caring for dying patients domain)	53,13 (6,924)	54,00 (4,050)	53,48 (5,896)	0,618
Domain II (Perception of patient care based on family and patient domains)	48,29 (5,162)	52,44 (5,151)	49,95 (5,491)	0,017*
FATCOD BI Total	101,42 (6,646)	106,44 (7,633)	103,43 (7,393)	0,034*

*have differences

Table 2 described the nurses' attitudes towards EOLC in the ED and ICU at rural hospitals. Overall the average attitude in the ICU (106.44) is higher than in the ED (101.42). Furthermore, according to the FATCOD-BI domain, the domain I described that attitude toward EOLC in ICU (54.00) was higher than in the ED (53.13). Domain II also showed that the attitude toward EOLC in ICU (52.44) was higher than in the emergency room (48.29). Moreover, Table 2 described that there is a difference between nurses' attitudes towards EOLC in the ED and ICU with a p-value=0.034 ($p < 0.05$). However, if it was tested in each domain, domain I showed no difference attitude with a p-value=0.618 ($p > 0.05$), and domain II showed a different attitude between nurses in ED and ICU with a p-value=0,017 ($p < 0.05$).

DISCUSSION

The results showed that there were nurse's attitude differences in ED and ICU towards EOLC. The attitude of nurses in the ICU was more positive

rather than in the ED. Based on each domain, the results showed a difference in domain II (Perception of patient care based on family and patients). Tripathy et al. (2017) stated that in the ICU, most family members had a contribution for EOLC, especially supportive care. This study also revealed that families often discuss with nurses and always appreciate the role of nurses in the ICU (Tripathy et al., 2017).

Moreover, Beckstrand et al. (2017) stated that there are many obstacles experienced by emergency nurses in applying EOLC. These obstacles will affect the attitude of the nurses in the ED towards EOLC. Some of these obstacles are family who contacts the nurse frequently to ask about the patient's condition. At the same time, nurses also have to take care of other patients, and another obstacle is the emergency room design that does not allow for the privacy of patients with dying or grieving family members. Therefore, it will affect the attitude of nurses and perceptions of dying patient care based on family and patients. The difference in nurses' attitudes toward EOLC in ED and ICU is also influenced by the lack of time for dying in the emergency room, low priority, lack of privacy, and an inappropriate ER environment (Decker et al., 2015). Nurses cannot apply EOLC in one patient only because many patients need help quickly and appropriately. Otherwise, in rural hospitals, the quantity of specialized nurses is a central problem. Nurses also seldom discussed with patients and family about the feelings because of the lack of privacy in ED (Decker et al., 2015). Barriers and obstacles in EOLC can affect the ability of nurses to communicate with patients and families regarding EOLC (Gurung & Timalsina, 2018). Moreover, The results of this study were supported by previous research, which states that the nurse's work environment can influence nurses' attitudes towards EOLC (Kassa & Murugan, 2014).

Homogeneity test in the characteristics of respondents between nurses in the ED and ICU showed that religion, age, marital status, experience caring for dying patients, and the experience of receiving dying care training was homogenous. However, sex, work experience, and level of education were no homogeneous, or there are differences in proportions between nurses in the ED and ICU. Based on these results, gender, education level, and work experience factors might influence

nurses' attitudes towards EOLC. Furthermore, based on previous research, gender is one of the factors that influence a nurse's attitude towards EOLC. According to Yaqoob et al. (2018), there was a relationship between gender and the attitude of nursing students towards EOLC. Other studies also revealed that women are considered to have a more positive attitude towards caring for dying patients than men (Lancaster et al., 2017). In Addition, feminism could affect the attitude of caring for dying patients (A'la et al., 2018; Dunn et al., 2005).

Work experience may affect attitude toward EOLC and affect the differences of nurses in the ED and ICU. The result showed that Nurses in the ICU have more extensive working experience compared to emergency room nurses so that they can influence attitudes towards EOLC. Also, these results were supported by previous studies. In Turkey, the result showed that there were differences in attitudes between groups of nurses who worked from 2 years to 5 years and groups from 5 years to 10 years (Ozcelik et al., 2018).

Education level has also could influence nurses' attitudes towards EOLC. According to Dunn et al. (2005), who analyzed nurses' attitudes in Sweden, found that education level is the best predictor of positive attitudes in caring for dying patients (Lancaster et al., 2017). Another study in Iran also revealed a significant relationship between education level and attitudes towards death (Ali & Ayoub, 2010).

Balung and Kalisat hospitals are hospitals located in rural areas or peripheral hospitals in Jember Regency. In rural areas, there are many emergency cases, so that rural hospitals are the first referral for local communities, even though human resources and facilities are still incomplete and under the standard. This situation may increase the risk of death of patients both in the ED and ICU due to delays in the referral of patients who are in an emergency and require prompt, appropriate, and adequate health facilities (Green et al., 2017). Therefore, the positive attitude of nurses towards EOLC in the ED and ICU in rural hospitals is needed to improve the quality of service to patients. The development of EOLC in rural hospitals also needs to be developed both in the ED and ICU. A positive

attitude towards EOLC in the ED nurses and ICU nurses could be an initial step in the development of EOLC services in rural hospitals.

According to researchers, the differences in nurses' attitudes towards EOLC in the ED and ICU may be caused by the environmental factor in both areas. in the ED, the patient lacked privacy so that affecting EOLC services. Otherwise, in the ICU, the time for caring is 24 hours, and in general, patient in ICU is treated for a long time. For further research, it needs to examine the relationship between sex, education level, and also the experience of working nurses with nurses' attitudes towards EOLC because it can be factors that influenced attitude toward EOLC.

CONCLUSION

There is a difference between nurses' attitudes towards EOLC in the ED and ICU room at rural hospitals. Moreover, the ICU nurses have a more positive attitude than ED nurses. The proportion of characteristics respondents based on sex, education level, and length of work was no homogeneous. Therefore, it could be a factor in differences of attitude toward EOLC. The development of EOLC in rural hospitals can begin with improving attitudes of caring for dying patients through training or refreshing knowledge to nurses. However, this study still has limitations because it was conducted on heterogeneous samples. For further research, it needs to examine the relationship between sex, education level, work experience with nurses' attitudes towards EOLC to heterogeneous samples.

REFERENCES

- A'la, M. Z. (2016). The Frommelt Attitudes toward Care Of The Dying Care Form B (FATCOD-B) Indonesia version: measurement validity using factor analysis in nursing students. *Nurseline Journal*, 1(1), 73-82. <https://jurnal.unej.ac.id/index.php/NLJ/article/view/5995>
- A'la, M. Z., Setioputro, B., & Kurniawan, D. E. (2018). Nursing Students' Attitudes towards Caring for Dying Patients. *Nurse Media Journal of Nursing*, 8(1), 25-34. <https://doi.org/10.14710/nmjn.v8i1.17270>

- Ali, W. G. M., & Ayoub, N. S. (2010). Nurses' attitudes toward caring for dying patient in Mansoura university hospitals. *Journal of Medicine and Biomedical Sciences*, 1(1), 16-23.
- Bakitas, M. A., Elk, R., Astin, M., Ceronsky, L., Clifford, K. N., Dionne-Odom, J. N., ... & Ritchie, C. (2015). Systematic review of palliative care in the rural setting. *Cancer Control*, 22(4), 450-464. <https://doi.org/10.1177/107327481502200411>
- Baliza, M. F., Bousso, R. S., Poles, K., Santos, M. R. D., Silva, L., & Paganini, M. C. (2015). Factors influencing intensive care units nurses in end-of-life decisions. *Revista da Escola de Enfermagem da USP*, 49(4), 0572-0579. <https://doi.org/10.1590/S0080-623420150000400006>
- Beckstrand, R. L., Smith, K. E., Luthy, K. E. B., & Macintosh, J. L. (2017). Rural emergency nurses' suggestions for improving end-of-life care. *Journal of Emergency Nursing*, 43(3), 214-220. <https://doi.org/10.1016/j.jen.2017.03.012>
- Buglass, E. (2010). Grief and bereavement theories. *Nursing Standard (through 2013)*, 24(41), 44-47. <https://doi.org/10.7748/ns2010.06.24.41.44.c7834>
- Decker, K., Lee, S., & Morphet, J. (2015). The experiences of emergency nurses in providing end-of-life care to patients in the emergency department. *Australasian Emergency Nursing Journal*, 18(2), 68-74. <https://doi.org/10.1016/j.aenj.2014.11.001>
- Dunn, K. S., Otten, C., & Stephens, E. (2005, January). Nursing experience and the care of dying patients. In *Oncology nursing forum*, 32(1), 97-104. <https://doi.org/10.1188/05.ONF.97-104>
- Decision, I. N. E. O. L., Gallagher, A., Bousso, R. S., McCarthy, J., Kohlen, H., Andrews, T., ... & Arber, A. (2015). Negotiated reorienting: a grounded theory of nurses' end-of-life decision-making in the intensive care unit. *International Journal of Nursing Studies*, 52(4), 794-803. <https://doi.org/10.1016/j.ijnurstu.2014.12.003>
- Gama, G., Vieira, M., & Barbosa, F. (2012). Factors influencing nurses' attitudes toward death. *International Journal of Palliative Nursing*, 18(6), 267-273. <https://doi.org/10.12968/ijpn.2012.18.6.267>
- Gastmans, C. (2012). Nursing ethics perspectives on end-of-life care. *Nursing ethics*, 19(5), 603-604. <https://doi.org/10.1177/0969733012453234>
- Gloss, K. (2017). End of life care in emergency departments: a review of the literature. *Emergency Nurse*, 25(2), 29-38. <https://doi.org/10.778/en.2017.e1707>
- Green, M. A., Dorling, D., Minton, J., & Pickett, K. E. (2017). Could the rise in mortality rates since 2015 be explained by changes in the number of delayed discharges of NHS patients?. *J Epidemiol Community Health*, 71(11), 1068-1071. <https://doi.org/10.1136/jech-2017-209403>
- Grubb, C., & Arthur, A. (2016). Student nurses' experience of and attitudes towards care of the dying: A cross-sectional study. *Palliative medicine*, 30(1), 83-88. <https://doi.org/10.1177/0269216315616762>
- Gurung, M., & Timalisina, R. (2018). Attitudes of Nepalese hospital nurses towards care of dying people. *J Aging Stud Ther*, 1(1), 1-4. <https://doi.org/10.16966/jast.101>
- Henoch, I., Browall, M., Melin-Johansson, C., Danielson, E., Udo, C., Sundler, A. J., ... & Strang, S. (2014). The Swedish version of the Frommelt Attitude Toward Care of the Dying Scale: Aspects of validity and factors influencing nurses' and nursing students' attitudes. *Cancer nursing*, 37(1), E1-E11. <https://doi.org/10.1097/NCC.0b013e318279106b>
- Higgs, C. (2010). *The Palliative care handbook: Advice on clinical management* (7th ed.). Hierographics Ltd.
- Kassa, H., & Murugan, R. (2014). Assessment of knowledge, attitude and practice and associated factors towards palliative care among nurses working in selected hospitals, Addis Ababa, Ethiopia. *BMC Palliative Care*, 13(6), 1-11.

- <http://bmcpalliatcare.biomedcentral.com/articles/10.1186/1472-684X-13-6>
- Lancaster, R. J., Kautzmann, C., Micheal, J. C. J., Chandrasekaran, L., Jambunathan, J., & Chapin, T. M. (2017). Attitudes of nurses towards care of the dying patient in India. *International Journal of Palliative Nursing*, 23(11), 558-566. <https://doi.org/10.12968/ijpn.2017.23.11.558>
- National Consensus Project for Quality Palliative care. (2009). *Clinical practice guidelines for quality palliative care* (2nd ed.). National Consensus Project for Quality Palliative Care.
- Özcelik, H., Aksoy, F., Sonmez, E., & Fadiloglu, Ç. (2018). Attitudes to death of nurses in Turkey and factors affecting them. *Hos Pal Med Int Jnl*, 2(6), 303-309. <https://doi.org/10.15406/hpmij.2018.02.00112>
- Polit, D. F., & Beck, C. T. (2009). *Nursing research: Principles and methods* (7th Ed). Lippincott Williams & Wilkins.
- Rainsford, S., MacLeod, R. D., Glasgow, N. J., Phillips, C. B., Wiles, R. B., & Wilson, D. M. (2017). Rural end-of-life care from the experiences and perspectives of patients and family caregivers: A systematic literature review. *Palliative Medicine*, 31(10), 895-912. <https://doi.org/10.1177/0269216316685234>
- Tait, V., Higgs, M., Magann, L., Dixon, J., Davis, J. M., & Fernandez, R. (2015). Attitudes of nonpalliative care nurses towards palliative care. *International Journal of Palliative Care*, 2015, 469174. <https://doi.org/10.1155/2015/469174>
- Tripathy, S., Routray, P. K., & Mishra, J. C. (2017). Intensive care nurses' attitude on palliative and end of life care. *Indian J Crit Care Med*. 21(10): 655-659. https://doi.org/10.4103/ijccm.IJCCM_240_16
- World Health Organization. & Alliance, W. P. C., (2014). *Global atlas of palliative care at the end of life*. London: Worldwide Palliative Care Alliance.
- Yaqoob, M., Nasaif, H., & Kadhom, H. (2018). Undergraduate final year nursing students' attitudes toward caring for dying patients: Bahrain's experience. *Clinical Nursing Studies*, 6(4), 53-58. <https://doi.org/10.5430/cns.v6n4p53>