


dentika

DENTAL JOURNAL

VOLUME 16 NUMBER 2 DECEMBER 2011



Open Menu

dentika

DENTAL JOURNAL

[Skip to main content](#)

[Skip to main navigation menu](#)

[Skip to site footer](#)

[Home](#) / [Archives](#) / Vol. 16 No. 2 (2011): Dentika Dental Journal

Vol. 16 No. 2 (2011): Dentika Dental Journal




ISSN : 1693-671X

dentika

DENTAL JOURNAL

VOLUME 16 NUMBER 2 DECEMBER 2011



dentika DENTAL JOURNAL

Vol. 16

No. 2

P. 103 - 202

Medan
December 2011

ISSN
1693 - 671X

Terbitan ini merupakan pendigitalan dokumen jurnal Dentika Dental Journal Vol 16 No 2 (2011) yang pernah terbit secara cetak oleh Dentika Dental Journal pada Tahun 2011.

Published: 2011-12-07

Articles

PENGARUH DURASI PERENDAMAN RESIN AKRILIK DALAM MINUMAN KOPI ACEH ULEE KARENG TERHADAP KEKERASAN PERMUKAAN

EFFECT OF ACRYLIC RESIN'S IMMERSION DURATION COFFEE TO SURFACE HARDNESS

Viona Diansari, Iin Sundari, Rini Defika Putri

103-107

[PDF](#)

PENGARUH PASTA GIGI YANG MENGANDUNG ENZIM TERHADAP PENURUNAN KADAR VOLATILE SULFUR COMPOUND

EFFECT OF ENZYME TOOTH PASTE IN DECREASING VOLATILE SULFUR COMPOUND LEVEL

Zulfan Muhammad Alibasyah, Irene Sukardi, Dewi Nurul Mustaqimah

108-110

[PDF](#)

EFEK EKSTRAK ALOE VERA TERHADAP CANDIDA ALBICANS YANG DIISOLASI DARI RONGGA MULUT PENDERITA HIV/AIDS DENGAN TEKNIK TUBE DILUTION

EFFECT OF ALOE VERA LEAF EXTRACT TO CANDIDA ALBICANS ISOLATED FROM ORAL CAVITY OF HIV/AIDS PATIENTS BY TUBE DILUTION TECHNIQUE

Elizabeth Fitriana Sari, Gus Permana Subitha

111-115

[PDF](#)

SIFAT OSTEOINDUKTIF SILIKA AMORPHOUS SEKAM PADI OSTEOINDUCTIVITY OF AMORPHOUS SILICA FROM RICE HUSK

Didin Erma Indahyani, Zahreni Hamzah, Izzata Barid

116-120

[PDF](#)

HUBUNGAN POLIMORFISME GEN VITAMIN D RESEPTOR DENGAN PERIODONTITIS KRONIS

POLYMORPHISM OF VITAMIN D RECEPTOR GENE IS ASSOCIATED WITH CHRONIC PERIODONTITIS

Nurlindah Hamrun

121-125

[PDF](#)

OUTER MEMBRANE PROTEIN OF PORPHYROMONAS GINGIVALIS AS BASIC OF DIAGNOSTIC TOOL IN AGGRESSIVE PERIODONTITIS

PROTEIN MEMBRAN BAGIAN LUAR PORPHYROMONAS GINGIVALIS SEBAGAI ALAT DIAGNOSTIK PERIODONTITIS AGRESIF

Desi Sandra Sari, Candra Bumi, Yuliana Mahdiyah DA

126-129

[PDF](#)

BAHAN BONDING DAN KAMFOROQUINON MAMPU MENINGKATKAN KEKERASAN RESIN KOMPOSIT SINAR-TAMPAK YANG MENGALAMI KEMAMPUAN MENGALIR

ABILITY OF BONDING AGENT AND CHAMPHOROGUINONE TO INCREASE VISIBLE LIGHT COMPOSITE RESIN HARDNESS IN FLOW READILY

Ekiyantini Widowati, Dwi Warna Aju Fatmawati

130-133

[PDF](#)

ANTIBACTERIAL EFFECT OF GLASS IONOMER AND ZINC OXIDE CHKM CEMENT AS ROOT CANAL FILLER TOWARD STREPTOCOCCUS VIRIDANS

EFEK ANTIBAKTERI SEMEN IONOMER KACA DAN ZINK OKSID CHKM SEBAGAI BAHAN PENGISI SALURAN AKAR TERHADAP STREPTOCOCCUS VIRIDANS

Dwi Warna Aju Fatmawati, Ekiyantini Widowati

134-138

[PDF](#)

ACTIVITY TEST OF IMMUNOMODULATORY COMPONENT FROM NEEM LEAVES TO CANDIDA ALBICANS

UJI AKTIVITAS KOMPONEN IMUNOMODULATOR DARI NEEM TERHADAP CANDIDA ALBICANS

I Dewa Ayu Ratna Dewanti, I Dewa Ayu Susilawat, Purwanto, Depi Praharani

139-143

[PDF](#)

EFEK EKSTRAK BUAH JAMBLANG TERHADAP PERTUMBUHAN STREPTOCOCCUS MUTANS SEBAGAI PENYABAB UTAMA KARIES

EFFECT OF JAMBLANG'S FRUIT EXTRACT TI STREPTOCOCCUS MUTANS GROWTH AS MAIN AGENT OF CARIES

Santi Chismirina, Poppy Andriyani, Nopi Yanti Fitri

144-148



EFFECT OF CASEIN PHOSPHO PEPTIDE-AMORPHOUS CALCIUM FLUOR PHOSPHATE FOR INHIBITING STREPTOCOCCUS MUTANS GROWTH IN YOUNG ADULT PATIENTS

EFEK CASEIN PHOSPO PEPTIDE-AMORPHOUS CALCIUM FLUOR PHOSPHATE UNTUK MENGHAMBAT PERTUMBUHAN STREPTOKOKUS MUTANS PADA PASIEN DEWASA MUDA

Ratna Meidyawati, Anggraeni

149



KEBUTUHAN PERAWATAN PENYAKIT PERIODONTAL DAN PERILAKU PEMELIHARAAN GIGI PADA MASYARAKAT DI KECAMATAN PANGURURAN SAMOSIR

PERIODONTAL TREATMENT NEED AND DENTAL HEALTH CARE BEHAVIOR IN SOCIETY OF PANGURURAN SAMOSIR

Simson Damanik, Josevina Silalahi

154



PENGARUH EKSTRAK KELOPAK BUNGA ROSELLA TERHADAP PERTUMBUHAN CANDIDA ALBICANS DI RONGGA MULUT

EFFECT OF HIBISCUS SABDARIFFA TO THE GROWTH OF CANDIDA ALBICANS IN THE MOUTH

Poppy Andriany, Santi Chismirina, Faolyna

159



PERAN ORANGTUA TERHADAP PEMELIHARAAN KESEHATAN GIGI DAN MULUT ANAK DAN STATUS KESEHATAN GIGI DAN MULUT ANAK KELAS II SD MEDAN

PARENT'S ROLE OF DENTAL ORAL HEALTH CARE AND DENTAL HEALTH STATUS OF THE SECOND GRADE STUDENT IN MEDAN

Lina Natamiharja, Margaret

163



EFFECT AMOUNT OF CIGARETTE SMOKING ON GINGIVAL EPITHELIUM THICKNESS
EFEK JUMLAH ROKOK SIGARET TERHADAP KETEBALAN EPITHEL GINGIVAL

Asam Khalifa Mohammed, Suryono, Siti Sunarintyas

168



INFLAMMATION RESPONSE OF MECHANICALLY EXPOSED PULP AFTER DIRECT PULP CAPPING WITH CALCIUM HYDROXIDE CEMENT AND PLATELET RICH PLASMA
RESPONS PERADANGAN PADA PULPA TERBUKA SECARA MEKANIK SETELAH PULP CAPPING DIRECT DENGAN SEMEN KALSIUM HIDROKSID DAN PLATELET RICH PLASMA

Sartika Puspita, Totok Utoro, Tetiana Haniastuti

171-174



PENATALAKSANAAN MINOR ERYTHEMA MULTIFORME
MANAGEMENT OF MINOR ERYTHEMA MULTIFORME

Setya Wardani, Mintarsih Djamhari K, Dwi Setianingtyas

175-179



BLEACHING KASUS DISKOLORASI TETRASIKLIN DERAJAT 3
BLEACHING OF THE THIRD DEGREE TETRACYCLINE DISCOLORATION

Any Setyawati

180-183



TREATMENTS OF CLASS III SEVERE CROWDED TEETH WITH DAMON BRACKET SYSTEM: EXTRACTION VS NON EXTRACTION
PERAWATAN GIGI GELIGI BERLAPIS KELAS III DENGAN SISTEM BRAKET DAMON: DENGAN DAN TANPA PENCABUTAN GIGI

Erliera

184-188

PENATALAKSANAAN MOUTH PREPARATION PADA ANAK DENGAN KELAINAN JANTUNG BAWAAN TIPE ATRIAL SEPTAL DEFECT DI BAWAH ANESTESI UMUM
MOUTH PREPARATION MANAGEMENT IN CHILDREN WITH ATRIAL SEPTA DEFECT TYPE OF CONGENITAL HEART DISEASE UNDER GENERAL ANAESTHESIA

Inne Suherna Sasmita, Anissa Maya Kania, Kirana Lina Gunawan

189-193



ROLE OF PALATAL RUGAE PATTERNS (RUGOSCOPY) IN FORENSIC FIELD
PERAN POLA RUGAE PALATAL (RUGOSKOPI) DALAM BIDANG FORENSIK

Subhaini Jakfar

194-196



PERANAN KASEIN DALAM PENCEGAHAN KARIES GIGI
ROLES OF CASEIN IN DENTAL CARIES PREVENTION

Darmayanti Siregar

197-202



Publisher:



Access Policy:



Reference Manager:



Indexed by:



Accredited by:



Visitor:



[View My Stats](#)

Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

Dentika Dental Journal

Faculty of Dental Medicine
Universitas Sumatera Utara

Jl. Alumni No. 2, Kampus USU Pd. Bulan
Medan 20155, Sumatera Utara - Indonesia
Phone : +62 61 8216131 psw. 21
Fax : +62 61 8213421
Email : dentika_journal@usu.ac.id



This work is licensed under a [Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/).

Platform &
workflow by
OJS / PKP

Open Menu

dentika

DENTAL JOURNAL

[Skip to main content](#)

[Skip to main navigation menu](#)

[Skip to site footer](#)

[Home](#) / Editorial Team

Editorial Team

CHIEF EDITOR

Wilda Hafni Lubis (Universitas Sumatera Utara)

[UNIVERSITY PAGE](#) | [SINTA](#) | [ORCID](#) | [GOOGLE](#)

EDITORIAL TEAM

Tadeus Arufan Jasrin (Universitas Padjadjaran)

[UNIVERSITY PAGE](#) | [SINTA](#) | [SCOPUS](#) | [GOOGLE](#)

Ardo Sabir (Universitas Hasanuddin)

[UNIVERSITY PAGE](#) | [SCOPUS](#) | [GOOGLE SCHOLAR](#)

Pitu Wulandari (Universitas Sumatera Utara)

[UNIVERSITY PAGE](#) | [SINTA](#) | [ORCID](#) | [SCOPUS](#)

Nurdiana (Universitas Sumatera Utara)

[UNIVERSITY PAGE](#) | [SINTA](#) | [GOOGLE SCHOLAR](#)

Darmayanti Siregar (Universitas Sumatera Utara)

[UNIVERSITY PAGE](#) | [SINTA](#) | [ORCID](#) | [SCOPUS](#)

Pocut Astari (Universitas Sumatera Utara)

[UNIVERSITY PAGE](#) | [SINTA](#) | [GOOGLE SCHOLAR](#)

Zahra Kharina Nasution (Universitas Sumatera Utara)



Publisher:



Access Policy:

OPEN  ACCESS

Reference Manager:



Indexed by:





Accredited by:



Visitor:



[View My Stats](#)

Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

Dentika Dental Journal


Faculty of Dental Medicine
Universitas Sumatera Utara

Jl. Alumni No. 2, Kampus USU Pd. Bulan
Medan 20155, Sumatera Utara - Indonesia
Phone : +62 61 8216131 psw. 21
Fax : +62 61 8213421
Email : dentika_journal@usu.ac.id



This work is licensed under a [Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/).

Platform &
workflow by
OJS / PKP



OUTER MEMBRANE PROTEIN OF PORPHYROMONAS GINGIVALIS AS BASIC OF DIAGNOSTIC TOOL IN AGGRESSIVE PERIODONTITIS

(PROTEIN MEMBRAN BAGIAN LUAR PORPHYROMONAS GINGIVALIS SEBAGAI ALAT DIAGNOSTIK PERIODONTITIS AGRESIF)

Desi Sandra Sari*, Candra Bumi**, Yuliana Mahdiyah DA*

* Departement of Periodontic
Faculty of Dentistry, University of Jember
** Faculty of Public Health, University of Jember
Jl. Kalimantan 37 Jember,
E-mail: desisandrasari@yahoo.com

Abstract

Porphyromonas gingivalis is a gram-negative anaerobic bacteria and has been shown previously to be one of the major pathogens in aggressive periodontitis. Outer membrane protein is the major virulence factor of *P. gingivalis* and plays role in the host immune response impair againts *P. gingivalis*, which in turns, causing tissue destruction and bone resorption. This study was aimed to investigate the isolation and characteristic outer membrane protein of *P. gingivalis*. Protein of OMP *P. gingivalis* ATCC 33277 was isolated using sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and than continued by hemagglutination test. The result showed that protein profil in SDS-PAGE of OMP protein was 40 kDa molecular weight dan hemagglutination test was positive in titer of hemagglutination 1/8. In conclusion, outer membrane protein molecular mass of 40kDa produced by *P. gingivalis* is a key virulence factor involved in the co-aggregation activity of *P. gingivalis*

Key words: polyclonal antibody, *Porphyromonas. gingivalis*, outer membrane protein

INTRODUCTION

Periodontitis is a chronic immunoinflammatory disease of the periodontium that results a progressive loss of gingival tissue, the periodontal ligament, and adjacent supporting alveolar bone. Chronic inflammation of the periodontium is initiated by complex subgingival biofilms containing several likely periodontal pathogens. The biofilm generally contains a portion of the gram negative anaerobic commensal microbiota as well as opportunistic pathogens of the oral cavity, including *Porphyromonas gingivalis*.¹

Black-pigmented anaerobe is a pathogen bacteria group in oral cavity and related with gingivitis, periodontitis, endodontic infection and odontogenic abscess. *P. gingivalis* and *P. intermedia* are one of the *black-pigmented* anaerobe bacteria. They are pathogen bacteria in early and advanced periodontal disease.²

Outer membrane protein (OMP) bacteria of *P. gingivalis* caused the increasing of humoral immune response, so it can stimulate inflammatory cytokine

expression, such as TNF- α , IL-1 β , and IL-6 in monocyte and fibroblast of gingiva. It also induced bone resorption activity.³

Imai et al. reseach⁴ showed outer membrane protein found in negative gram bacteria is RagA, RagB and OMPA-like protein. RagB has the strongest virulent ability in periradicular lesion subject. Whole cell of *P. gingivalis* bacteria, cell extract or OMP immunization can reduce periodontal tissue destruction that caused by *P. gingivalis* bacteria. Rat's study showed that 40 kDa OMP antibody is potential to kill *P. gingivalis* bacteria.⁵

The aim of this study was to investigate the isolation and characteristic outer membrane protein of *P. gingivalis* bacteria as the making dot enzyme immunoassay (EIA) material to examine aggressive periodontitis.

MATERIALS AND METHODS

This study was done at the Microbiology Laboratory of Dental Faculty, Jember University and

physiology Laboratory of Medicine Faculty, Brawijaya University. *P. gingivalis* bacteria culture in phosphate buffer saline (PBS) was soaked by *fortex mixer* for 10 seconds, then 0.5 mL. The mixture was taken and cultured on blood medium. Blood medium composition were 2 mL *brucella broth*, 0.4 µl/ml K₁ vitamin, and 5 µl/ml hemin and 10% *sheep blood*. Medium and bacteria were put in *anaerobic jars* then were incubated in 37°C room condition that contained 10% CO₂ by *GasPack CO₂ generating sachet* for 7-14 days. Then it was cultured in biphasic medium (BHI/TCG).

Before OMP isolation, the pili was cut. Bacterial collection result was added with trichloro-acetic-acid (TCA) and centrifuged 6000 rpm for 10 minutes in 4°C. Pellet was taken and resuspended with pH 7.4 PBS and the ratio was 1:10. Bacteria was blend by mixer. Pili in the top part was taken. OMP isolation used Evan's modification method. Sample modification was the top sediment of pili cutting treatment in the last centrifugation, pellet was resuspended in pH 7,4 PBS until its volume reached 15 times, then added with n-octyl B-D-glucopyranoside (NOG) until its concentration reached 0.5%. Supernatant fluid was taken and dialyzed. The fluid was dialyzed on the first 24 hours using H₂O and on the second 24 hours using pH 7.4 PBS.

Monitoring of molecule weight used sodium dodecyl sulphate polyacrylamide gel electrophoresis. Protein samples were heated 100°C for 5 minutes in buffer solution that contained 5 mM Tris HCl pH 6,8-2 mercapto etanol 5%, w/v sodium dodecyl sulfate 2,5%, v/v gliserol 10% with detection color bromophenol blue. Selected mini-slab gel of 12.5% with 4% tracking gel used 125 mV voltages. Material used *coomassive brilliant blue* material and standard molecule sigma *low range marker*.

The results of SDS-PAGE gel straight cut at 40 kDa were collected and included in the dialysis membrane and performed electroelusion and dialysis. Dilution of samples were prepared with a concentration 1/2 on microplate V, the volume of each well 50µl. Each of the wells murine blood concentration of 0.5% was placed in room temperature until 1 hour. The amount of titer was determined by observation of blood agglutination.

OMP 40 kDa protein in wistar rats was injected by intra-peritoneal injection. The first injection was mixed with complete *Freund's adjuvant* (Booster I). The second and further were mixed with incomplete *Freund's adjuvant* for 1 week (Booster II). 40 kDa protein antigen injection *P.gingivalis* until booster V. After the wistar rats were injected ketamine, blood was collected from the heart. Blood di-

sentifuse 1500 rpm for 15 minutes. Serum taken from the supernatant, obtained polyclonal antibodies anti-40 kDa OMP of *P. gingivalis* serum from wistar rats.

RESULTS

After culturing in blood medium, we obtained the result of colony of *P. gingivalis* bacteria. Figure 1 showed the results of bacterial culture isolation of *P. gingivalis* ATCC 33277 on day 14, which on this day the bacterial harvesting of isolated OMP. Colonies of these bacteria have a characteristic brown or black pigment on blood agar.




Figure 1. Bacterial colony of *P. gingivalis* ATCC 33277 on the 14th day




Figure 2. Result of SDS-PAGE of OMP *P. gingivalis*
1. Protein marker, 2. OMP *P. gingivalis*

Protein profile on SDS-PAGE of OMP *P. gingivalis* showed 40 kDa protein. Lanes 1 is a protein marker and lanes 2 is the OMP *P.gingivalis* with protein band is 40 kDa. Hemagglutination test results used eritrosit of mice can be seen in Table 1.

Table 1. Hemagglutination test 40 kDa OMP protein of *P. gingivalis* ATCC 33277

Molecular Weight Proteins (kDa)	Dilution										
	1/2	1/4	1/8	1/16	1/32	1/64	1/128	1/256	1/512	1/1024	K(-)
40	+	+	+	-	-	-	-	-	-	-	-

(+): No Coagulation, (-): Coagulation, K(-): Control negative




Figure 3. The result hemagglutination test against 40 kDa OMP *P. gingivalis*

Table 1 and Figure 3 Showed that protein with a 40 kDa molecular weight provide the highest hemagglutinin titer of 1/8. Hemagglutination protein is considered as one of the virulence factors of pathogenic bacteria and as a basis for making antibodies.

DISCUSSION

Porphyromonas species was an anaerobic *black-pigmented* bacteria that is pathogen and was related specifically with advanced periodontitis lesion. *P. gingivalis* was a virulent bacteria in vivo and in vitro, and almost found in subgingival plaque of periodontitis subject. This species had brown or black pigmented characteristic in blood medium.⁶

The amount of anaerobic *black-pigmented* bacteria in aggressive periodontitis subject and healthy subject showed that was significant different in both of the groups. The amount of *black-pigmented* bacteria in aggressive periodontitis subject was higher than healthy subject.⁷ Subsequently, amount of *black-pigmented* bacteria in aggressive periodontitis subject had strong positive relationship with pocket depth, that deepest pocket had higher amount *black-pigmented* bacteria.⁸

Porphyromonas gingivalis had virulent factors such as lipopolysaccharide (LPS), fimbriae, toxic metabolic product, and protease. Virulent factors can cause disease directly or not by host cell activation to release inflammatory mediators.⁹ *Porphyromonas gingivalis* can produce protease extracellular, such as arginine-specific *protease* and lysine-specific *protease*, both of them were coded as polyprotein that contained *protease* and *adhesin* domain. Both of the *proteases* had C-terminal *adhesin* domain that was involved in attachment of host protein.¹⁰

Forty kDa molecule weight of outer membran protein of *P. gingivalis* was virulent factor for co-

aggregation. Forty kDa outer membran protein was found in some strains of *P. gingivalis*. It also described that anti opsonisation of 40 kDa-OMP antibody of *P. gingivalis* was as phagocyte target of neutrophil cell, so several studies indicated that 40 kDa OMP can be used as candidate of anti vaccine of periodontitis.¹¹ It can be concluded that outer membran protein of *P. gingivalis* bacteria can be used as polyclonal antibody of making diagnostic tool of aggressive periodontitis in 40 kDa molecule weight.

ACKNOWLEDGEMENTS

Ministry of National Education Republic of Indonesia, in this case through the University of Jember Research DIPA that has funded this research and the Research Institute of the University of Jember, which facilitates the study.

References

1. Pihlstrom BS, Michalowicz, Johnson. Periodontal diseases. *Lancet* 2005; 366 (9499): 1809-20.
2. Soukos NS, Som S, Abernethy AD, Ruggiero A, Dunham J, Lee C, Doukas AG, Goodson JM. Phototargeting oral black-pigmented bacteria. *Antimicrobial Agent and Chemotherapy* 2005; 49(4): 1391-6.
3. Kuehn M, Kesty N. Bacterial outer membrane vesicles and the host-pathogen interaction. *Genes and Development* 2005; 19: 2645-55.
4. Imai M, Murakami Y, Nagano K, Nakamura H, Yoshimura F. Major outer membrane proteins from *Porphyromonas gingivalis*: strain variation, distribution, and clinical significance in periradicular lesions. *European Oral Sciences* 2005; 113(5): 391-9.
5. Tsurumi Y, Hayakawa M, Shibata Y, Abiko Y. Production of antibody against a synthetic peptide of *Porphyromonas gingivalis* 40-kDa outer membrane protein. *Oral Science* 2003; 45(2): 111-6.
6. Samaranayake LP. *Essential microbiology for dentistry* 2nd ed. Philadelphia: Churchill Livingstone, 2002:122-3.
7. Sari DS. Jumlah bakteri black-pigmented anaerob cairan sulkus gingival pada penderita aggressive periodontitis. *Spirulina* 2010; 5(1):53-66.
8. Sari DS. Calprotectin in gingival crevicular fluids correlates with clinical parameter and the count of black-pigmented anaerobes bacteria on the aggressive periodontitis. Poster Presentation 6th FDI-

- IDA Joint Meeting, 2010: 31.
9. Sugawara S, Nemoto E, Tada H, Miyake K, Imamura T, Takada H. Proteolysis of human monocyte CD 14 by cystein proteinase (gingipains) from Porphyromonas gingivalis leading to lipopolysaccharide hyporesponsiveness, Immunology 2000; 165: 411-8.
 10. Mars P. Martin MV. Oral microbiology, 4th ed., Oxford: Wright, 2000: 195.
 11. Maeba S, Otake S, Namikoshi J, Hayakawa M. Transcutaneous immunization with Porphyromonas gingivalis surface protein antigen induces T helper 2 responses in systemic tissue. Int J Oral-Med Sci 2004; 3(2): 67-74.

