






# Barekeng

jurnal ilmu matematika dan terapan

pISSN 1978 - 7227 | eISSN 2615 - 3017

## CALL FOR PAPER

**S2**  
sinta  
Science and Technology Index

### Focus & Scope

Pure Mathematics , Applied Mathematics, Statistics, Actuarial Science, Logic, Geometry & Topology, Numerical Analysis, Mathematic Computation and Mathematics Education.

Vol 17 No 1, March 2023  
Submission Deadline: 30<sup>th</sup> November 2022

Indexed by :










 <https://ojs3.unpatti.ac.id/index.php/barekeng>
 [barekeng.math@yahoo.com](mailto:barekeng.math@yahoo.com)
 085243358669

### Journal Identity

Original journal title	BAREKENG : Jurnal Ilmu Matematika dan Terapan
English journal title	BAREKENG : Journal of Mathematics and Its Application
Country	Indonesia
Subject	Mathematics, Applied Mathematics
Language	English
ISSN	<u>2615-3017</u> (online), <u>1978-7227</u> (print)
Frequency	4 issues per year (March, June, September, December)
DOI	Prefix 10.30598/ <u>Crossref</u>
Acreditation	<u>SINTA 2</u> , Decree Number: 158/E/KPT/2021, 9th December 2021
Editor-in-Chief	Yopi Andry Lesnussa [ <u>Sinta</u> ][ <u>Scopus</u> ][ <u>Google Scholar</u> ]
Publisher	<u>Pattimura University</u> .
Citation Analysis	<u>Google Scholar</u>   <u>Dimensions</u>
OAI	<a href="https://ojs3.unpatti.ac.id/index.php/barekeng/oai">https://ojs3.unpatti.ac.id/index.php/barekeng/oai</a>

## EDITOR IN CHIEF



**Yopi Andry Lesnussa, M.Si.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

## ASSOCIATE EDITOR



**Muhammad Yahya Matdoan, M.Si.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)



**Meilin Imelda Tilukay, M.Si**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)



**Berny Pebo Tomasouw, M.Si**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

## EDITORIAL BOARD

**Prof. Dr. Basuki Widodo, M.Sc.**  
ITS Surabaya, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Prof. Dr. Atje Setiawan Abdullah, MS., M.Kom.**  
Universitas Padjadjaran, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Subchan, M.Sc., Ph.D.**  
ITS Surabaya, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Sobri Abusini, MT.**  
Universitas Brawijaya, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Rr. Kurnia Novita Sari, M.Si.**  
Institut Teknologi Bandung, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Hengameh R. Dehkodi, Ph.D.**  
Universidade Federal do ABC, Brazil  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

**Prof. Dr. Budi Nurani Ruchjana, M.Si.**  
Universitas Padjadjaran, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Prof. Dr. Guisheng Zhai**  
Shibaura Institute of Technology, Japan  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

**Dr. Fajar Adi Kusumo, M.Si.**  
Universitas Gadjah Mada, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Sumardi, M.Si.**  
Universitas Gadjah Mada, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Yuwadee Klomwises**  
King Mongkul's Institute of Technology, Thailand  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

**Dr. Antonio Comi**  
Universita degli Studi di Roma "Tor Vergata", Roma  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

**Prof. Dr. Shakir Ali**  
Aligarh Muslim University, India  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

**Prof. Drs. Marjono, M.Phil., Ph.D.**  
Universitas Brawijaya, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Prof. Dr. Tanwey Gerson Ratumanan, M.Pd**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Ir. Bib Paruhum Silalahi, M.Kom.**  
Institut Pertanian Bogor, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Ikha Magdalena, M.Si.**  
Institut Teknologi Bandung, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Prof. Dr. Theresia Lourens, M.Pd.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dr. Ahmad Fadillah bin Embong**  
Universiti Teknologi Malaysia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#)

## ASSISTANT EDITORS

**Norisca Lewaherilla, M.Si.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Jefri Esna Thomas Radjabaycolle, M.Cs.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Venn Yan Ishak Ilwaru, M.Si**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

**Dyana Patty, M.Sc.**  
Universitas Pattimura, Indonesia  
[\[Scopus ID\]](#) [\[Google Scholar ID\]](#) [\[SINTA ID\]](#)

## Editorial Address:



**BAREKENG: Jurnal ilmu matematika dan terapan** (*Original name*)  
**BAREKENG: Journal of Mathematics and Its Applications** (*English name*)  
 Ex. UT Building, 2<sup>nd</sup> Floor, Pattimura University

Ir. M. Putuhera Street, Kampus Unpatti, Poka – Ambon City, 97233, Maluku Province, Indonesia

<https://ojs3.unpatti.ac.id/index.php/barekeng/>

+62 852 4335 8669 (Yopi)

+62 823 9798 0021 (Dyana)

[barekeng.math@yahoo.com](mailto:barekeng.math@yahoo.com)

[barekeng.jurmath@gmail.com](mailto:barekeng.jurmath@gmail.com)


[barekeng.journal@mail.unpatti.ac.id](mailto:barekeng.journal@mail.unpatti.ac.id)

Jurnal Barekeng

jurnal\_barekeng

## Current Issue

### Vol 16 No 4 (2022): BAREKENG: Jurnal Ilmu Matematika dan Terapan



PATTIMURA UNIVERSITY

BAREKENG : Journal of Mathematics and Its Applications, volume 16 issue 4 (December 2022) was officially published in English (also journal name officialy in english).

BAREKENG : Journal of Mathematics and Its Applications, has been published four (4) times a year in March, June, September, and December (4 issues). This issue contain 40 articles available online. Special for this issue consists of authors from 2 countries (Indonesia, and Malaysia).

**PUBLISHED:** 2022-12-15

## Articles

### THE HARMONIOUS, ODD HARMONIOUS, AND EVEN HARMONIOUS LABELING

Ahmad Lasim, Ikhsanul Halikin, Kristiana Wijaya

1131-1138

pdf download

Abstract views 12 | Pdf Download downloads 13 |

### ANALYSIS OF OPTIMUM CONTROL ON THE IMPLEMENTATION OF VACCINATION AND QUARANTINE ON THE SPREAD OF COVID-19

Agusyarif Rezka Nuha, Novianita Achmad, Gusti Arviana Rahman, Syarif Abdullah, Sri Istiyarti Uswatun Chasanah, Nina Valentika, La Ode Nashar

1139-1146

pdf download

Abstract views 65 | Pdf Download downloads 24 |

### FLUID FLOW MODELLING WITH FREE SURFACE

Anjeryan Sapta Pratama, Evi Noviani, Yudhi Yudhi

1147-1158

pdf download

Abstract views 9 | Pdf Download downloads 10 |

### MODIFICATION OF POLLARD RHO ALGORITHM USING NEGATION MAPPING

Sa'aadah Sajjana Carita, Herman Kabetta

1159-1166

pdf download


Abstract views 7 | Pdf Download downloads 7 |



### HILL CLIMBING ALGORITHM ON BAYESIAN NETWORK TO DETERMINE PROBABILITY VALUE OF SYMPTOMS AND EYE DISEASE

Ria Puan Adhitama, Dewi Retno Sari Saputro, Sutanto Sutanto

1271-1282

 pdf download

 Abstract views 8 |  Pdf Download downloads 11 |

### ANALYSIS OF BANKING DEPOSIT COST IN THE DYNAMICS OF LOAN: BIFURCATION AND CHAOS PERSPECTIVES

Moch. Fandi Ansori, Susilo Hariyanto

1283-1292


 pdf download

 Abstract views 7 |  Pdf Download downloads 11 |

### MULTI-STATE MODEL FOR CALCULATION OF LONG-TERM CARE INSURANCE PRODUCT PREMIUM IN INDONESIA

Hendra Perdana, Neva Satyahadewi, Dadan Kusnandar, Ray Tamtama

1293-1302

 pdf download


 Abstract views 12 |  Pdf Download downloads 11 |

### MAGNETOHYDRODYNAMICS NANOFERRO FLUID FLOWS PASSING THROUGH A MAGNETIC POROUS SPHERE UNDER THERMAL RADIATION EFFECT

Basuki Widodo, Eirene Juwita Ningtyas Pamela, Dieky Adzkiya, Chairul Imron, Tri Rahayuningsih

1303-1312


 pdf download



 Abstract views 9 |  Pdf Download downloads 8 |

### THE PROMINENCE OF VECTOR AUTOREGRESSIVE MODEL IN MULTIVARIATE TIME SERIES FORECASTING MODELS WITH STATIONARY PROBLEMS

Embay Rohaeti, I Made Sumertajaya, Aji Hamim Wigena, Kusman Sadik

1313-1324


 pdf download



 Abstract views 6 |  Pdf Download downloads 8 |

### OPTIMIZATION OF PORTFOLIO USING FUZZY SELECTION

Rahmania Ayu Wardani, Sugiyarto Surono, Goh Kang Wen

1325-1336


 pdf download

 Abstract views 6 |  Pdf Download downloads 8 |

### CREDIT CARD FRAUD DETECTION USING LINEAR DISCRIMINANT ANALYSIS (LDA), RANDOM FOREST, AND BINARY LOGISTIC REGRESSION

Muhammad Ahsan, Tabita Yuni Susanto, Tiza Ayu Virania, Andi Indra Jaya

1337-1346


 pdf download

 Abstract views 15 |  Pdf Download downloads 10 |

### A NOTE ON THE SOLUTION OF THE CHARACTERISTIC EQUATION OVER THE SYMMETRIZED MAX-PLUS ALGEBRA

Gregoria Ariyanti

1347-1354

 pdf download

 Abstract views 7 |  Pdf Download downloads 8 |

## THE HARMONIOUS, ODD HARMONIOUS, AND EVEN HARMONIOUS LABELING

Ahmad Lasim<sup>1</sup>, Ikhsanul Halikin<sup>2</sup>, Kristiana Wijaya<sup>3\*</sup>

<sup>1,2,3</sup>Graph and Algebra Research Group, Mathematics Department,  
Faculty of Mathematics and Natural Sciences, Universitas Jember  
Jl. Kalimantan 37 Jember, 68121, Jawa Timur, Indonesia

Corresponding author's e-mail: <sup>3\*</sup> [kristiana.fmipa@unej.ac.id](mailto:kristiana.fmipa@unej.ac.id)

---

**Abstract.** Suppose  $G$  is a simple and connected graph with  $q$  edges. A harmonious labeling on a graph  $G$  is an injective function  $f: V(G) \rightarrow \{0, 1, 2, \dots, q-1\}$  so that there exists a bijective function  $f^*: E(G) \rightarrow \{0, 1, 2, \dots, q-1\}$  where  $f^*(uv) = f(u) + f(v) \pmod{q}$ , for each  $uv \in E(G)$ . An odd harmonious labeling on a graph  $G$  is an injective function  $f$  from  $V(G)$  to non-negative integer set less than  $2q$  so that there is a function  $f^*(uv) = f(u) + f(v)$  where  $f^*(uv) \in \{1, 3, 5, \dots, 2q-1\}$  for every  $uv \in E(G)$ . An even harmonious labeling on a graph  $G$  is an injective function  $f: V(G) \rightarrow \{0, 1, 2, \dots, 2q\}$  so that there is a bijective function  $f^*: E(G) \rightarrow \{0, 2, 4, \dots, 2q-2\}$  where  $f^*(uv) = f(u) + f(v) \pmod{2q}$  for each  $uv \in E(G)$ . In this paper, we discuss how to build new labeling (harmonious, odd harmonious, even harmonious) based on the existing labeling (harmonious, odd harmonious, even harmonious).

**Keywords:** even harmonious labeling, harmonious labeling, odd harmonious labeling.

---

**Article info:**

Submitted: 31<sup>st</sup> January 2022

Accepted: 2<sup>nd</sup> October 2022

**How to cite this article:**

A. Lasim, I. Halikin and K. Wijaya, "THE HARMONIOUS, ODD HARMONIOUS, AND EVEN HARMONIOUS LABELING", *BAREKENG: J. Math. & App.*, vol. 16, iss. 4, pp. 1131-1138, Dec., 2022.

---



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).  
Copyright © 2022 Author(s)



## 1. INTRODUCTION

Sedlacek first introduced graph labeling in 1963. Graph labeling is a function that maps a set of vertices or edges of a graph to a non-negative integer that fulfills specific properties/rules. This particular nature/rule leads to different types of labeling. Thousands of papers on graph labeling have been produced. Gallian [1] surveyed this. In this paper, the research focus is on harmonious, odd harmonious labeling, and even harmonious labeling, all three of which are vertex labeling.

A harmonious labeling was first introduced by Graham and Sloane [2] in 1980. A graph  $G$  with  $q$  edges is said to be harmonious if there is a one-to-one (injective) function  $f$  that maps the set of vertices  $V(G)$  to the set  $\{0, 1, 2, \dots, q-1\}$  so that each edge gets the label of the sum of the two vertices labels incident with that edge in modulo  $q$ , which are all different. When  $G$  is a tree,  $G$  has  $(q + 1)$  vertices, so it is clear that the one-to-one function condition is not satisfied. Some researchers give exceptions to trees by changing the existence of a one-to-one function into a surjective function [1]. Therefore, in this paper, the focus of the discussion is only on connected graphs that are not trees. It means that the graph always contains a closed path. Not many graphs can be labeled harmoniously [1]. Graham and Sloane [2] proved that only a cycle graph with an odd length is a harmonious graph; the wheel  $W_n$  is a harmonious graph for every  $n$ . Other results regarding harmonious labeling can be seen in [1].

Furthermore, harmonious labeling inspires the emergence of odd and even harmonious labeling. It is said to be odd harmonious because the label of each edge must be odd. It is said to be even harmonious because all of its edges must be labeled even. Odd harmonious labeling was introduced by Liang and Bai in 2009. A graph  $G$  with  $q$  edges is said to be odd harmonious if there is a one-to-one function  $f$  that maps every element in  $V(G)$  to the set of non-negative integers less than  $2q$ , so that each edge gets an odd label that is all different from the set  $\{1, 3, 5, \dots, 2q-1\}$ , which is obtained from the sum of the two vertices labels incident to that edge. The following is a theorem related to odd harmonious labeling.

**Theorem 1.** (Liang and Bai [3])

- (1) *If  $G$  is an odd harmonious graph, then  $G$  is a bipartite graph.*
- (2) *Let  $G$  be a graph with  $p$  vertices and  $q$  edges. If  $G$  is an odd harmonious graph, then  $2\sqrt{q} \leq p \leq 2q-1$ .*

Several other studies on odd harmonious labeling that have been carried out can be seen, among others: [4], [5], [6], [7], [8], [9], [10].

An even harmonious labeling of a graph was introduced by Sarasija and Binthiya [11] in 2011. The graph  $G$  is an even harmonious if there is a one-to-one function  $f: V(G) \rightarrow \{0, 1, 2, \dots, 2q\}$  so that each edge gets a different even label in the set of integers modulo  $2q$  as the result of the sum of the two labels of the vertices incident to that edge. Even harmonious graphs that have been studied can be seen in [11], [12], [13], [14], [15].

The graph studied in this paper was a simple and connected. The basic theory for the graph can be seen in Chartrand and Zhang [16]. In this paper, we study more deeply about harmonious, odd harmonious, and even harmonious labeling on a graph. The previous studies emphasize more on the class of graphs studied. Then, this study emphasizes the labeling itself, namely how to build a new label based on the current labeling. In particular, if it is given labeling of harmonious, odd harmonious, or even harmonious on a graph, a new label will be built based on that labeling so that the resulting labeling fulfills the labeling requirements for harmonious, odd harmonious, or even harmonious labeling.

## 2. RESEARCH METHOD

The method in this research was exploratory research which aimed to build new labeling from the current labeling of harmonious, odd, and even harmonious. The research was begun by analyzing the definitions of the three types of labeling and what properties each labeling must fulfill. Next, a new label based on the existing labeling was built. The final step was proving that the new labeling met the rules of the labeling, whether the labeling was harmonious, odd harmonious, or even harmonious.



### 3. RESULT AND DISCUSSION

In this section, three types of vertex labeling were discussed, namely harmonious labeling, odd harmonious, and even harmonious labeling. From the three labels given, a new function was built based on the existing labeling so that the resulting labeling meets the labeling properties of harmonious, odd harmonious, or even harmonious labeling.

#### 3.1. Harmonious Labeling

A harmonious labeling on a graph  $G$  with  $q$  edges is a one-to-one function (injective)  $f: V(G) \rightarrow \{0, 1, 2, \dots, q-1\}$  so that it produces a one-to-one correspondence (bijective function)  $f^*: E(G) \rightarrow \{0, 1, 2, \dots, q-1\}$  with  $f^*(uv) = (f(u) + f(v)) \pmod{q}$ , for each  $uv \in E(G)$ . A graph is said to be harmonious if it can be labeled with the harmonious labeling rule. The following two theorems explain how to construct a new harmonious label based on the known harmonious labeling.

**Theorem 2.** Let  $G$  be a graph with  $q$  edges and  $f$  be a harmonious labeling of  $G$ . Suppose that the function  $g_f: V(G) \rightarrow \{0, 1, \dots, q-1\}$  with  $g_f(u) = q - f(u)$  for each  $u \in V(G)$ . Therefore,  $g_f$  is a harmonious labeling on the graph  $G$ .

**Proof.** Suppose  $f$  is a harmonious labeling on a graph  $G$  with  $q$  edges and the function  $g_f: V(G) \rightarrow \{0, 1, \dots, q-1\}$  with  $g_f(u) = q - f(u)$  for each  $u \in V(G)$ . First, it will be shown that  $g_f$  is a one-to-one function. Since  $f$  is a one-to-one function, it is clear that any two distinct vertices in  $G$  will be labeled differently by the  $g_f$  function. Thus  $g_f$  is a one-to-one function.

Then, it will be proven that for each edge  $uv \in E(G)$  get a different label from  $\{0, 1, 2, \dots, q-1\}$  by the rules  $g_f^*(uv) = g_f(u) + g_f(v) \pmod{q}$ .

$$\begin{aligned} g_f^*(uv) &= (g_f(u) + g_f(v)) \pmod{q} \\ &= ((q - f(u)) + (q - f(v))) \pmod{q} \\ &= 2q - (f(u) + f(v)) \pmod{q} \\ &= 2q - f^*(uv) \pmod{q} \\ &= -f^*(uv) \pmod{q} \end{aligned}$$

This means  $g_f^*$  is the summation inverse of  $f^*$  on the set of integers modulo  $q$ . This shows that each edge  $uv \in E(G)$  get a different label from  $\{0, 1, 2, \dots, q-1\}$ . Because each vertex in  $G$  gets a different label with the function rule  $g_f$ , each edge in  $G$  also gets a different label in  $\{0, 1, 2, \dots, q-1\}$  by the rule  $g_f^*(uv) = g_f(u) + g_f(v) \pmod{q}$ . Therefore, it is proven  $g_f$  is a harmonious labeling on the graph  $G$ . ■

For example, in Figure 1 (a), it is given a labeling of a wheel with 8 edges,  $W_4$ . Furthermore, if the label of vertices in Figure 1 (a) is changed by the rule  $8 - f(u)$  for each  $u \in V(W_4)$ , it will produce the harmonious labeling in Figure 1 (b).




Figure 1. Two harmonious labelings of wheel  $W_4$  to illustrate Theorem 2

**Theorem 3.** Suppose the function  $f$  is a harmonious labeling on a graph  $G$  with  $q$  edges. Then the function  $g_f: V(G) \rightarrow \{0, 1, \dots, q-1\}$  with  $g_f(u) = (f(u) + k) \pmod{q}$  for each  $u \in V(G)$  and  $k \in [1, q-1]$  is a harmonious labeling on the graph  $G$ .



**Proof.** First, it will be proved that  $g_f$  is a one-to-one function of the set  $V(G)$  to the set  $\{0, 1, 2, \dots, q - 1\}$ . Take any  $u, v \in V(G)$  that fulfills  $g_f(u) = g_f(v)$ . Therefore,  $(f(u) + k) \pmod q = (f(v) + k) \pmod q$  for  $k \in \{1, 2, \dots, q - 1\}$ . Thus, we get  $f(u) = f(v)$ . Because  $f$  is a one-to-one function, then  $u = v$ . Therefore,  $g_f$  is a one-to-one function.

Next, we will prove that with the function  $g_f^*$ , each edge in  $G$  gets a different label from  $0, 1, 2, \dots, q - 1$ . Suppose there are two different edges  $uv$  and  $u_0v_0$  in  $E(G)$  which has the same label, i.e.  $g_f^*(uv) = g_f^*(u_0v_0)$ . It is obtained  $(f^*(uv) + 2k) \pmod q = (f^*(u_0v_0) + 2k) \pmod q$ . Consequently,  $f^*(uv) = f^*(u_0v_0)$ . This is contrary to  $f^*$  one-to-one function. It is proven that  $g_f^* : V(G) \rightarrow \mathbb{Z}_q$  is bijective function. Thus,  $g_f$  is a harmonious labeling on the graph  $G$ . ■

For example, consider the harmonious labeling of the 15-edges Petersen graph in Figure 2(a). If the labels of each vertex in Figure 2 (a) are added to 5 in modulo 15, then the harmonious labeling is obtained as in Figure 2 (b).




Figure 2. Two harmonious labelings of a Petersen graph to illustrate Theorem 3

### 3.2. Odd Harmonious Labeling

An odd harmonious labeling on a graph  $G$  with  $q$  edges is a one-to-one function  $f: V(G) \rightarrow \{0, 1, 2, \dots, 2q - 1\}$  so that there is a one-to-one correspondence  $f^*: E(G) \rightarrow \{1, 3, 5, \dots, 2q - 1\}$  with  $f^*(uv) = f(u) + f(v)$  for each  $uv \in E(G)$  [8]. The following is a theorem related to odd harmonious labeling. According to Theorem 1, an odd harmonious graph is a bipartition graph. Since a bipartition graph does not contain odd cycles, therefore an odd harmonious graph will not contain odd cycles. In addition, based on the definition of odd harmonious labeling that each edge  $e = uv$  gets a label from the number of labels of two vertices incident to that edge, it is clear that the edge labeled 1 is only obtained from the vertex with label 0 and the vertex with label 1. Thus, the vertex with a label 0 must be adjacent to the vertex with label 1. Furthermore, the following theorem discusses odd harmonious labeling, which is built from existing odd harmonious labeling.

**Theorem 4.** Let  $G$  be a graph with  $q$  edges. If  $f$  is an odd harmonious labeling on a graph  $G$ , then the function  $g_f: V(G) \rightarrow \{0, 1, \dots, 2q - 1\}$  which is defined as

$$g_f(u) = \begin{cases} f(u) - 1 & \text{if } f(u) \text{ odd,} \\ f(u) + 1 & \text{if } f(u) \text{ even,} \end{cases}$$

for each  $u \in V(G)$  is an odd harmonious labeling on graph  $G$  with the same edge label as labeling  $f$ .

**Proof.** First, it will be shown that the function  $g_f$  is one-to-one. Based on the definition of  $g_f$ , label  $g_f(u)$  is odd if and only if  $f(u)$  is even and vice versa, label  $g_f(u)$  is even if and only if  $f(u)$  is odd. Thus, since  $f$  is a one-to-one function, obviously a function  $g_f$  is also one-to-one. Next, it will be proved that each edge  $uv \in E(G)$  gets odd labels that are all different from  $1, 3, \dots, 2q - 1$  by the rule  $g_f^*(uv) = g_f(u) + g_f(v)$ . Because  $g_f^*(uv)$  must be odd, one of  $g_f(u)$  and  $g_f(v)$  must be odd and others even. Based on the function definition  $g_f$ , label  $g_f(u)$  is odd if and only if  $f(u)$  is even and vice versa, label  $g_f(u)$  is even if and only if  $f(u)$  is odd. Without loss of generality, if  $g_f(u)$  is odd and  $g_f(v)$  is even, it is obtained  $g_f^*(uv) = g_f(u) + g_f(v) = (f(u) - 1) + (f(v) + 1) = f(u) + f(v) = f^*(uv)$ . Because  $f^*$  is a





bijjective function that maps each edge  $uv \in E(G)$  to the set  $\{1, 3, \dots, 2q - 1\}$ , so that  $g_f^*$  is also a bijective function. Therefore, it is proven that  $g_f$  is an odd harmonious labeling on a graph  $G$ , with the same edge label as the function  $f$ . ■

For example, in Figure 3 (a), it is given an odd harmonious label to a star  $S_8$ . The odd harmonious labeling in Figure 3 (b) is obtained from Figure 3 (a) with the rules written in Theorem 4, namely the odd label is minus one, while the even label is added one.




Figure 3. Two odd harmonious labelings with the same edge label on a star to illustrate Theorem 4

### 3.3. Even Harmonious Labeling

An *even harmonious labeling* on a graph  $G$  is a one-to-one function  $f: V(G) \rightarrow \{0, 1, 2, \dots, 2q\}$  so that there is a one-to-one correspondence  $f^*: E(G) \rightarrow \{0, 2, 4, \dots, 2q - 2\}$  with  $f^*(uv) = (f(u) + f(v)) \pmod{2q}$  for each edge  $uv \in E(G)$  [13]. A graph  $G$  is said to be even harmonious if the set of vertices and edges in graph  $G$  can be labeled with an even harmonious labeling rule. It is said to be even harmonious because all of its edges are labeled even. Because even numbers are obtained from the sum of both even or odd numbers, all vertices must be labeled even or all vertices must be labeled odd in this even harmonious labeling. Thus, when  $G$  is a tree, all vertices of  $G$  must be labeled even. In the following two theorems, it is proved that if  $f$  is even harmonious labeling on a graph  $G$  with  $q$  edges, then the two labels constructed from  $f$ , namely for each  $u \in V(G)$ ,  $g_f(u) = (f(u) + k) \pmod{2q}$  for  $k \in [1, 2q - 1]$  or  $g_f(u) = 2q - f(u)$ , are even harmonious labeling.

**Theorem 5.** Let  $G$  be a graph that has  $q$  edges. Suppose  $f$  is an even harmonious labeling on  $G$  and the function  $g_f: V(G) \rightarrow \{0, 1, \dots, 2q\}$  is defined as  $g_f(u) = (f(u) + k) \pmod{2q}$  for each  $u \in V(G)$  and  $k \in [1, 2q - 1]$ . If  $f: V(G) \rightarrow \{1, 2, \dots, 2q - 1\}$  so  $g_f$  is an even harmonious labeling on graph  $G$ .

**Proof.** Suppose  $f: V(G) \rightarrow \{1, 2, \dots, 2q - 1\}$  is even harmonious labeling on a graph  $G$ . Suppose again that  $g_f(u) = (f(u) + k) \pmod{2q}$  for each  $u \in V(G)$  and  $k \in [1, 2q - 1]$ . Suppose  $g_f$  is not one-to-one function, there are two distinct vertices  $u$  and  $v$  in  $G$  so that  $g_f(u) = g_f(v)$ , namely  $(f(u) + k) \pmod{2q} = (f(v) + k) \pmod{2q}$  for some  $k \in [1, 2q - 1]$ . Since the sum is in modulo  $2q$ , this equation is fulfilled when  $f(u) = 0$  and  $f(v) = 2q$  or vice versa. However, this contradicts to  $f: V(G) \rightarrow \{1, 2, \dots, 2q - 1\}$ . Therefore,  $g_f$  must be one-to-one function. Next, we consider  $g_f^*: E(G) \rightarrow \{0, 2, \dots, 2q - 2\}$  with  $g_f^*(uv) = (g_f(u) + g_f(v)) \pmod{2q}$  for each edge  $uv \in E(G)$ . Therefore,  $g_f^*(uv) = (g_f(u) + g_f(v)) \pmod{2q} = (f(u) + f(v) + 2k) \pmod{2q} = (f^*(uv) + 2k) \pmod{2q}$ . This shows that the edge label is based on the function rule  $g_f^*$  different to  $2k$  with edge label based on the function rules  $f$ . As a note, the edge label of both functions will be the same when  $k = q$ . ■

As an illustration, in the Figure 4 is given an even harmonious labeling on cycle  $C_5$  with different vertex label but produces the same edge labels. The labeling of the vertices in Figure 4 (b) is obtained from the labeling rules in Theorem 5, namely the labeling of the vertices in Figure 4 (a) plus 5 in modulo 10, while the labeling of the edges do not change.




Figure 4. Two Even harmonious labelings with different vertex labels on a cycle  $C_5$

**Theorem 6.** Suppose  $f$  is an even harmonious labeling on a graph  $G$  with  $q$  edges. Then, the function  $g_f: V(G) \rightarrow \{0, 1, \dots, 2q\}$  with  $g_f(u) = 2q - f(u)$  for each  $u \in V(G)$  is an even harmonious labeling on the graph  $G$ .

**Proof.** Since  $f$  is a one-to-one function from the set of vertices  $V(G)$  to the set of labels  $\{0, 1, \dots, 2q\}$ , it is obvious that function  $g_f(u) = 2q - f(u)$  for each  $u \in V(G)$ , is one-to-one. Next, for each edge  $uv \in E(G)$  will get label  $g_f^*(uv) = (g_f(u) + g_f(v)) \pmod{2q} = (2q - f(u) + 2q - f(v)) \pmod{2q} = (4q - (f(u) + f(v))) \pmod{2q} = - (f(u) + f(v)) \pmod{2q}$ . This means the edge label based on function rules  $g_f$  is the inverse of addition in modulo  $2q$  of the edge label according to the rule of function  $f$ . Thus, each edge in  $G$  gets a different label and even. Therefore, it is proven that  $g_f$  is an even harmonious labeling on the graph  $G$ . ■

As an illustration of Theorem 6, if given an even harmonious labeling in Figure 5 (a), then we can construct an even harmonious labeling by rule  $10 - f(u)$  for each vertex label, as shown in Figure 5 (b). It can be seen that the edge label on the graph in Figure 5 (b) is the inverse of the integer sum modulo 10 of the edge label on the graph in Figure 5 (a).




Figure 5. Two even harmonious labelings on a 5-edges graph to illustrate Theorem 6

#### 4. CONCLUSION

In this paper, we have constructed two new labelings (harmonious and even harmonious) from the existing (harmonious and even harmonious) labeling. However, only one new labeling (odd harmonious) is obtained from the existing (odd harmonious) labeling. It is due to the edge label in odd harmonious obtained from the addition of two integers that are not in modulo  $2q$ , with  $q$  is the number of edges in the graph, so the vertex labeled zero must be adjacent to the vertex labeled one. However, it is possible to construct another odd harmonious labeling. Therefore, this study is still open to other researchers to construct a new labeling from the existing labeling according to the harmonious labeling.



## REFERENCES

- [1] J. A. Gallian, "A Dynamic Survey of Graph Labeling," 2017.
- [2] R. L. Graham and N. J. A. Sloane, "On Additive Bases and Harmonious Graphs," *SIAM Journal on Algebraic Discrete Methods*, vol. 1, no. 4, pp. 382–404, Dec. 1980, doi: 10.1137/0601045.
- [3] Z.-H. Liang and Z.-L. Bai, "On the odd harmonious graphs with applications," *J. Appl.Math. Comput*, vol. 29, pp. 105–116, 2009.
- [4] M. Bača and M. Z. Youssef, "On harmonious labeling of corona graphs," *Journal of Applied Mathematics*, vol. 2014, 2014, doi: 10.1155/2014/627248.
- [5] P. Jeyanthi and S. Philo, "Odd Harmonious Labeling of Some New Families of Graphs," *Electronic Notes in Discrete Mathematics*, vol. 48, pp. 165–168, Jul. 2015, doi: 10.1016/j.endm.2015.05.024.
- [6] D. A. Pujiwati, I. Halikin, and K. Wijaya, "Odd harmonious labeling of two graphs containing star," in *AIP Conference Proceedings*, Feb. 2021, vol. 2326. doi: 10.1063/5.0039644.
- [7] G. A. Saputri, K. A. Sugeng, and D. Froncek, "The Odd Harmonious Labeling of Dumbbell and Generalized Prism Graphs," *AKCE International Journal of Graphs and Combinatorics*, vol. 10, no. 2, pp. 221–228, 2013, doi: 10.1080/09728600.2013.12088738.
- [8] S. S. Sarasvati, I. Halikin, and K. Wijaya, "Odd Harmonious Labeling of  $P_n \cong C_4$  and  $P_n \cong D_2(C_4)$ ," *Indonesian Journal of Combinatorics*, vol. 5, no. 2, p. 94, Dec. 2021, doi: 10.19184/ijc.2021.5.2.5.
- [9] K. A. Sugeng, S. Surip, and R. Rismayati, "On odd \ labeling of  $m$ -shadow of cycle, gear with pendant and Shuriken graphs," in *AIP Conference Proceedings*, Dec. 2019, vol. 2192. doi: 10.1063/1.5139141.
- [10] S. K. Vaidya and N. H. Shah, "Odd Harmonious Labeling of Some Graphs," 2012.
- [11] P. B. Sarasija and R. Binthiya, "Even Harmonious Graphs with Applications," *Internat. J. Comput. Sci. Infor. Security*, vol. 9, pp. 161–163, 2011.
- [12] J. A. Gallian, L. A. Schoenhard, and S. Arumugam, "Even Harmonious Graphs," *AKCE International Journal of Graphs and Combinatorics*, vol. 11, no. 1, pp. 27–49, 2014, doi: 10.1080/09728600.2014.12088761.
- [13] J. A. Gallian and D. Stewart, "Even harmonious labelings of disjoint unions with even sequential graphs," *J. Graph Labeling*, vol. 1, no. 1, pp. 1–10, 2015.
- [14] J. A. Gallian and D. Stewart, "Even harmonious labelings of disjoint graphs with a small component," *AKCE J. Graphs Combin*, vol. 12, no. 2–3, pp. 204–215, 2015.
- [15] J. A. Gallian and D. Stewart, "Properly even harmonious labelings of disconnected graphs," *AKCE International Journal of Graphs and Combinatorics*, vol. 12, no. 2–3, pp. 193–203, Nov. 2015, doi: 10.1016/j.akcej.2015.11.015.
- [16] G. Chartrand and P. Zhang, *A first course in graph theory*. Dover Publications, 2012.

