

ISSN (p) 1857-1727
ISSN (e) 2345-1688

Volume 13

No. 2, 2018

CHEMISTRY

JOURNAL OF MOLDOVA

General, Industrial and Ecological Chemistry

Editor-in-Chief: Gheorghe DUCA

NCAA national scientific journals ranking: Category A

Academy of Sciences of Moldova
Institute of Chemistry
Moldova State University

CHEMISTRY JOURNAL OF MOLDOVA

General, Industrial and Ecological Chemistry

is included/abstracted/indexed in:

Web of Science (Thomson Reuters-Emerging Source Citation Index)
SCOPUS (Elsevier international indexing database)
SJR (Scimago Journal and Country Rank)
CAS (Chemical Abstracts Service/ SciFinder)
VINITI (Russian Scientific and Technical Information Institute)
DOAJ (Directory Open Access Journals)
ROAD (Directory of Open Access Scholarly Resources)
SHERPA/ROMEO
OAJI (Open Academic Journals Index)
EVISA (European Virtual Institute for Speciation Analysis)
EIFL-OA Moldova (On-line Journals of Moldova)
BASE (Bielefeld Academic Search Engine)
Cite Factor (Directory Indexing of International Research Journals)
PubsHub Journals & Congresses
Academic Keys
Scholar Steer
Library of Moldova State University
Electronic Journals Library Database
Electronic Catalog of Academy of Sciences of Belarus
V. I. Vernadsky National Library of Ukraine
ZDB (Berlin State Library)
German National Library of Science and Technology University Library Hannover (Germany)
Temple University Libraries (USA)
The Royal Library Copenhagen/Copenhagen University Library (Denmark)
Digital library of "Dunarea de Jos" University (Romania)
Digital library of Ghent University (Belgium)
Digital library of Wroclaw University (Poland)
Digital library of Helsinki University (Finland)
Digital library of KTH Royal Institute of Technology (Sweden)
Digital library of Manchester University (UK)
Digital library of Cambridge University (UK)
Digital library of Sheffield University (UK)
Digital library of University of Texas at Austin (USA)
Digital library of Indiana University (USA)
Digital library of Simon Fraser University (Canada)
Digital library of University of Saskatchewan (Canada)
Digital library of Laval University (Canada)
Digital library of Auckland University (New Zealand)
Digital library of Queensland University of Technology (Australia)
Digital library of Victoria University (Australia)

Editorial production and secretariat:

Dr. Lilia Anghel
Dr. Larisa Postolachi
Dr. Olesea Cuzan-Munteanu
Iolanta Balan
Natalia Bolocan

Editorial office address:

Institute of Chemistry, 3, Academiei Str., Chisinau, MD-2028, Republic of Moldova
Tel: + 373 22 725490; Fax: +373 22 739954;
E-mail: chemjm.submission@gmail.com; chemjm@gmail.com; chemjm_correspondence@yahoo.com;
Web: <http://ejm.asm.md>

CHEMISTRY JOURNAL OF MOLDOVA

General, Industrial and Ecological Chemistry

Editor-in-Chief: Academician Gheorghe DUCA, Academy of Sciences of Moldova
Editors: Academician Tudor LUPASCU, Institute of Chemistry
Dr. Viorica GLADCHI, Moldova State University

Local Editorial Board:

Dr. Hab. A. ARICU
Institute of Chemistry

Dr. Hab. I. BULHAC
Institute of Chemistry

Dr. G. DRAGALINA
Moldova State University

Corr. member I. GERU
Institute of Chemistry

Dr. Hab. M. GONTA
Moldova State University

Acad. A. GULEA
Moldova State University

Dr. Hab. V. KULCITKI
Institute of Chemistry

Dr. Hab. F. MACAEV
Institute of Chemistry

Dr. R. NASTAS, scientific secretary
of the Editorial Board,
Institute of Chemistry

Dr. Hab. I. POVAR
Institute of Chemistry

Dr. Hab. V. RUSU
Institute of Chemistry

Dr. Hab. V. SARAGOV
A. Russo State University of Balti

Dr. Hab. R. STURZA
Technical University of Moldova

Dr. Hab. N. UNGUR
Institute of Chemistry

International Editorial Board:

Acad. S. ALDOSHIN
Russian Academy of Sciences

Acad. S. ANDRONATI
National Academy of Sciences of Ukraine

Acad. M. ANDRUH
University of Bucharest, Bucharest, Romania

Prof. V. ARION
University of Vienna, Austria

Prof. M. BAHADIR
Technical University of Braunschweig, Germany

Prof. A. BENNISTON
Newcastle University, UK

Acad. I. BERSUKER
University of Texas at Austin, USA

Prof. J. de BOER
Vrije University Amsterdam, The Netherlands

Prof. L. CIBOTARU
Katholieke Universiteit Leuven, Belgium

Prof. I. CRETESCU
Gh. Asachi Technical University of Iasi, Romania

Prof. A. GARABADZHIU
St-Petersburg State Institute of Technology, Russia

Acad. N. KARTEL
National Academy of Sciences of Ukraine

Prof. I. SANDU
Δ.I. Cuza University, Iasi, Romania

Acad. B. SIMIONESCU
Romanian Academy, Iasi, Romania

Corr. member B. TSUKERBLAT
Ben-Gurion University of the Negev, Israel

SCOPUS Metrics for 2017 are CiteScore 0.07, SNIP 0.164, SJR 0.105

Journal Quality Factor for the year 2017 is 1.20

Infobase Index Factor for the year 2017 is 3.5

Cosmos Impact Factor for the year 2017 is 5.224

ISSUE CONTENTS LIST WITH GRAPHICAL ABSTRACTS

NEWS AND EVENTS

7

**FIRST ANNOUNCEMENT - INTERNATIONAL CONFERENCE
“ACHIEVEMENTS AND PERSPECTIVES OF MODERN CHEMISTRY”**
October 2019, Chisinau, Republic of Moldova

REVIEW PAPER

NATURAL PRODUCT CHEMISTRY AND SYNTHESIS

8

CHEMICAL COMPOSITION AND BIOLOGICAL ACTIVITIES OF SOME *MARRUBIUM* SPECIES ESSENTIAL OIL: A REVIEW

Benalia Yabrir

This paper reviews information on essential oil of *Marrubium* species (except *M. vulgare*) described until now regarding extraction, chemical composition and biological activities. *Marrubium* essential oils, although quantitatively poor, are rich in chemical composition. This composition consists especially of sesquiterpenoids and a little amount of monoterpenes. It varies from one species to another, sometimes within same species. *Marrubium* essential oils exhibit antioxidant and antimicrobial activities. However, because the lack of literature concerning essential oil of these species, further studies are necessary, particularly regarding their activities.



RESEARCH PAPER


INDUSTRIAL CHEMISTRY

24

THE EFFECT OF RICE HUSK ON THE PHASE FORMATION AND CEMENT CLINKER PROPERTIES

Lev Chemyak, Liubov Melnyk, Natalia Dorogan

The features of the cement clinker containing rice husk are investigated. The dependence of the raw mixture composition on the set product burning characteristics has been analysed using the software “CLINKER”. The mixture compositions have been identified on the basis of the chalk–clay–rice husk system, with the introduction of man-made stock of 6.0–18.0 mass %. The features of the phase composition and the binder properties, by varying the rice husk content, mixture ratio and burning temperature were studied.



RESEARCH PAPER

INDUSTRIAL CHEMISTRY

32

METALLURGICAL COKEMAKING WITH THE IMPROVED PHYSICO-CHEMICAL PARAMETERS AT AVDEEVKA COKE PLANT

Oleg Zelenskii, Yuriy Vasil'ev, Alexey Sytnik, Natalia Desna,
Elena Spirina, Andrey Grigorenko

The article presents the results of studies regarding the improvement of the qualitative characteristics of blast furnace coke obtained from modified coal blend in industrial conditions of Avdeevka Coke Plant. Inorganic corundum powders are applied to modify the coal blend, namely electrocorundum (α -Al₂O₃) and carborundum (α -SiC). The introduction of a specific quantity (0.25 wt %) of non-clinkering additives allows the modification of the processes that occur when the coal blend is plastic, with consequent improvement in coke strength.



RESEARCH PAPER


INORGANIC AND COORDINATION CHEMISTRY

38

NMR STUDIES OF SOME ZN AND Cd COORDINATION COMPOUNDS BEARING 1,2-CYCLOHEXANEDIONEDIOXIME

Elena Gorincioi, Eduard Coropceanu

A series of homobi- and polynuclear zinc and cadmium coordination compounds supported by 1,2-cyclohexanedionedioxi- and bridging bidentate ligands: 4,4'-bipyridyl, 1,2-bis(4-pyridyl) ethane, 1,3-bis(4-pyridyl) propane and dipiridil sulfide were characterized by the experimental techniques of ¹H and ¹³C NMR spectroscopy. Individual NMR data of compounds are consistent with their assignment as complexes.



RESEARCH PAPER


INORGANIC AND COORDINATION CHEMISTRY

48

SYNTHESIS AND X-RAY CHARACTERISATION OF A NEW MIXED-VALENT TRINUCLEAR IRON CLUSTER

Olesea Cuzan-Munteanu, Silvia Melnic, Sergiu Shova

This paper reports on the synthesis of a new trinuclear homometalic mixed-valent iron carboxylate cluster with furan-2-carboxylic acid. The complex with the formula $[\text{Fe}_2^{\text{III}}\text{Fe}^{\text{II}}\text{O}(\text{C}_4\text{H}_3\text{O}_2\text{COO})_6(\text{H}_2\text{O})_3] \cdot 0.5\text{CH}_3\text{CN} \cdot 2.25\text{H}_2\text{O}$ was characterized by X-ray analysis which revealed that the compound crystallizes in the triclinic centrosymmetric group $P\bar{1}$, with the following unit cell parameters: $a = 10.2758(6)\text{ \AA}$, $b = 11.5991(9)\text{ \AA}$, $c = 19.7349(15)\text{ \AA}$, $\alpha = 105.060(7)^\circ$, $\beta = 94.216(6)^\circ$, $\gamma = 101.662(6)^\circ$.



RESEARCH PAPER


INORGANIC AND COORDINATION CHEMISTRY

56

SYNTHESIS AND CRYSTAL STRUCTURE OF BIS(CITRATO)GERMANATE AND STANNATE WITH TRIS(PHENANTHROLINE)NICKEL(II) CATION

Elena Martsinko, Inna Seifullina, Elena Chebanenko, Olha Pirozhok, Viktoriya Dyakonenko, Svitlana Shishkina

The new complexes $[\text{Ni}(\text{phen})_3][\text{Ge}(\text{HCit})_2] \cdot 2\text{H}_2\text{O}$ (1), $[\text{Ni}(\text{phen})_3][\text{Sn}(\text{HCit})_2] \cdot 3\text{H}_2\text{O}$ (2) (where phen is 1,10-phenanthroline, H_4Cit is citric acid) were obtained. The identity, composition, and thermal stability of the complexes were established by elemental analysis, thermogravimetry, and IR spectroscopy. According to the X-ray diffraction data, the bis(citrate)germanate / bis(citrate)stannate $[\text{Ge}/\text{Sn}(\text{HCit})_2]^{2-}$ carries out the role of the dianion while two halves of the nickel complexes with phenanthroline $[\text{Ni}(\text{Phen})_3]^{2+}$ are cations.



RESEARCH PAPER


NATURAL PRODUCT CHEMISTRY AND SYNTHESIS

63

CHEMICAL COMPOSITION AND ANTIMICROBIAL ACTIVITY OF THE LEVISTICUM OFFICINALE W.D.J. KOCH ESSENTIAL OIL

Alexandru Ciocarlan, Ion Dragalin, Aculina Aricu, Lucian Lupascu, Nina Ciocarlan, Violeta Popescu

The chemical composition of industrially obtained *Levisticum officinale* W.D.J. Koch (lovage) essential oil of Moldovan origin was analysed by means of chromatographic (GC-MS) and spectral (IR, ^1H and ^{13}C NMR) methods. The obtained results show that the main components of *L. officinale* essential oil are monoterpene hydrocarbons which make up to 53.50% of the total number of components. Antibacterial and antifungal activities of mentioned oil were evaluated *in vitro* on five strains of microorganisms. It was found that lovage volatile oil (*L. officinale*) exhibits high antibacterial and antifungal properties in the range of concentrations 0.015-0.030%.



RESEARCH PAPER


PHYSICAL CHEMISTRY AND CHEMICAL PHYSICS

69

ADSORPTION OF STRONTIUM IONS FROM AQUEOUS SOLUTIONS ON NUT SHELLS ACTIVATED CARBONS

Gheorghe Duca, Mihai Ciobanu, Tudor Lupascu, Igor Povar

The adsorption of strontium ions from aqueous solutions on nut shells activated carbons (samples CAN-7 and CAN-8) at different temperatures has been studied. The isotherm of adsorption of strontium ions from aqueous solutions on activated carbon CAN-7 has two inflection points at relatively small and high equilibrium concentrations. As the temperature increases, the adsorption values decrease, which indicates that the adsorption process is exothermic.



RESEARCH PAPER


PHYSICAL CHEMISTRY AND CHEMICAL PHYSICS

74

MULTICOMPONENT CRYSTALLIZATION OF KETOPROFEN-NICOTINAMIDE FOR IMPROVING THE SOLUBILITY AND DISSOLUTION RATE

Yudi Wicaksono, Dwi Setyawan, Siswandonno Siswandono

The study was aimed at increasing solubility and dissolution rate of ketoprofen by using multicomponent crystallization approach with nicotinamide as additional material. The multicomponent crystallization was carried out by solvent evaporation method using 2-propanol as solvent. The solubility and dissolution test showed that the ketoprofen-nicotinamide multicomponent crystal has solubility and dissolution rate significantly higher than the solubility and dissolution rate of the pure ketoprofen.



RESEARCH PAPER


PHYSICAL CHEMISTRY AND CHEMICAL PHYSICS

82

EFFECT OF THE NATURE OF SURFACTANT ON THE REACTIVITY OF C,N-DIPHENYLNITRONE TOWARDS ACRYLONITRILE IN DIFFERENT MICROEMULSION SYSTEMS

Kahina Hamza, Abdelkader Touati, Ahmed Ait-Yahia, Michel Baltas, Christiane André Barres, Saâd Moulay

The present work provides an insight into the effect of the nature of surfactant (cationic, anionic), a component of water- and oil-borne microemulsions, on the reaction rate of 1,3-dipolar cycloaddition of C,N-diphenylnitron with acrylonitrile. The electrostatically attractive character of the cationic surfactant, would bring the reactants closer to each other; hence, a rate enhancement would ensue, particularly within the water-rich zone. Besides the fact that acrylonitrile played a dual role, as a component of the microemulsion and a dipolarophile in the cycloaddition reaction, made the work-up advantageously sound. Additionally, the increase in reagents molar ratio was found to promote higher reactivity.



SHORT COMMUNICATION


PHYSICAL CHEMISTRY AND CHEMICAL PHYSICS

89

HYGROSCOPIC PROPERTIES OF ENOXIL-SILICA COMPOSITES

Oksana Stavinskaya, Iryna Laguta, Olga Kazakova, Pavlo Kuzema, Tudor Lupascu

Enoxil-silica composites with various Enoxil-to-silica ratios were prepared by mechanical mixing of the biologically active Enoxil and fumed silica powders. The hygroscopic properties of the composites were studied by the gravimetric method. It was found that hygroscopicity is reduced by 30÷40% when the Enoxil-to-silica mass ratio in the composites is 0.15÷0.35.



INSTRUCTIONS FOR AUTHORS

93

MULTICOMPONENT CRYSTALLIZATION OF KETOPROFEN-NICOTINAMIDE FOR IMPROVING THE SOLUBILITY AND DISSOLUTION RATE

Yudi Wicaksono ^{a*}, Dwi Setyawan ^b, Siswandono Siswandono ^b

^a Faculty of Pharmacy, University of Jember, 37, Jl. Kalimantan str., Jember 68121, Indonesia

^b Faculty of Pharmacy, Airlangga University, 4-6, Jl. Darmawangsa Dalam str., Surabaya 60286, Indonesia
e-mail: yudi.farmasi@unej.ac.id; phone: (+62 331) 32 17 36; fax: (+62 331) 32 17 36

Abstract. The purpose of this research was to improve the solubility and dissolution rate of ketoprofen by using the multicomponent crystallization approach with nicotinamide as coformer. Multicomponent crystallization of ketoprofen-nicotinamide with a 2:1 molar ratio was performed by solvent evaporation technique using 2-propanol as solvent. The characterization of the multicomponent crystal was performed using powder X-ray diffraction (PXRD), differential scanning calorimetry (DSC), Fourier transform infrared spectroscopy (FTIR) and scanning electron microscopy (SEM). The solubility and dissolution behaviour of the multicomponent crystal in distilled water were determined by the shake-flask and standard paddle method, respectively. The results of characterization by PXRD, DSC, FTIR and SEM have confirmed the formation of a new crystalline phase of ketoprofen-nicotinamide as the multicomponent crystal. The solubility of ketoprofen-nicotinamide multicomponent crystal was registered 1.3 times higher compared to pure ketoprofen. The multicomponent crystal dissolved about 64% within 60 minutes in comparison to the pure ketoprofen that showed a dissolution of only about 56% during the same period. The results of solubility and dissolution tests showed that the ketoprofen-nicotinamide multicomponent crystal is characterized by a solubility and dissolution rate significantly higher than those for the pure ketoprofen.

Keywords: ketoprofen, multicomponent crystallization, solvent evaporation, solubility, dissolution rate.

Received: 08 May 2018/ Revised final: 05 August 2018/ Accepted: 13 August 2018

Introduction

The solids of the active pharmaceutical ingredients (API) can be distinguished as crystalline and amorphous forms. API are commonly used in crystalline form, being thermodynamically more stable than the amorphous one [1,2]. A crystalline solid has three-dimensional long-range order and its physicochemical properties are strongly influenced by the molecular arrangement of components in the crystal lattice. Therefore, modification of the crystal lattice can be used to improve the physicochemical properties of crystalline API such as solubility, dissolution rate and hygroscopicity [3,4].

Currently, the modification of crystal lattice of API by the multicomponent crystallization approach is often used to improve the physicochemical properties of API [3-7]. The multicomponent crystallization approach is done by combining API with the other component called coformer [8]. In the multicomponent crystallization, the API and coformer are

crystallized together in the same crystal lattice through intermolecular interactions, such as hydrogen bonds and aromatic π - π stacking [8]. The formation of the new crystal lattice from API and coformer in the multicomponent crystal changes the crystal lattice and packing arrangement of the API, which induces the presence of new physicochemical properties of the API [8]. Therefore, one of the advantages of the multicomponent crystallisation approach is that it can be used to improve various physicochemical properties of API as needed by using the appropriate type of coformer.

Ketoprofen is a profen-type of non-steroidal anti-inflammatory drug, characterized by an asymmetric carbon center attached to a carboxylic acid, a methyl, and an aryl group. It is used as an analgesic and anti-pyretic agent to treat rheumatoid arthritis, osteoarthritis, dysmenorrhea, and to alleviate moderate pain [9]. It is a weak acid of propionic acid derivative, which is poorly soluble in water (0.253 mg/mL) [9]. It has been classified

as a Class II compound of the biopharmaceutical classification system, which means that its low water solubility is the limiting step for absorption and bioavailability [10]. Various methods are developed to improve the solubility of ketoprofen such as salt formation, micronization, formation of solid dispersions and re-crystallization [11-14]. However, the increased solubility of ketoprofen by using the above methods produces amorphous ketoprofen or ketoprofen with lower crystallinity, which is thermodynamically less stable.

The aim of the present study was to improve the solubility and dissolution rate of ketoprofen using the multicomponent crystallization approach. Nicotinamide was selected as coformer due to its acceptability for human consumption and it has effective functional groups to form intermolecular hydrogen bonds with various molecules [15,16]. The 2:1 ratio of ketoprofen-nicotinamide is used for the formation of the multicomponent crystal corresponding to the target stoichiometry and is commonly used for obtaining multicomponent crystals of profen-type drugs with nicotinamide as coformer [15,17,18]. The multicomponent crystallization of ketoprofen-nicotinamide was carried out by solvent evaporation technique using 2-propanol as solvent. The resulted multicomponent crystal was characterized by using PXRD, DSC, FTIR and SEM and was evaluated for solubility and dissolution behaviour by using the shake-flask and standard paddle method, respectively.

Experimental

Materials

Ketoprofen (purity $\geq 98.7\%$) was kindly assisted by Dexa Medica (Palembang, Indonesia). Nicotinamide (purity $\geq 99.6\%$) was purchased from Western Drugs Ltd (Udaipur, India) and 2-propanol (purity $\geq 99.8\%$) was purchased from Merck (Darmstadt, Germany).

Preparation of the multicomponent crystal

The multicomponent crystallization of ketoprofen and nicotinamide was carried out by solvent evaporation technique using 2-propanol as solvent. The mixture of ketoprofen-nicotinamide (2:1 molar ratio) was dissolved with 2-propanol in a beaker glass. The beaker glass was then covered with aluminium foil and the solution was stirred with a magnetic stirrer (Scilogex MS7-H550-Pro) for 30 minutes at room temperature. Shortly afterwards the aluminium foil was given small holes and settled at room temperature so the solvent evaporates slowly. The resulting solid was

then crushed using a mortar and pestle and passed through an 80 mesh sieve. The solid powder was stored in a desiccator until testing.

Characterization methods

The *PXRD diffractograms* were collected using a powder X-ray diffractometer (Philip Xpert) with $\text{CuK}\alpha$ radiation (1.54060 Å). The measurement was carried out at 5-50° in 2θ with a step size of 0.017° and a step time of 10 s/step. The condition of divergence slit and anti-scattering slit was 0.25° with diffraction experiment on the 10 mm sample size.

The *DSC analysis* was performed using a differential scanning calorimeter (Rigaku DSC 8230). Prior to measurement, the DSC instrument was calibrated for temperature and heat flow using high purity indium standard. The sample of approximately 2.0 mg was accurately weighed in a hermetically aluminium pan and then sealed. Measurements were performed from 30 to 200°C at a heating rate of 10°C/min under a dry nitrogen atmosphere (flow rate 50 mL/min).

The *FTIR spectra* were recorded using an attenuated total reflection Fourier transform infrared spectrometer (Bruker Alpha FTIR spectrometer). The measurements were performed with a resolution of 4 cm^{-1} in the range of 4000-400 cm^{-1} .

The *SEM analysis* was carried out using a scanning electron microscope (Hitachi TM-3000). The samples were firstly attached to a stub with double-sided carbon tape, and then coated with platinum using sputter coater ion (Hitachi E-1045) for 10 seconds. SEM analysis was performed at 15 kV accelerating voltage and 500x magnification.

Solubility and dissolution tests were determined using *UV-Vis analysis*.

Solubility and dissolution tests

Calibration curve. Ketoprofen stock solution (100 ppm) was prepared by dissolving 10 mg of pure ketoprofen in distilled water of 100 mL volumetric flask up to the mark. Ketoprofen standard solutions of concentration of 10, 20, 25, 40, 50 and 100 ppm were prepared by dilution of the stock solution with distilled water. The absorbance of the standard solutions was measured on a UV-Vis spectrophotometer (Hitachi U-2900) at 300 nm using 10 mm quartz cuvette. The measurements were carried out at 300 nm because nicotinamide had no absorbance at this wavelength and posed no interference to ketoprofen. The resulting calibration curve of ketoprofen in distilled water has the linear regression equation $y = 0.008x + 0.007$ with square correlation coefficient (R^2) of 0.9999 (Figure 1).




Figure 1. Calibration curve of ketoprofen in distilled water at $\lambda = 300$ nm.

Solubility test. The sample was tested for solubility in distilled water by the shake-flask method. An excess sample (about 50 mg) was fed into a 250-mL Erlenmeyer flask containing 50 mL of distilled water. Erlenmeyer flask was shaken using an incubator orbital (Stuart S1600) with a stirring speed of 150 rpm and a temperature of $37 \pm 0.5^\circ\text{C}$ for 12 hours. In the filtered solutions (0.45 μm filters) the ketoprofen concentration was determined from the calibration curve. Testing was conducted with three repetitions.

Dissolution test. The dissolution test was performed by paddle method using a dissolution apparatus (Logan UDT-804) in 900 mL of distilled water medium. The sample used was equivalent to 50 mg ketoprofen, while the stirring rate and the test temperature were 100 rpm and $37 \pm 0.5^\circ\text{C}$, respectively. At each time interval, 5 mL of the solution was withdrawn from the instrument and replaced by equal volume of distilled water. In the filtered solutions (0.45 μm filters) the ketoprofen concentration was determined from the calibration curve. Testing was repeated three times.

Statistical analysis

The statistical analysis of obtained data was conducted using one way analysis of variant (ANOVA) of SPSS version 16.0 for Windows. The test average values were considered significantly different at $p < 0.05$.

Results and discussion

Powder X-ray diffraction analysis

PXRD is a very important method in characterization of crystalline properties of solid-state samples. The formation of the new crystalline phase can be confirmed based on the unique fingerprint of the diffractogram [19].

The characteristic diffraction of multicomponent crystals is characterized by the appearance of new diffraction peaks and the absence of some diffraction peaks of individual components [20,21]. The overlay of powder X-ray diffractograms of ketoprofen, nicotinamide and ketoprofen-nicotinamide is shown in Figure 2. The diffractogram of ketoprofen has specific peaks of 2θ at 6.3, 14.3, 18.3, and 22.8° , while nicotinamide has 2θ specific peaks at 14.6, 14.8, 25.8, and 27.4° . The diffractogram of both materials indicated conformity with reports in previous studies [14,22,23].

The diffractogram of ketoprofen-nicotinamide exhibits new peaks at position 2θ 18.6° and 26.0° , which were absent in both the ketoprofen and the nicotinamide. The peaks at positions 11.2° and 27.2° present in the individual components were absent in the ketoprofen-nicotinamide. The powder X-ray diffractogram of ketoprofen-nicotinamide showed diffraction peaks different from those of the constituents. The diffraction peaks of ketoprofen-nicotinamide indicated the difference of its crystal lattice compared to the individual components. This indicated that the ketoprofen-nicotinamide was not a mechanical mixture, but had formed a multicomponent crystal with new solid crystalline phase [21,24].

DSC analysis

The DSC curves and the data of the melting temperature-enthalpy of fusion (ΔH) are shown in Figure 3 and Table 1, respectively. The DSC curve of ketoprofen showed a single endothermic peak at 96.2°C ($\Delta H = 94.8$ J/g), while the nicotinamide showed a single endothermic peak at 129.9°C ($\Delta H = 208.6$ J/g). These endothermic peaks were attributed to the melting points of the two ingredients. The melting points of the ketoprofen and nicotinamide are in good agreement with literature [22,25].

One indication of formation of a new crystalline phase is the emergence of different melting points with the initial components [26,27]. Ketoprofen-nicotinamide has a lower melting point than the melting point of both the individual components. The DSC curve of ketoprofen-nicotinamide has a melting endothermic peak at 72.4°C ($\Delta H = 27.6$ J/g) markedly different from the endothermic melting peak of ketoprofen and nicotinamide.




Figure 2. Powder X-ray diffractograms of pure ketoprofen (a), nicotinamide (b) and ketoprofen-nicotinamide (c).




Figure 3. DSC curves of ketoprofen (a), nicotinamide (b) and ketoprofen-nicotinamide (c).

Table 1

Melting temperature and enthalpy of fusion of ketoprofen, nicotinamide and ketoprofen-nicotinamide.		
Samples	Melting temperature ($^{\circ}\text{C}$)	Enthalpy of fusion (ΔH , J/g)
Ketoprofen	96.2	94.8
Nicotinamide	129.9	208.6
Ketoprofen-nicotinamide	72.4	27.6

These results indicated that ketoprofen and nicotinamide completely transformed into a multicomponent crystal as a new crystalline phase. In addition, ketoprofen-nicotinamide has a single endothermic peak indicated the absence of unbound component due to the mixture complete transformation into the crystalline phase of the multicomponent crystal [28].

FTIR analysis

FTIR spectroscopy is a technique commonly used to characterize intermolecular interactions in multicomponent crystals [27,29,30]. The obtained FTIR spectra of ketoprofen, nicotinamide and ketoprofen-nicotinamide are shown in Figure 4. The FTIR spectrum of ketoprofen has absorption peaks at 2979 and 2938 cm^{-1} (corresponding to a carboxylic -O-H stretching), 1695 cm^{-1} (-C=O acid stretching), 1654 cm^{-1} (-C=O ketone

stretching) and 1599 and 1451 cm^{-1} (aromatic ring -C=C- stretching). This is in agreement with previously reported research [23]. The FTIR spectra of nicotinamide has specific peaks at 3363 and 3166 cm^{-1} (-NH₂ stretching), 1676 cm^{-1} (-C=O amide stretching), 1397 cm^{-1} (-CN stretching), and 1340 cm^{-1} (ring stretching), which was also in accordance with previous research reports [31].

The spectra of ketoprofen-nicotinamide showed a shift of the absorption peaks compared to the spectra of individual components. The shift of the absorption peaks occurred in the -O-H and -C=C- stretching of ketoprofen (from 1451 cm^{-1} to 1445 cm^{-1}) and -NH₂ stretching (from 3166 cm^{-1} to 3177 cm^{-1}), -C=O amide stretching (from 1676 to 1695 cm^{-1}), -CN stretching (from 1397 to 1400 cm^{-1}) and pyridine ring stretching (from 1340 to 1371 cm^{-1}) of nicotinamide.




Figure 4. FTIR spectra of ketoprofen (a), nicotinamide (b) and ketoprofen-nicotinamide (c).

The shift of the absorption peaks of ketoprofen-nicotinamide indicated the formation of intermolecular interactions between ketoprofen and nicotinamide. Based on the shift of the absorption peaks, ketoprofen and nicotinamide formed a multicomponent crystal through intermolecular hydrogen bonds and π stacking interaction between the -O-H and the aromatic ring of ketoprofen with the -NH₂, -C=O, -CN, and the ring of nicotinamide [29,30,32-34].

Morphological analysis

The SEM images of ketoprofen, nicotinamide and the ketoprofen-nicotinamide multicomponent crystal are presented in Figure 5. The morphological changes of individual components in the ketoprofen-nicotinamide indicated the formation of a multicomponent crystal between ketoprofen and nicotinamide [7].




Figure 5. SEM images of ketoprofen (a), nicotinamide (b) and ketoprofen-nicotinamide (c).

Solubility assessment

The solubility of ketoprofen and ketoprofen-nicotinamide multicomponent crystal in distilled water was 181 ± 15 mg/mL and 241 ± 21 mg/mL, respectively. The ketoprofen-nicotinamide multicomponent crystal showed a significant increase in solubility ($p < 0.05$) of about 1.3 times higher compared to the solubility of ketoprofen. The increase of solubility of the multicomponent crystal is often associated

with a decrease of its melting point. A multicomponent crystal with a lower melting point tends to have higher solubility than the initial ingredient [35,36]. The DSC experiment showed that the ketoprofen-nicotinamide multicomponent crystal has a melting point at 72.46°C , which is lower by 24°C than ketoprofen. So it can be expected that the increased solubility of the ketoprofen-nicotinamide multicomponent crystal may be caused by the decrease of the melting point.

Dissolution behaviour

The dissolution behaviour of ketoprofen and ketoprofen-nicotinamide multicomponent crystal in distilled water was shown in Figure 6. The amount of ketoprofen released from the ketoprofen-nicotinamide multicomponent crystal within the first 30 minutes increased rapidly and subsequently formed a plateau profile indicating a saturation kinetics process [37]. The multicomponent crystal showed higher dissolution rate compared to ketoprofen with a cumulative amount released $64 \pm 5\%$ at 60 minutes. The dissolution profile in distilled water indicated that the amount of ketoprofen released from the ketoprofen-nicotinamide multicomponent crystal was higher than the pure ketoprofen (only about 56%).




Figure 6. Dissolution profile of ketoprofen (a) and ketoprofen-nicotinamide (b) in distilled water.

Conclusions

In the present study, the solubility and dissolution rate of ketoprofen was improved by the multicomponent crystallization approach using nicotinamide as a coformer in a 2:1 molar ratio. The formation of the multicomponent crystal of ketoprofen and nicotinamide was confirmed using PXRD, DSC, FTIR, and SEM methods.

The result of solubility and dissolution tests showed that the ketoprofen-nicotinamide

multicomponent crystal has higher solubility and dissolution rate than the pure ketoprofen. The multicomponent crystal showed a significant increasing in solubility ($p < 0.05$) of about 1.3 times higher compared to the solubility of ketoprofen (241±21 mg/mL and 181±15 mg/mL, respectively). The multicomponent crystal dissolution was about 64% within 60 minutes; however the pure ketoprofen showed a dissolution of only about 56% during the same period.

Acknowledgements

The authors would like to thank DRPM, the Ministry of research, technology, and higher education of the Republic of Indonesia for the research funding through doctoral dissertation research grant (Hibah Penelitian Disertasi Doktor) decree number 42/E/KPT/2017 under contract number 1867/UN25.3.1/LT/2017. The authors would also like to thank PT Dexa Medica for ketoprofen provision used in the study.

References

1. Dranca, I.; Povar, I.; Lupascu, T. Pharmaceutical amorphous organic materials characterization by using the differential scanning calorimetry and dynamic mechanical analysis. *Chemistry Journal of Moldova*, 2011, 6(2), pp. 91-95. DOI: [dx.doi.org/10.19261/cjm.2011.06\(2\).19](https://doi.org/10.19261/cjm.2011.06(2).19)
2. Vioglio, P.C.; Chierotti, M.R.; Gobetto, R. Pharmaceutical aspects of salt and cocrystal forms of APIs and characterization challenges. *Advanced Drug Delivery Reviews*, 2017, 117, pp. 86-110. DOI: <https://doi.org/10.1016/j.addr.2017.07.001>
3. Putra, O.D.; Furuishi, T.; Yonemochi, E.; Terada, K.; Uekusa, H. Drug-drug multicomponent crystals as an effective technique to overcome weaknesses in parent drugs. *Crystal Growth & Design*, 2016, 16(7), pp. 3577-3581. DOI: [10.1021/acs.cgd.6b00639](https://doi.org/10.1021/acs.cgd.6b00639)
4. Wicaksono, Y.; Wisudyaningsih, B.; Siswoyo, T.A. Enhancement of solubility and dissolution rate of atorvastatin calcium by co-crystallization. *Tropical Journal of Pharmaceutical Research*, 2017, 16(7), pp. 1497-1502. DOI: <https://dx.doi.org/10.4314/tjpr.v16i7.6>
5. Chadha, R.; Saini, A.; Jain, D.S.; Venugopalan, P. Preparation and solid-state characterization of three novel multicomponent solid forms of oxcabazepine: improvement in solubility through saccharin cocrystal. *Crystal Growth & Design*, 2012, 12(8), pp. 4211-4224. DOI: [10.1021/cg3007102](https://doi.org/10.1021/cg3007102)
6. Umeda, Y.; Fukami, T.; Furuishi, T.; Suzuki, T.; Tanjoh, K.; Tomono, K. Characterization of multicomponent crystal formed between indomethacin and lidocaine. *Drug Development and Industrial Pharmacy*, 2009, 35(7), pp. 843-851. DOI: <https://doi.org/10.1080/03639040802660489>
7. Ainurofiq, A.; Mauludin, R.; Mudhakhir, D.; Soewandhi, S.N. Synthesis, characterization, and stability study of desloratadine multicomponent crystal formation. *Research in Pharmaceutical Sciences*, 2018, 13(2), pp. 93-102. DOI: [http://doi.org/10.4103/1735-5362.223775](https://doi.org/10.4103/1735-5362.223775)
8. Boldyreva, E.V. Multicomponent organic crystals at high pressures. *Zeitschrift für Kristallographie*, 2014, 229(3), pp. 236-245. DOI: <https://doi.org/10.1515/zkri-2013-1699>
9. Shohin, I.F.; Kulinich, J.I.; Ramenskaya, G.V.; Abrahamsson, B.; Kopp, S.; Langguth, P.; Polli, J.E.; Shah, V.P.; Groot, D.W.; Barends, D.M.; Dressman, J.B. Biowaiver monographs for immediate-release solid oral dosage forms: Ketoprofen. *Journal of Pharmaceutical Sciences*, 2012, 101(10), pp. 3593-3603. DOI: <https://doi.org/10.1002/jps.23233>
10. Rençber, S.; Karavana S.Y.; Özyazici, M. Bioavailability file: Ketoprofen. *FABAD Journal of Pharmaceutical Sciences*, 2009, 34(4), pp. 203-216. <http://dergi.fabad.org.tr/tr/2009-volume-34-issue-4/>
11. Hildebrand, G.E.; Muller-Goymann, C.C. Ketoprofen sodium: Preparation and its formation of mixed crystals with ketoprofen. *Journal of Pharmaceutical Sciences*, 1997, 86(7), pp. 854-857. DOI: <https://doi.org/10.1021/js9602781>
12. Dixit, M.; Kulkarni, P.K.; Gowtham, V.; Shivakumar, H.G. Preparation and characterization of spray dried microparticle and chilled spray dried particle of ketoprofen by spray drying method. *Asian Journal of Pharmaceutical and Clinical Research*, 2011, 4(1), pp. 138-142. <https://innovareacademics.in/journal/ajpcr/>
13. Sherikar, A. Improvement of physicochemical characteristics and dissolution profile of poorly water soluble drug: ketoprofen by solid dispersion technique. *International Journal of Research in Pharmaceutical Sciences*, 2010, 1(4), pp. 450-453. <https://www.pharmascope.org/index.php/ijrps/article/view/558>
14. Dixit, M.; Kulkarni, P.K.; Vaghela, R.S. Effect of different crystallization techniques on the dissolution behavior of ketoprofen. *Tropical Journal of Pharmaceutical Research*, 2013, 12(3), pp. 317-322. DOI: [http://dx.doi.org/10.4314/tjpr.v12i3.7](https://dx.doi.org/10.4314/tjpr.v12i3.7)
15. Ando, S.; Kikuchi, J.; Fujimura, Y.; Ida, Y.; Higashi, K.; Moribe, K.; Yamamoto, K. Physicochemical characterization and structural evaluation of a specific 2:1 cocrystal of naproxen-nicotinamide. *Journal of Pharmaceutical Sciences*, 2012, 101(9), pp. 321-3221. DOI: <https://doi.org/10.1002/jps.23158>
16. Berry, D.J.; Seaton, C.C.; Clegg, W.; Harrington, R.W.; Colcs, S.J.; Horton, P.N.; Hursthouse, M.B.; Storey, R.; Jones, W.; Friscic, T.; Blagden, N. Applying hot-stage microscopy to co-crystal screening: A study of nicotinamide with seven active pharmaceutical ingredients. *Crystal Growth & Design*, 2008, 8(5), pp. 1697-1712. DOI: [10.1021/cg800035w](https://doi.org/10.1021/cg800035w)

17. Neurohr, C.; Marchivie, M.; Lecomte, S.; Cartigny, Y.; Couvrat, N.; Sanselme, M.; Subra-Paternault, P. Naproxen–nicotinamide cocrystals: Racemic and conglomerate structures generated by CO₂ antisolvent crystallization. *Crystal Growth & Design*, 2015, 15(9), pp. 4616-4626. DOI: [10.1021/acs.cgd.5b00876](https://doi.org/10.1021/acs.cgd.5b00876)
18. Blagden, N.; Coles, S.J.; Berry, D.J. Pharmaceutical co-crystals - Are we there yet? *CrystEngComm*, 2014, 16(26), pp. 5753-5761. DOI: [10.1039/C4CE00127C](https://doi.org/10.1039/C4CE00127C)
19. Padrela, L.; de Azevedo, E.G.; Velaga, S.P. Powder X-ray diffraction method for the quantification of cocrystals in the crystallization mixture. *Drug Development and Industrial Pharmacy*, 2012, 38(8), pp. 923-929. DOI: <https://doi.org/10.3109/03639045.2011.633263>
20. Hasa, D.; Miniussi, E.; Jones, W. Mechanochemical synthesis of multicomponent crystals: One liquid for one polymorph? A myth to dispel. *Crystal Growth & Design*, 2016, 16(8), pp. 4582-4588. DOI: <https://doi.org/10.1021/acs.cgd.6b00682>
21. Chadha, R.; Bhalla, Y.; Nandan, A.; Chadha, K.; Karan, M. Chrysin cocrystals: Characterization and evaluation. *Journal of Pharmaceutical and Biomedical Analysis*, 2017, 134, pp. 361-371. DOI: <https://doi.org/10.1016/j.jpba.2016.10.020>
22. Patil, S.P.; Modi, S.R.; Bansal, A.K. Generation of 1:1 carbamazepine: nicotinamide cocrystals by spray drying. *European Journal of Pharmaceutical Sciences*, 2014, 62, pp. 251-257. DOI: <https://doi.org/10.1016/j.ejps.2014.06.001>
23. Yadav, P.S.; Kumar, V.; Singh, U.P.; Bhat, H.R.; Mazumder, B. Physicochemical characterization and *in vitro* dissolution studies of solid dispersions of ketoprofen with PVP K30 and D-mannitol. *Saudi Pharmaceutical Journal*, 2013, 21(1), pp. 77-84. DOI: <https://doi.org/10.1016/j.jsps.2011.12.007>
24. Sangeetha, M.; Mathammal, R. Establishment of the structural and enhanced physicochemical properties of the cocrystal-2-benzyl amino pyridine with oxalic acid. *Journal of Molecular Structure*, 2017, 1143, pp. 192-203. DOI: <https://doi.org/10.1016/j.molstruc.2017.04.085>
25. Wicaksono, Y.; Seryawan, D.; Siswandono, S. Formation of ketoprofen-malonic acid cocrystal by solvent evaporation method. *Indonesian Journal of Chemistry*, 2017, 17(2), pp. 161-166. DOI: <https://doi.org/10.22146/ijc.24884>
26. Qiao, N.; Li, M.; Schlindwein, W.; Malek, N.; Davies, A.; Trappitt, G. Pharmaceutical cocrystals: An overview. *International Journal of Pharmaceutics*, 2011, 419(1-2), pp. 1-11. DOI: <https://doi.org/10.1016/j.ijpharm.2011.07.037>
27. Gao, Y.; Gao, J.; Liu, Z.; Kan, H.; Zu, H.; Sun, W.; Zhang, J.; Qian, S. Cofomer selection based on degradation pathway of drugs: A case study of adefovir dipivoxil–saccharin and adefovir dipivoxil–nicotinamide cocrystals. *International Journal of Pharmaceutics*, 2012, 438(1-2), pp. 327-335. DOI: <https://doi.org/10.1016/j.ijpharm.2012.09.027>
28. Manin, A.N.; Voronin, A.P.; Drozd, K.V.; Manin, N.G.; Bauer-Brandl, A.; Perlovich, G.L. Cocrystal screening of hydroxybenzamide derivatives with benzoic acid derivatives: A comparative study of thermal and solution-based methods. *European Journal of Pharmaceutical Sciences*, 2014, 65, pp. 56-64. DOI: <https://doi.org/10.1016/j.ejps.2014.09.003>
29. Chadha, R.; Bhandari, S.; Haneef, J.; Khullar, S.; Mandal, S. Cocrystals of telmisartan: characterization, structure elucidation, *in vivo* and toxicity studies. *CrystEngComm*, 2014, 16(36), pp. 8375-8389. DOI: [http://dx.doi.org/10.1039/c4ce00797b](https://doi.org/10.1039/c4ce00797b)
30. Hong, C.; Xie, Y.; Yao, Y.; Li, G.; Yuan, X.; Shen, H. A novel strategy for pharmaceutical cocrystal generation without knowledge of stoichiometric ratio: Myricetin cocrystals and a ternary phase diagram. *Pharmaceutical Research*, 2015, 32(1), pp. 47-60. DOI: <https://doi.org/10.1007/s11095-014-1443-y>
31. Rahman, Z.; Agarabi, C.; Zidan, A.S.; Khan, S.R.; Khan, M.A. Physico-mechanical and stability evaluation of carbamazepine cocrystal with nicotinamide. *AAPS PharmSciTech*, 2011, 12(2), pp. 693-704. DOI: <https://doi.org/10.1208/s12249-011-9603-4>
32. Masuda, T.; Yoshihashi, Y.; Yonemochi, E.; Fujii, K.; Uekusa, H.; Terada, K. Cocrystallization and amorphization induced by drug–excipient interaction improves the physical properties of acyclovir. *International Journal of Pharmaceutics*, 2012, 422(1-2), pp. 160-169. DOI: <https://doi.org/10.1016/j.ijpharm.2011.10.046>
33. Sowa, M.; Slepokura, K.; Matczak-Jon, E. Solid-state characterization and solubility of a genistein-caffeine cocrystal. *Journal of Molecular Structure*, 2014, 1076, pp. 80-88. DOI: <https://doi.org/10.1016/j.molstruc.2014.07.036>
34. Saha, S.; Rajput, L.; Joseph, S.; Mishra, M.K.; Ganguly, S.; Desiraju, G.R. IR spectroscopy as a probe for C–H···X hydrogen bonded supramolecular synthons. *CrystEngComm*, 2015, 17(6), pp. 1273-1290. DOI: [http://dx.doi.org/10.1039/C4CE02034K](https://doi.org/10.1039/C4CE02034K)
35. Parmar, V.K.; Shah, S.A. Hydrochloride salt co-crystals: preparation, characterization and physicochemical studies. *Pharmaceutical Development and Technology*, 2013, 18(2), pp. 443-453. DOI: <https://doi.org/10.3109/10837450.2012.696270>
36. Mulye, S.P.; Jamadar, S.A.; Karekar, P.S.; Pore, Y.V.; Dhawale, S.C. Improvement in physicochemical properties of ezetimibe using a crystal engineering technique. *Powder Technology*, 2012, 222, pp. 131-138. DOI: <https://doi.org/10.1016/j.powtec.2012.02.020>
37. Bolla, G.; Sanphui, P.; Nangia, A. Solubility advantage of tenoxicam phenolic cocrystals compared to salts. *Crystal Growth & Design*, 2013, 13(5), pp. 1988-2003. DOI: [http://dx.doi.org/10.1021/cg4000457](https://doi.org/10.1021/cg4000457)