

## Letter to the Editor

# Alleviating the impact of the geohelminthiases in the COVID-19 crisis

Sir,

Alleviating geohelminthiases impact during 2019 coronavirus disease (COVID-19) is complicated. Various efforts have been made to reduce them.<sup>1</sup> Chemoprevention such as annual or biennial mass preventive drugs (MPD) with albendazole or mebendazole as a single dose for at-risk populations, including preschool children, school-age children, adolescent girls, childbearing age women, and pregnant women, is the form of short-term action. Long-term actions are providing clean water, improving sanitation and toilets, banning on the usage of stool as fertilizer, and counseling on a clean and healthy lifestyle (CHL) accompanied by implementing community based total sanitation (CBTS). The challenges of them are inconstancy in observing the programs, chemoprevention coverage growth in at-risk groups, drug resistance manifestation, and weak diagnostic methods.<sup>2</sup>

Moreover, MPD for at-risk communities was also delayed due to inconsistencies in anthelmintic distribution.<sup>3</sup> If coinfection between the geohelminthiases parasites and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) increases complications and there is a shift in the comorbid age pattern to a younger age, then the burden of COVID-19 in endemic countries may be much worse than expected. However, the challenges of the COVID-19 crisis actually encourage these activities to be continued and integrated into ongoing COVID-19 prevention. However, every effort to increase the frequency and scale of MPD must be accompanied by programs to increase access to clean water, sanitation, and hygiene (WASH).<sup>4</sup>

The interruption and delay of the deworming control program due to COVID-19 temporarily impacted the progress of eliminating geohelminthiases as a public health problem (EHP) by 2030.<sup>5</sup> Implementation of chemoprevention should be restarted as soon as possible to minimize the time without chemoprevention as quickly as possible, even if it is before the following chemoprevention schedule under normal circumstances. COVID-19 crisis can be turned into opportunities to increase the likelihood of achieving targets by implementing appropriate alleviation strategies. A year's period of chemoprevention in the general population after discontinuation could accelerate the achievement of the target for reducing morbidity by 2030.<sup>6</sup>

Stimulating efforts to control geohelminthiases by consolidating integrated interventions and collaborations through the expansion of MPD, CHL, WASH and CBTS

in line with COVID-19 prevention efforts with alleviation innovations shifting from controlling morbidity to terminate the transmission of geohelminthiases.

This method accelerates progress towards the goal of geohelminthiases EHP in 2030. It is recommended to integrate the expansion of MPD coverage and the program's implementation to increase access to WASH which focuses on supporting the COVID-19 health protocol. Obstacles to alleviating geohelminthiases due to the COVID-19 crisis must be minimized, and the program should be restarted immediately.

**Pilar Menara Falah<sup>1\*</sup>, Yudha Nurdian<sup>2</sup>**

<sup>1</sup>Department of Basic Dental Sciences, Faculty of Dentistry, University of Jember, Jember, Indonesia

<sup>2</sup>Department of Basic Medical Sciences, Faculty of Medicine, University of Jember, Jember, Indonesia

**\*Correspondence to**

Pilar Menara Falah,

E-mail: [Pilarmenara18@gmail.com](mailto:Pilarmenara18@gmail.com)

## REFERENCES

1. Fauziyah S, Putri S, Salma Z, Wardhani H, Hakim F, Sucipto T, et al. How should Indonesia consider its neglected tropical diseases in the Covid-19 era? hopes and challenges (Review). *Biomed Rep*. 2021;14(6):53.
2. Toor J, Adams ER, Aliee M, Amoah B, Anderson RM, Ayabina D, et al. Predicted impact of COVID-19 on neglected tropical disease programs and the opportunity for innovation. *Clin Infect Dis*. 2021;72(8):1463-6.
3. Hollingsworth TD, Mwinzi P, Vasconcelo A, de Vlas SJ. 2021. Evaluating the potential impact of interruptions to neglected tropical disease programmes due to COVID-19. *Transact Royal Soc Tropic Med Hyg*. 2021;115(3):201-4.
4. Brooker SJ, Ziumbe K, Negussu N, Crowley S, Hammami M. Neglected tropical disease control in a world with COVID-19: An opportunity and a necessity for innovation. *Transact Royal Soc Tropic Med Hyg*. 2021;115(3):205-7.
5. Mationg MLS, Tallo VL, Williams GM, Gordon CA, Celements ACA, McManus DP, et al. The control of soil-transmitted helminthiases in the Philippines: The story continues. *Infect Dis Pover*. 2021;10(1):85.

6. Malizia V, Giardina F, Vegvari C, Bajaj S, McRae-McKee K. Modelling the impact of COVID-19-related control programme interruptions on progress towards the WHO 2030 target for soil-transmitted helminths. *Transact Royal Soc Tropic Med Hyg*. 2021;115(3):253-60.

**Cite this article as:** Falah PM, Nurdian Y. Alleviating the impact of the geohelminthiases in the COVID-19 crisis. *Int J Community Med Public Health* 2023;10:1285-6.

