



International
Pharmaceutical
Federation

FIP Education
(FIPEd)



Pharmacy Education
*an international
peer review journal*

Vol. 23 No. 2 (2023): IAI Special Edition

Published: 15-05-2023

PKP Publishing
Services

About the Journal

The *Pharmacy Education* journal is a peer-reviewed, open-access journal. The journal does not have article publication charges (APC) or subscription fees. It is published by the [International Pharmaceutical Federation \(FIP\)](#) and is aligned with a global mission of advancing education, practice, and science. The journal adheres to the principles of clinical and social pharmacy practice research as described in the [Granada Statements](#).

Pharmacy Education covers all aspects of pharmacy and pharmaceutical-related education, training, and workforce development. The journal's scope includes educational matters related to professional practice and pharmaceutical science.

The *journal* is listed in EBSCO and indexed in the [Emerging Sources Citation Index](#) (ESCI – Web of Science), EMBASE, and [SCOPUS](#).

Dr Marwan El Akel, Editor

Dr Sherly Meilianti, Production Editor

Professor Ian Bates, Editor-in-Chief

Editorial Team

Editor in Chief

Prof Ian Bates, FIP Education, United Kingdom

Journal Editor

Dr Marwan El Akel, Pharmacy Education

Senior Associate Editors

Dr Andreia Bruno-Tomé, Monash University, Australia

Assoc Prof Jennifer Marriott, Monash University, Australia

Production Editor

Dr Sherly Meilianti, Pharmacy Education

Associate Editors

Prof Joyce Addo-Atuah, Touro College of Pharmacy, USA

Prof Patricia Acuna-Johnson, University of Valparaiso, Chile

Dr Syed Imran Ahmed, University of Lincoln, United Kingdom

Prof Alba Mahmoud Albsoul-Younes, The University of Jordon, Jordan

Dr Ammar Almaaytah, Middle East University, Jordan

Dr Filipa Alves Da Costa, University of Lisbon, Portugal

Mr Chima Amadi, Pharmacists Council of Nigeria, Nigeria

Dr Mudassar Iqbal Arain, , University of Sindh, Pakistan.

Prof Lilian M. Azzopardi, University of Malta, Malta

Prof Rula Darwish, The University of Jordon, Jordan

Dr Ruth Edwards, Aston University, UK

Dr Divakar Goli, Acharya Institutes, India

Prof Yahdiana Harahap, University of Indonesia, Indonesia,

Prof Martin Henman, Trinity College Dublin, Ireland

Dr Shazia Jamshed, International Islamic University Malaysia, Malaysia

Dr Abby Kahaleh, Roosevelt University, USA

Prof Silvana Nair Leite, Federal University of Santa Catarina, Brazil

Dr Subhash Chandra Mandal, Directorate of Drugs Control, India

Mr Khalid Garba Mohammed, University of Milan, Italy

Dr Hana Morrissey, University of Wolverhampton, UK

Dr Christos Petrou, University of Nicosia, Cyprus

Dr Ukamaka Okafor, Pharmacists Council of Nigeria, Nigeria

Dr Carl Schneider, The University of Sydney, Australia

Prof Bruno Sepodes, University of Lisbon, Portugal

Dr James Scott, Western University of Health Sciences, USA

Prof M Chandra Sekar, University of Findlay, USA

Dr Rajani Shakya, Kathmandu University, Nepal

Dr Lixin Shu, Naval Medical University, China

Dr Judilynn Solidum, University of the Philippines, Philippines

Dr Kyle Wilby, University of Otago, New Zealand

Dr Sarah Willis, The University of Manchester, UK

Prof Shigeo Yamamura, Josai International University, Japan

Dr M. Nazli Sencan, Acibadem University, Istanbul, Turkey

Prof Abeer Al-Ghananeem, Sullivan University College of Pharmacy & Health Sciences
Kentucky, USA

Vol. 23 No. 2 (2023): IAI Special Edition

Published: 15-05-2023

Special Edition

- [Evaluation of COVID-19 vaccine storage at community health centres in Mataram city](#)

Baiq Lenysia Puspita Anjani, Cyntiya Rahmawati, Baiq Nurbaety, Baiq Leny Nopitasari , Anna Pradiningsih (Author)

p. 37-41

- [Impact of appropriate empirical antibiotics therapy on the clinical outcome of patients with urinary tract infections \(UTIs\)](#)

Fivy Kurniawati, Stephanus Manunggaling Ayun, Jesslyn Patricia (Author)

p. 184-189

- [Optimisation of lozenge formulation from Citronella \(*Cymbopogon nardus* \(L.\) Rendle\) extract with various binding materials using a simplex lattice design method](#)

Reynelda Juliani Sagala, Pretty Falena Atmanda Kambira, Untung Gunawan, Sondang, G.O Ambarita (Author)

p. 194-199

- [Profile of online drug purchasing in marketplace by students at a Faculty of Medicine and Health Sciences in Jakarta, Indonesia](#)

Fonny Cokro, Laurentine Belinda Arfenda, Hadiyanto (Author)

p. 144-148

- [Acute and sub-chronic toxicity study of green coffee extract \(*Coffea canephora* L.\) on liver function of wistar rats](#)

Fransiska Maria Christianty, Fifteen Aprilia Fajrin, Yearrika Rahayu Putra, Aida Nurmalita (Author)

p. 78-83

- [Translation, validation, and reliability of the Indonesian version Assessment of Quality of Life-4 Dimensions \(AQoL-4D\)](#)

Gusti N.V. Achmad, R.M, Husna, Y. Priyandani, E. Zairina (Author)

p. 9-13

- [Anti-inflammatory activity of *Eucheuma denticulatum* from Warambadi coast: In-vivo study model of carrageenan-induced paw oedema](#)

Erlia Anggrainy Sianipar, Adeline Jap (Author)

p. 216-222

- [Total flavonoid content and in vitro study on the sunscreen activity of extracts of leaves of *Elaeocarpus floribundus* blume](#)

Rahayu Utami, Raynaldi Syahputra, Rahma Dona, Haiyul Fadhli, Mustika Furi, Ihsan Ikhtiarudin (Author)

p. 118-121

- [Development of objective structured clinical examination-based assessment methods in drug information services lectures](#)

Yosef Wijoyo, Putu Dyana Christasani (Author)

p. 113-117

- [Formulation of a gambier catechin-loaded nanophytosome and the MTT assay on HeLa cell lines](#)

Henny Lucida , Suci Hasani , Meri Susanti , Friardi Ismed (Author)

p. 19-24

- [Modification of purple sweet potato starch \(*Ipomoea batatas* L. Poir\) with praelatination and acetylation methods as disintegrant of paracetamol tablets](#)

Budipratiwi Wisudyaningsih, Nina Wijiani, Vita Anggraeni (Author)

p. 207-211

- [Predicting factor analysis of gastrointestinal bleeding complication among hospitalised ischemic stroke patients](#)

Fivy Kurniawati, Erna Kristin, Rizaldy T. Pinzon, Sri Awalia Febriana (Author)

p. 139-143

- [The effect of *Aspergillus oryzae* and *Rhizopus aspergillus* fermentation on daidzein content in edamame \(glycine max\)](#)

Endah Puspitasari, Fanitika Imansari, Banun Kusumawardani, Ika Rahmawati Sutejo (Author)

p. 14-18

- [Analysis of total flavonoid content and antibacterial activity of the ethanolic extract of apu-apu herb \(*Pistia stratiotes*\) against *Salmonella typhimurium*](#)

Dewi Dianasari, Lilis Sapta Eka Lestari, Indah Yulia Ningsih (Author)

p. 212-215

- [The impact of COVID-19 on the management of medicines at a public health centre: A showcase of pharmacist resilience](#)

Firiyal Okta Safarah, Yuni Priyandani, Umi Athiyah, Abdul Rahem, Anila Impian Sukorini, Andi Hermansyah (Author)

p. 223-226

- [Correlation between clozapine use and metabolic syndrome in schizophrenic patients](#)

Woro Harjaningsih, Zullies Ikawati, Mustofa, Fitriani, Rina Silfiana Khoirunnisa, Nida Istivada (Author)

p. 103-112

- [Synthesis, molecular docking study, and in vivo biological evaluation of pyrazolopyridines derived from monocarbonyl curcumin analogues as potential anti-inflammatory agents](#)

Enda Mora, Hilwan Yuda Teruna, Neni Frimayanti, Ihsan Ikhtiarudin, Noval Herfindo, Adel Zamri (Author)

p. 200-206

- [Evaluation of drug information services in self-medication services with the patient simulation method at community pharmacies](#)

Devi Ristian Octavia , Sri Bintang Sahara Maha Putra Kusuma Negara, Primanitha Ria Utami (Author)

p. 92-97

- [In vitro antiplasmodial and toxicological activities of Vittaria anguste-elongata Hayata extracts](#)

Rudi Hendra, Rohimatul Khodijah, Hilwan Yuda Teruna (Author)

p. 190-193

- [Pharmacy student knowledge level regarding the beyond-use date](#)

Baiq Nurbaety, Cyntiya Rahmawati, Baiq Lenysia Puspita Anjani, Baiq Leny Nopitasari, Dwi Monika Ningrum (Author)

p. 60-64

- [The association between knowledge level and common cold self-medication behaviour among students of non-health faculty](#)

Devi Ristian Octavia , Pinasti Utami , Fitriana Yuliasuti (Author)

p. 149-155

- [Adverse drug reactions evaluation of antimicrobials in COVID-19 inpatients using Modified Trigger Tool and Naranjo Algorithm](#)

Larasati Arrum Kusumawardani, Nisa Maria, Yoga Amarta (Author)

p. 1-8

- [Cost effectiveness analysis of ceftriaxone and levofloxacin for therapy of urinary tract infection at Soebandi public hospital](#)

Ika Norcahyanti, Auralia Putri Pratama, Ema Rachmawati (Author)

p. 98-102

- [Antioxidant and toxicological activities of *Pyrossia lanceolata* \(L.\) Farw. extracts](#)

Ika Ria Indriani, Hilwan Yuda Teruna, Rudi Hendra (Author)

p. 174-178

- [Synthesis of 3'-methoxy flavonol and its derivatives as potential inhibitors for Dengue NS2B/NS3 and molecular insight into binding interaction](#)

Neni Frimayanti, Ihsan Ikhtiarudin, Rahma Dona, Istiazah Putri, Abdi Wira Septama (Author)

p. 231-243

- [Drug related problems in patients with pneumonia at Jasa Kartini Tasikmalaya city hospital](#)

Nur Rahayuningsih, Amalia Rahayu, Muharam Priatna (Author)

- [Hand sanitiser activity test of eucalyptus \(*Eucalyptus globulus*\) oil extract against *Bacillus subtilis* and *Enterococcus faecalis* bacteria](#)

Lukky Jayadi , Sandry Kesuma, Muhammad Hasan Wattiheluw (Author)

p. 179-183

- [Morphology and physicochemical properties of starch extracted from Indonesian ginger](#)

Indah Yulia Ningsih , Dewi Dianasari , Mochammad Amrun Hidayat (Author)

p. 47-52

- [Antibiotic use and antibiotic resistance profile of bacteria isolated from out-patients of Pakem primary healthcare, Yogyakarta](#)

Daru Estiningsih, Ika Puspitasari, Titik Nuryastuti, Endang Lukitaningsih (Author)

p.156-162

- [The effectiveness of using digital applications for diabetes mellitus with augmented reality models as learning media in pharmacy education](#)

Adin Hakim Kurniawan, Nanda Puspita, Yusmaniar, Febrain Rajendra (Author)

p. 53-59

- [Antibacterial properties of *Pyrrhosia longifolia* extracts](#)

Rohimatul Khodijah, Yuli Haryani , Hilwan Yuda Teruna, Rudi Hendra (Author)

p. 168-173

- [Public perceptions about telemedicine services for COVID-19 self-isolating patients](#)

Devi Ristian Octavia , Liza Pristianty, Andi Hermansyah (Author)

p. 227-230

- [Curcumin-mediated gene expression changes in Drosophila melanogaster](#)

Nurfadhilah Asfa, Ahmad Shayful Widiyanto, Muhammad Khadafi Anugrah Pratama, Reski Amalia Rosa, Ahmad Mu'arif, Risfah Yulianty, Firzan Nainu (Author)

p. 84-91

- [Antioxidant activity of Loranthus ferrugineus twigs extracts](#)

Desy Ariani, M. Almurdati, Rudi Hendra, Hilwan Yuda Teruna (Author)

p. 244-247

- [Fibrinolytic activity and molecular identification of PB-12 isolate from Papuma Coastal at Jember regency](#)

Evi Umayah Ulfa, Sattya Arimurti (Author)

p. 31-36

- [Validation and determination of Daidzein in Aspergillus oryzae using thin-layer chromatography-densitometry](#)

Endah Puspitasari, Sutatik, Banun Kusumawardani, Ika Rahmawati Sutejo (Author)

p. 42-46

- [Pineapple fruit extract \(Ananas comosus L. Merr\) as an antioxidant and anti-acne agent made with the nano-emulsion gel delivery system](#)

Lutfi Chabib, Arman Suryani, Loly Sintia Dewi, Herdwi Noviani, Widya Husna Puspa Maharani, Aina Anasta Indraswari (Author)

p.126-132

- [The correlation between knowledge, attitude and family support on compliance of outpatients with hypertension in a healthcare centre in Indonesia](#)

Liza Pristianty, Yuni Priyandani, Abdul Rahem (Author)

p. 25-30

- [Analysis of polypharmacy events and drug-drug interaction in COVID-19 therapy](#)

Anna Pradiningsih, Nurul Qiyaam, Baiq Leny Nopitasari (Author)

p. 133-138

- [The effect of e-booklet education on treatment behaviour of tuberculosis patients at Denpasar City health centre](#)

Ida Ayu Manik Partha Sutema, Ni Putu Aryati Suryaningsih, Gde Palguna Reganata, IGAR Widowati (Author)

p. 163-167

- [Formulation and in vitro study of nanoparticles loaded Anredera cordifolia leaf extract as anti-acne](#)

Lina Winarti, Adelia Amanda Safitri, Syafira Az-Zahro, Lusia Oktora RKS (Author)

p. 65-70

- [Antioxidant activities of medang lendir \(Litsea glutinosa\) stem bark](#)

Nabilah, Almurdani M. , Rudi Hendra, Hilwan Yuda Teruna (Author)

p. 122-125

Pharmacy Education

COUNTRY

Netherlands

Universities and research institutions in Netherlands

Media Ranking in Netherlands

SUBJECT AREA AND CATEGORY

Health Professions
Pharmacy

Pharmacology, Toxicology and
Pharmaceutics
Pharmaceutical Science

Social Sciences
Education

PUBLISHER

International Pharmaceutical
Federation

H-INDEX

19

PUBLICATION TYPE

Journals

ISSN

14772701, 15602214

COVERAGE

1973-1978, 2002-2022

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

pej@fp.org



IAI SPECIAL EDITION

RESEARCH ARTICLE

Validation and determination of Daidzein in *Aspergillus oryzae* using thin-layer chromatography-densitometry

Endah Puspitasari¹, Sutatik¹, Banun Kusumawardani², Ika Rahmawati Sutejo³¹ Pharmaceutical Biology Department, Faculty of Pharmacy, University of Jember, Jember, Indonesia² Biomedical Department, Faculty of Dentistry, University of Jember, Jember, Indonesia³ Biochemistry Department, Faculty of Medicine, University of Jember, Jember, Indonesia**Keywords**

Aspergillus oryzae
Daidzein content determination
TLC densitometry
Validation

Correspondence

Endah Puspitasari
Faculty of Pharmacy
University of Jember
Jember
Indonesia
e.puspitasari@unej.ac.id

Abstract

Background: Daidzin (glycosidic isoflavone) bioconversion into daidzein (aglyconic isoflavone) increases its bioactivity. This bioconversion can be carried out using fermentation, using a fungus that has been widely used for making tempeh in Indonesia, *Aspergillus oryzae*. **Objective:** This study aimed to validate the method of analysis used and to determine the daidzein content in *A. oryzae*-fermented edamame using TLC densitometry. **Method:** Daidzein content determination was carried out on non-fermented (D0) and fermented edamame on days one to four (D1-D4) using the TLC densitometry method. The validation method of analysis was done previously using linearity, the limit of detection (LOD), the limit of quantification (LOQ), selectivity, specificity, precision, and accuracy parameters. **Result:** The TLC densitometry method used met all the validation method of analysis parameters, except for the selectivity. The daidzein content in edamame from D0 to D4 were 0.0433 ± 0.00221 ; 0.0336 ± 0.00698 ; 0.0276 ± 0.00039 ; 0.0219 ± 0.00041 ; 0.1002 ± 0.00924 % w/w, respectively. **Conclusion:** The highest daidzein level was obtained in day four of *A. oryzae*-fermented edamame.

Introduction

The life expectancy of Indonesian women is increasing, from 73.33 years in 2010 to 73.55 years in 2016, resulting in an increase in the number of menopausal women (Badan Pusat Statistik, 2021). Menopausal women often experience uncomfortable symptoms i.e., hot flushes, insomnia, and vaginal dryness (Bruce & Rymer, 2009). They are seeking ways to decrease those symptoms, 76.1% are undergoing therapies including taking dietary soy and isoflavone supplements with 22.9% proportion. Among women using therapies, 89-100% found it beneficial (Newton *et al.*, 2002).

Soybean contains several natural isoflavones, and so does edamame, the vegetable soybean (Mentreddy *et al.*, 2002). Soybean isoflavones are found mostly in the glycosidic form. Glycosidic isoflavone is more bioactive

than glycosidic ones. Fermentation can remove the glycosidic group of glycosidic isoflavone resulting in a glycosidic isoflavone (Barnes, 2010).

This research was conducted to determine the daidzein (the major glycosidic isoflavone in soybeans) content in *Aspergillus oryzae*-fermented edamame using the thin-layer chromatography (TLC) method. *A. oryzae* was chosen because it is the common fungus used to make fermented soybean products, including tempeh and soy sauce (Yoneya, 2003). The validation method of analysis was done prior to the analysis since the determination of daidzein content in *A. oryzae*-fermented edamame has never been done before.

Method

Material

The SPMI edamame variety used in this study was obtained from PT. Mitra Tani Dua Tujuh and Jember. It was cultivated in Jember and harvested at 63rd-68th days. Other materials used were daidzein standard (Sigma), potato dextrose agar (BD Difco), Tween 80, TLC plate silica gel 60 F254 (Merck), toluene (Smart Lab Indonesia), ethyl acetate (Smart Lab Indonesia), acetone (Smart Lab Indonesia), n-hexane (Smart Lab Indonesia), ethanol 70%, methanol (Smart Lab Indonesia), and aqua dest.

Fermentation using *A. oryzae*

The edamame was peeled, washed, and sterilised using an autoclave. The fermentation was done on the cooled edamame using 10 ml of 10⁶ *A. oryzae*'s spore/ml per 500 g edamame, then incubated for one to four days (D1-D4) at 30°C 95% RH until solid-state fermentation was obtained (Lee, Hung, & Chou, 2008).

Extraction

The non-fermented (D0) and fermented edamame (D1-D4) were dried at 60°C for 30 hours in the oven, pulverised, and sifted. The defatting was done using 1:5 of n-hexane for three hours in the soxhlet apparatus. The defatted edamame was then air-dried overnight and extracted using ultrasonication and 1:6 of sample: 70% ethanol for one hour, followed by deposition using a centrifuge at 2,600 rpm for ten minutes (Miao, Qi, & Zhao, 2005). The extraction was repeated three times on the residue. The filtrate was collected and concentrated using a rotary evaporator until a thick extract was obtained (Luthria, Biswas, & Natarajan, 2007). The extract yield was calculated using the following formula:

$$\text{Yield} = \frac{\text{Extract weight}}{\text{Edamame weight}} \times 100\%$$

Analytical condition optimisation

The optimisation of the analytical condition was done prior to the validation method of analysis. The optimisation was done on solvent, stationary phase, eluent, λ_{max} , and sample concentration. The optimisation on the solvent was done by comparing the following eluent: n-hexane – ethyl acetate – formic acid (1:5:0.05), n-hexane – ethyl acetate – water (2:5:0.25), n-hexane – ethyl acetate (3:5), and n-hexane – ethyl acetate – acetate acid (2:5:0.15). λ_{max} was determined by wavelength scanning on the analyte at 200–400 nm. While the sample concentration optimisation was done by using three sample concentrations and comparing

them for the biggest N value (theoretical plate number) and smallest H value (height equivalent of a theoretical plate).

Validation method of analysis

The validation method of analysis on daidzein determination using TLC was done on linearity, LOD, LOQ, selectivity, specificity, precision, and accuracy. Those parameters were calculated using the validation method of the analysis programme. The selectivity was determined using the resolution (Rs) parameter, it was calculated based on the separation between the daidzein peak and two other peaks, i.e., genistein and glycitein on the sample chromatogram. The resolution was calculated by dividing the distance between two spots by the sum of the two spot radii. The parameters used to justify whether the method is valid or not are those from AOAC International (AOAC International, 2012).

Determination of daidzein content in non-fermented and *A. oryzae*-fermented edamame

The determination of daidzein content in non-fermented and *A. oryzae*-fermented edamame was done using the validated method of analysis. The data were analyzed using ANOVA followed by LSD post hoc ($\alpha = 0.01$).

Result

Mycelium growth was observed from D1 fermentation (Figure 1). It was originally whiteish (D1 and D2 fermentation), then turned brownish on D3 to D4 fermentation where a more compact mass was formed.




Figure 1: Morphological characteristic of non-fermented (D0) and *A. oryzae*-fermented (D1-D4) edamame

Extraction

The extract obtained from each sample is shown in Table I.

Table I: Extract obtained from edamame

Extract	Yield (% w/w)	Colour
D0	9.690	Light yellow
D1	17.185	Brownish yellow
D2	23.749	Dark brown
D3	12.570	Dark brown
D4	20.531	Dark brown

Analytical condition optimisation

The optimum condition of analysis is shown in Table II.

Table II: Optimum condition of analysis using TLC method

Analytical condition	Optimum condition of analysis
Solvent	Methanol p.a.
Stationary phase	Silica gel 60 F ₂₅₄
Eluent	n-hexane : ethyl acetate : acetic acid (2:5:0.15)
λ_{max}	273 nm
Sample concentration	80 mg/mL

The validation method of analysis

The standard curve obtained from the linearity test was $y=15.169x - 234.522$ (the sample concentrations used were six points ranging from ten to 100 µg/ml) with r value (correlation coefficient) of 0.999, $Vx0$ (the coefficient of variation of the regression function) = 3.068%, and $Xp = 55.675$. The LOD and LOQ value was 23.694 ng and 71.089 ng, respectively. The selectivity test resulted in the R_s value of daidzein-glycitein of 0.128, and in the R_s value of daidzein-genistein of 0.198. The purity and identity test for specificity resulted in both $r(s,m)$ and $r(m,e)$ values of 0.99. The precision parameter calculated from the six samples used was repeatability (4.467%) and intermediate precision (4.121%). While the recovery was used as an accuracy parameter (99.015%). It was calculated using the standard addition method.

Determination of daidzein content in *A. oryzae*-fermented edamame

The TLC profile of daidzein content determination in non-fermented and *A. oryzae*-fermented edamame is as shown in Figure 2. While the calculation on daidzein content is shown in Figure 3. The daidzein content in non-fermented and D1 of fermentation using *A. oryzae* was the same, but it was increasing significantly on day two to day four of fermentation and reached its peak on day four of fermentation.




(1) Daidzein 10 µg/ml; (2) Daidzein 30 µg/ml; (3) Daidzein 50 µg/ml; (4) Daidzein 70 µg/ml; (5) Daidzein 90 µg/ml; (6) Daidzein 100 µg/ml; (7) D0 sample replication 1; (8) D0 sample replication 2; (9) D0 sample replication

(1) D1 sample replication 1; (2) D1 sample replication 2; (3) D1 sample replication 3; (4) D2 sample replication 1; (5) D2 sample replication 2; (6) D2 sample replication 3; (7) Daidzein 10 µg/ml; (8) Daidzein 30 µg/ml; (9) Daidzein 50 µg/ml; (10) Daidzein 70 µg/ml; (11) Daidzein 90 µg/ml; (12) Daidzein 100 µg/ml; (13) D3 sample replication 1; (14) D3 sample replication 2; (15) D3 sample replication 3; (16) D4 sample replication 1; (17) D4 sample replication 2; (18) D4 sample replication 3

(a) Non-fermented edamame

(b) *A. oryzae*-fermented edamame

Figure 2: TLC profile of non-fermented (a) and *A. oryzae*-fermented (b) edamame under UV 254 nm



The data are shown as mean ± SD (n=3). Different annotation shows a significant difference (LSD, $p < 0.01$)

Figure 3: Daidzein content in non-fermented and *A. oryzae*-fermented edamame

Discussion

This research was conducted to mimic the process of making tempeh, one of the Indonesian traditional foods rich in isoflavones (Mani & Ming, 2017). However, we used edamame instead of soybean as the original source. The growth phase in edamame fermentation occurs at zero to 30 hours of fermentation, this phase is characterised by the growth of mycelia on the surface of the seeds. The next phase is the transition phase which occurs at 30-50 hours of fermentation, this phase is characterised by the presence of a specific odour such as the typical smell of tempeh (Dwinanto, 2011).

The extracts obtained from non-fermented and *A. oryzae*-fermented edamame are shown in Table I. The highest yield was obtained on the second day of fermentation (D2), which was 23.749 % w/w. The large yield indicates the quantity of compound in the extract but is not directly correlated to its activity. Extract yield only describes the total number of compounds taken,

but does not represent the concentration of one particular compound (Nuri *et al.*, 2020).

The validation method of analysis fulfilled all the criteria from AOAC International (2012), except for the specificity. To be considered specific, the Rs value should be > 1.5 (AOAC International, 2012). There is no literature pinpointing the Rs value of daidzein separation. The resolution value that did not meet the acceptance range may be due to the close proximity of the Log P values of the three components i.e., daidzein 2.5, genistein 2.7, and glycitein 2.4, making the separation of these three components difficult (National Center for Biotechnology Information, 2004a; National Center for Biotechnology Information, 2004b; National Center for Biotechnology Information, 2005).

Interestingly, the fermentation using *A. oryzae* on edamame fermentation made the daidzein content decrease on the first three days of fermentation, then drastically increased on the fourth day of fermentation using *A. oryzae* (Figure 3). The daidzein content then significantly increased on day four, with the value of

0.1002 ± 0.0092 % w/w. These findings are contradictive with previous research stating that fermentation would increase aglyconic isoflavone over time (Chun *et al.*, 2007; Chun, Kim, & Kim, 2008). Further research needs to be done to determine the exact mechanism of these phenomena.

Conclusion

The method of daidzein analysis fulfilled all the validation parameters, except for the selectivity. The highest daidzein content was that on the fourth day of fermentation

Acknowledgement

This article was presented orally at the 2022 Annual Scientific Conference of the Indonesian Pharmacist Association "Expanding The Role of Pharmacists to Strengthen National Resilience in a Dynamic Health Care and Health Development".

Source of funding

This research was funded by The Ministry of Research, Technology, and Higher education through Penelitian Produk Terapan scheme No. 0476/UN25.3.1/LT/ 2017.

Reference

AOAC International. (2012). *Official Methods of Analysis of AOAC International* (19th ed.). Washington DC: Association of Official Analytical Chemists

Badan Pusat Statistik. (2021). The Gender Development Index (GDI) 2019-2021. Retrieved August 6, 2022, from <https://www.bps.go.id/indicator/40/463/1/indeks-pembangunan-gender-ippg-.html>

Barnes, S. (2010). The biochemistry, chemistry and physiology of the isoflavones in soybeans and their food products. *Lymphatic research and biology*, *8*(1), 89–98

Barnes, S., Prasain, J., D'Alessandro, T., Arabshahi, A., Botting, N., Lila, M. A., Jackson, G., *et al.* (2011). The metabolism and analysis of isoflavones and other dietary polyphenols in foods and biological systems. *Food & function*, *2*(5), 235–244

Bruce, D., & Rymer, J. (2009). Symptoms of the menopause. *Best practice & research Clinical obstetrics & gynaecology*, *23*(1), 25–32

Chun, J., Kim, G. M., Lee, K. W., Choi, I. D., Kwon, G., Park, J., Jeong, S., *et al.* (2007). Conversion of isoflavone glucosides

to aglycones in soymilk by fermentation with lactic acid bacteria. *Journal of Food Science*, *72*(2), M39–M44

Chun, J., Kim, J. S., & Kim, J. H. (2008). Enrichment of isoflavone aglycones in soymilk by fermentation with single and mixed cultures of *Streptococcus infantarius* 12 and *Weissella* sp. 4. *Food chemistry*, *109*(2), 278–284

Dwinanto, S. (2011). Tempe. *Wacana Didaktika*, *1*(6), 25–32

Lee, I.-H., Hung, Y.-H., & Chou, C.-C. (2008). Solid-state fermentation with fungi to enhance the antioxidative activity, total phenolic and anthocyanin contents of black bean. *International Journal of Food Microbiology*, *121*(2), 150–156

Luthria, D. L., Biswas, R., & Natarajan, S. (2007). Comparison of extraction solvents and techniques used for the assay of isoflavones from soybean. *Food Chemistry*, *105*(1), 325–333

Mani, V., & Ming, L. C. (2017). Tempeh and other fermented soybean products rich in isoflavones. *Fermented foods in health and disease prevention* (pp. 453–474)

Mentreddy, S. R., Mohamed, A. I., Joshee, N., & Yadav, A. K. (2002). Edamame: a nutritious vegetable crop. *Trends in new crops and new uses. Proceedings of the Fifth National Symposium, Atlanta, Georgia, USA, 10-13 November, 2001* (pp. 432–438)

Miao, H., Qi, T., & Zhao, H. (2005). Methods for extracting, separating, identifying and quantifying daidzein and genistein. *Chinese Journal of Applied and Environmental Biology*, *11*(3), 293–295

National Center for Biotechnology Information. (2004a). Daidzein. *PubChem Compound Summary*

National Center for Biotechnology Information. (2004b). Genistein. *PubChem Compound Summary*. Retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/Genistein>

National Center for Biotechnology Information. (2005). Glycitein. *PubChem Compound Summary*. Retrieved from <https://pubchem.ncbi.nlm.nih.gov/compound/Glycitein>

Newton, K. M., Buist, D. S. M., Keenan, N. L., Anderson, L. A., & LaCroix, A. Z. (2002). Use of alternative therapies for menopause symptoms: results of a population-based survey. *Obstetrics & Gynecology*, *100*(1), 18–25

Nuri, N., Puspitasari, E., Hidayat, M. A., Ningsih, I. Y., Triatmoko, B., & Dianasari, D. (2020). Pengaruh Metode Ekstraksi terhadap Kadar Fenol dan Flavonoid Total, Aktivitas Antioksidan serta Antilipase Daun Jati Belanda (*Guazuma ulmifolia*). *Jurnal Sains Farmasi & Klinis*, *7*(2), 143–150

Yoneya, T. (2003). Fermented Soy Products: Tempeh, Nattos, Miso, and Soy Sauce. In Y. H. Hui, S. Ghazala, D. M. Graham, K. D. Murell, & W.-K. Nip (Eds.), *Handbook of Vegetable Preservation and Processing* (1st ed., pp. 251–272)