



## Protective effect of *Moringa oleifera* leaves extract on cardiac fibrosis of streptozotocin-induced diabetic rats

Zadia Maheswari Salsabila<sup>1</sup>, Suryono<sup>2\*</sup>, Pulong Wijang Pralampita<sup>3</sup>

<sup>1</sup>Faculty of Medicine, University of Jember, Jember, Indonesia, <sup>2</sup>Department of Cardiology and Cardiovascular Medicine, Faculty of Medicine, University of Jember/Dr. Soebandi Regional Hospital, Jember, Indonesia, <sup>3</sup>Department of Clinical Pathology, Faculty of Medicine, University of Jember, Jember, Indonesia

### ABSTRACT

Submitted: 2023-01-10  
Accepted : 2023-07-16

Diabetes mellitus (DM) is a metabolic disease characterized by chronic hyperglycemia that induces excessive reactive oxygen species (ROS) production and causes oxidative stress. Diabetic cardiomyopathy is a diabetic complication characterized by structural and functional changes of the myocardium. Fibrosis is one of the pathological features of diabetic cardiomyopathy. *Moringa oleifera* leaves have been reported to possess antidiabetic and antioxidant activities which could prevent diabetic complications such as cardiomyopathy. A previous study reported that *M. oleifera* leaves extract have protective effects to the kidneys and liver of rats exposed to oxidative stress. This study aimed to investigate the protective effect of the *M. oleifera* leaves extract on cardiac fibrosis of rats induced by streptozotocin (STZ). This was an experimental study using a posttest-only control group design. Thirty-three male Wistar rats were randomly divided into three groups i.e. normal control group (Group 1) were administered normal saline, diabetic control group (Group 2) were administered normal saline, and diabetic treatment group (Group 3) were administered *M. oleifera* leaves extract. Diabetes induction of rats was conducted by intraperitoneally injection of STZ at dose of 45 mg/kg BW. The *M. oleifera* leaves extract at a dose of 1000 mg/kg BW was administered orally one time a day for 28 days. Statistical analysis was performed using the Kruskal-Wallis test followed by Mann Whitney. A significant difference in cardiac fibrosis occurrence between three groups was observed ( $p < 0.05$ ). No cardiac fibrosis was observed in normal control group, In figure 3 we explained that fibrosis was observed in 8 rats of diabetic control group. Only 2 rats in the treatment group (G3) had cardiac fibrosis. In conclusion, *M. oleifera* leaves extract can inhibit cardiac fibrosis in STZ-induced diabetic rats.

### ABSTRAK

Diabetes melitus (DM) merupakan penyakit metabolik yang ditandai dengan hiperglikemia kronis yang menyebabkan produksi spesies oksigen reaktif (ROS) berlebihan dan menyebabkan stres oksidatif. Kardiomiopati diabetik adalah komplikasi diabetes yang ditandai dengan perubahan struktural dan fungsional miokardium. Fibrosis adalah salah satu gambaran patologis kardiomiopati diabetik. Daun kelor (*M. oleifera*) dilaporkan memiliki aktivitas antidiabetik dan antioksidan yang dapat mencegah komplikasi diabetes seperti kardiomiopati. Penelitian sebelumnya melaporkan bahwa ekstrak daun kelor memiliki efek protektif terhadap ginjal dan hati tikus yang terpapar stres oksidatif. Penelitian ini bertujuan untuk mengetahui efek protektif ekstrak daun kelor terhadap fibrosis jantung tikus yang diinduksi streptozotocin (STZ). Penelitian ini merupakan penelitian eksperimental dengan menggunakan rancangan *posttest-only control group*. Tiga puluh tiga ekor tikus Wistar jantan dibagi secara acak menjadi tiga kelompok yaitu kelompok kontrol normal (K1) yang disuntik dan diberikan larutan garam normal, kelompok kontrol diabetes (K2) diberikan larutan garam normal, dan kelompok perlakuan diabetes (Kelompok 3) diberikan ekstrak daun kelor. Induksi diabetes pada tikus dilakukan dengan cara penyuntikan STZ secara intraperitoneal dengan dosis 45 mg/kg BB. Ekstrak daun kelor dosis 1000 mg/kg BB diberikan secara oral satu kali sehari selama 28 hari. Analisis statistik dilakukan dengan menggunakan uji Kruskal-Wallis yang diikuti oleh Mann Whitney. Terdapat perbedaan yang signifikan dalam kejadian fibrosis jantung antara tiga kelompok yang diamati ( $p < 0,05$ ). Tidak ada fibrosis jantung yang diamati pada kelompok kontrol normal (K1), sedangkan fibrosis jantung diamati pada semua tikus (9 tikus) dari kelompok kontrol diabetes (K2). Hanya 2 tikus pada kelompok perlakuan (K3) yang mengalami fibrosis jantung. Kesimpulannya, ekstrak daun kelor dapat menghambat fibrosis jantung pada tikus diabetes yang diinduksi STZ.

### Keywords:

diabetes;  
oxidative stress;  
cardiac fibrosis;  
antioxidant;  
*Moringa oleifera*

## INTRODUCTION

Diabetes mellitus (DM) is a chronic metabolic disease characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both.<sup>1</sup> According to International Diabetes Foundation (IDF) in 2019, DM in Indonesia ranked seventh in the world with 10.7 million cases.<sup>2</sup> Uncontrolled DM can cause complications affecting the heart, kidneys, nerves and blood vessels.<sup>3</sup> Diabetic cardiomyopathy is a diabetic complication characterized by structural and functional changes of the myocardium, one of them is cardiac fibrosis.<sup>4</sup> The cardiac fibrosis can increase the risk of ventricular stiffness, which impairs cardiac contractility.<sup>5</sup>

According to a cohort study conducted over 5.5 years in 1.9 million patients with type 2 diabetes mellitus (T2DM), heart failure is the second most common manifestation of cardiovascular disease after peripheral arterial disease.<sup>6</sup> Another study reported that 32.3% of patients with T2DM experienced heart failure in less than 5 yr, and 67.7% of patients after 5 yr or more.<sup>7</sup> The cardiomyopathy also cause impairment of cardiac function which characterized by diastolic dysfunction. It was reported that the prevalence of diastolic dysfunction in patients with T2DM is between 40-60%.<sup>8</sup>

Hyperglycemia induces excessive production of reactive oxygen species (ROS) which further caused oxidative stress.<sup>9</sup> Oxidative stress has an important role in the pathogenesis of cardiac fibrosis through activation of the transforming growth factor (TGF)- $\beta$  leads to the activation of the suppressor of mothers against decapentaplegic (SMAD) signaling pathway included SMAD-2/3.<sup>10</sup> In the nucleus, the SMAD-2/3 complexes regulate the transcription of target profibrotic genes encoding proteins involved in extracellular matrix production, including collagen.<sup>11</sup>

Streptozotocin (STZ) is a diabetogenic agent that widely used in experimental studies.<sup>12</sup> Histopathological evaluation of the cardiac tissue of diabetic animal induced by STZ showed that diabetes causes structural changes in myocardium. These structural changes are caused by the excessive production of ROS in the myocardium due to hyperglycemic conditions.<sup>13</sup> A previous study also reported that diabetic rats induced by STZ show an increase in the collagen deposition.<sup>14</sup>

*Moringa oleifera* leaves have been reported to possess antidiabetic and antioxidant activities, which are helpful for the treatment of diabetes and its complications.<sup>15</sup> Quercetin in *M. oleifera* leaves extract was reported as one of the active constituents that is responsible for its high antioxidant activity.<sup>16</sup> *Moringa oleifera* exhibits cardioprotective effects in cardiac damage and vascular dysfunction due to its antioxidant activity.<sup>17</sup> In addition, the rich polyphenolic content of *M. oleifera* reduces the myocardial damage and decreases the oxidative stress.<sup>18</sup> Previous studies reported that administration of *M. oleifera* leaves extract at a dose of 1000 mg/kg BW exhibits protective effects to the kidneys and liver of rats exposed to oxidative stress.<sup>19,20</sup> The aim of this study was to investigate the protective effect of the *M. oleifera* leaves extract on cardiac fibrosis of rats induced by STZ.

## MATERIAL AND METHODS

### *M. oleifera* leaves extract preparations

The leaves of *M. oleifera* were obtained from a farmer in Wuluhan District, Jember Regency, and were authenticated at the Herbal Materia Medica Laboratory, Batu, East Java. The leaves were then washed and dried in oven at a temperature of 55-60°C. Once dried, the leaves were ground into a fine powder using a blender and

sieve. The dried and powdered leaves were extracted by maceration using 96% ethanol for 72 h. The mixture was stirred using an orbital shaker to facilitate the extraction process. After 72 h of maceration, the mixture was filtered using Whatman filter paper. The filtrate was then evaporated using water bath at a temperature of 70 °C. The dried extract obtained was then kept in a refrigerator until used.

### Animals and design

This was an experimental laboratory study using a posttest-only control group design. Thirty-three male Wistar rats with weighing 200-300 g at 8-12 wk of age were randomly divided into three groups i.e. normal control rats' group (Group 1) were injected and administered normal saline, diabetic control rats' group (Group 2) were administered normal saline, and diabetic treatment rats' group (Group 3) were administered *M. oleifera* leaves extract. The rats were housed in individual cages, which were cleaned every 2-3 d in a wk. Rats were given standard feed and water *ad libitum*. The *M. oleifera* leaves extract at dose of 1000 mg/BW was orally administered daily started from day 4 and continued for the next 28 d. The rats' body weight was measured five times during the study using a digital scale. This study has been approved by the Medical and Health Research Ethics Committee, Faculty of Medicine, University of Jember, with ref:1599/H25.1.11/KE/2022.

### Induction of diabetes by streptozotocin

Diabetes induction was performed by a single intraperitoneal injection of freshly STZ at a dose of 45 mg/kg BW by dissolved in 0.1 M of cold citrate buffer at pH 4.5 to overnight fasted male Wistar rats. After STZ induction, the rats were provided with 10% dextrose overnight to prevent hypoglycemic shock. Three days

after STZ induction, the fasting blood glucose levels were measured using EasyTouch glucometer. Rats with fasting blood glucose level (BGL) of greater than 200 mg/dL were considered as diabetic rats and included in this study.

### Histological examination

The histopathological examination of was conducted at the Biomedical Laboratory, Faculty of Dentistry, University of Jember. Rats were sacrificed by intraperitoneal injection of pentobarbital at dose of 150 mg/kg BW after 4 wk of treatment. The heart were cleaned with 0.9% NaCl solution to remove any remaining blood or dirt and were preserved in a jar containing a 10% buffer neutral formalin solution.<sup>21</sup> Masson's trichrome staining was used to identify collagen deposition in the cardiac tissue. Each preparation was examined under a microscope at magnification ranging from 100x to 400x and at five fields of view to evaluate the collagen deposition in cardiac tissue. Collagen deposition was stained blue, whereas the cardiac muscle was stained red.<sup>22</sup> The histopathological examination was carried out by two individuals using a single blind technique under the supervision of an anatomical pathologist.

### Statistical analysis

Statistical analysis was performed using SPSS version 23 software. Statistical differences were assessed using the Kruskal-Wallis test, followed by Mann Whitney Test, due to not normal data distribution. Differences between groups were considered significant if a p value of <0.05.

## RESULTS

### Streptozotocin induction

The BGL of normal control group

(G1), diabetic control group (G2) and treatment group (G3) are presented in TABLE 1. The mean BGL of G1 is  $108.91 \pm 9.101$  mg/dL (<200 mg/dL) which considered as normal rats. The mean BGL of G2 and G3 are  $437.82 \pm 38.126$  mg/dL;  $424.91 \pm 41.469$  mg/dL (>200 mg/dL) which considered as diabetic rats (TABLE 1).

**Body weight of rats after STZ induction**

The mean growth of rats body weight in all groups are presented in FIGURE 1. No significant difference in the body weight of rats in all groups in the week 0 ( $p > 0.05$ ). A significant difference in the body weight of rats between normal control group (G1) with diabetic control group (G2) and treatment group (G3) was observed ( $p < 0.05$ ). The growth of rats body weight was only observed in normal control group (G1), whereas in diabetic control group (G2) and treatment group

(G3), the loss of rats body weight during study in the week 1 to 4 was observed.

**Protective effect of *M. oleifera* leaves extract**

Among 11 rats induced by STZ, only 9 rats in each group were survive and eligible for histopathological examination. Histopathological features of cardiac tissues of rats in each group are presented in FIGURE 2 and the results of histopathological examinations are presented in FIGURE 3. Significantly different in cardiac fibrosis between the three groups was observed ( $p < 0.05$ ) No fibrosis was observed in normal control group (G1), In figure 3 we explained that fibrosis was observed in 8 rats of diabetic control group and in 2 rats of treatment group (G3).

TABLE 1. Rats fasting BGL (mg/dL) in all of groups after STZ induction

Group	n	Mean $\pm$ SEM	Min	Max
Normal control group (G1)	11	$108.91 \pm 9.101$	66	156
Diabetic control group (G2)	11	$437.82 \pm 38.126$	248	600
Treatment group (G3)	11	$424.91 \pm 41.469$	208	600





FIGURE 1. Mean body weight of rats



A. Normal control B. Diabetic control C. Treatment  
 FIGURE 2. Histopathological features of cardiac tissues of rats in each group after microscopic examination with 400x magnification. A. Normal control: no fibrosis was observed; B. Diabetic control: fibrosis was observed (yellow arrow); C. Treatment: no fibrosis was observed.




FIGURE 3. The results of histopathological examination of cardiac tissues of rats in each group.

## DISCUSSION

In this study STZ is used to induce diabetes of rats. Streptozotocin-induced diabetes model is widely used in various

studies related to diabetes in animal model. Streptozotocin is a naturally occurring chemical, a broad-spectrum antibiotic that is specifically toxic to the insulin-producing  $\beta$ -cells of the

pancreas lead to diabetes of the animal model.<sup>23,24</sup> Lew *et al.*<sup>25</sup> reported that the characteristics of STZ-induced diabetes rats include hyperglycemia, polyphagia, polyuria, polydipsia, and significant weight loss. The animal model is considered diabetes if the BGL greater than 200 mg/dL. This study showed that three days after STZ induction, the rats BGL of the diabetic control group (G2) and the treatment group (G3) range from 208 to 600 mg/dL (TABLE 1).

In addition, a significant weight loss was also observed in this study after STZ induction (FIGURE 1). The significant weight loss of the STZ-induced diabetes rats occurred due to an increased protein breakdown in muscle and tissue.<sup>26</sup> Diabetes leads to changes in protein metabolism.<sup>27</sup> The elevated protein catabolism for gluconeogenesis that cause weight loss occurs in diabetes.<sup>28</sup>

In figure 3 we explained that fibrosis was observed in 8 rats of diabetic control group and the part of rats in treatment group (G3) in this study (FIGURE 3). Cardiac fibrosis is an injury condition in the cardiac muscle that is characterized by excessive deposition of type 1 collagen.<sup>29</sup> Cardiac fibrosis in STZ-induced diabetic rats occurs due to the differentiation of cardiac fibroblast into myofibroblast.<sup>30</sup> Furthermore, the myofibroblasts secrete excessive amount of ECM such as collagen.<sup>31</sup> It was also reported that an increased expression of TGF- $\beta$ ,  $\alpha$ -SMA, and connective tissue growth factor (CTGF) that contributed in the development of cardiac fibrosis are observed in the hearts of STZ-induced diabetic rats.<sup>30,32,33</sup> The CTGF is a fibrogenic factor that promotes fibroblast proliferation and interstitial collagen deposition.<sup>34</sup>

Hyperglycemia in STZ-induced diabetic rats increased ROS production

which cause imbalance between ROS and antioxidant levels lead to oxidative stress.<sup>9,13,24,34</sup> The levels of MDA of diabetic rats increase lead to stimulate fibroblast proliferation and other fibrotic signaling pathways in cardiac tissue.<sup>31,34</sup> Oxidative stress triggers the activation of the TGF- $\beta$  pathway, which, in turn, activates the SMAD signaling pathway, specifically SMAD-2/3.<sup>10</sup> Within the nucleus, SMAD-2/3 complexes govern the transcription of target profibrotic genes responsible for encoding proteins involved in extracellular matrix production, such as collagen.<sup>11</sup>

This study showed that the administration of *M. oleifera* leaves extract inhibits the incidence of cardiac fibrosis in diabetic rats. A better histological feature of cardiac tissue of diabetic rats treated by *M. oleifera* leaves extract was observed compared to that untreated diabetic rats. *Moringa oleifera* leaves extract can ameliorate the histological feature of cardiac tissue of diabetic rats as almost its normal condition. Previous study reported that *M. oleifera* leaves extract has protective effect on cardiac antioxidant status and lipid peroxidation in STZ-induced diabetic rats.<sup>35</sup>

*Moringa oleifera* leaves has been reported to contain active antioxidant compounds such as vitamin C, vitamin E, and flavonoids.<sup>36-38</sup> Quercetin, a potent antioxidant flavonoid found in high concentration in *M. oleifera*, has therapeutic activities.<sup>39,18</sup> The quercetin is responsible for free radical scavenging activity. Moreover, it was also reported that the quercetin in *M. oleifera* leaves extract reduce the cardiac necrosis biomarkers levels and normalize the myocardium structure in both *in vitro* and *in vivo* studies.<sup>18</sup>

Previous studies reported that *M.*

*oleifera* leaves extract at a dose of 1000 mg/kg BW significantly showed positive effects to the rat organs exposed to oxidative stress.<sup>19,20,40</sup> It was also reported that *M. oleifera* leaves extract has a renoprotective effect on the glomerulus of diabetic rats. The *M. oleifera* leaves extract restored the glomerulus damage from a score of 4 to a score of 1 to 0 or return to its normal condition. Another study reported that *M. oleifera* leaves extract has a hepatoprotective effect against oxidative stress by restoring the structure of liver tissue close to normal.<sup>20</sup> *Moringa oleifera* leaves extract restored the structure of pancreatic islet cells of hypercholesterolemic rats model.<sup>40</sup>

## CONCLUSION

In conclusion, *M. oleifera* leaves extract can inhibit cardiac fibrosis in STZ-induced diabetic rats.

## ACKNOWLEDGEMENTS

We would like to express our gratitude to the staff at the Faculty of Medicine and Faculty of Dentistry University of Jember, for valuable assistances during the study.

## REFERENCES

- Petersmann A, Müller-Wieland D, Müller UA, Landgraf R, Nauck M, Freckmann G, *et al.* Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes* 2019; 127(S 01):S1-7.  
<https://doi.org/10.1055/a-1018-9078>
- Kementerian Kesehatan Republik Indonesia. Tetap produktif, cegah dan atasi diabetes mellitus. Pusat Data dan Informasi Kementerian Kesehatan Republik Indonesia. 2020.
- WHO. Diabetes [Internet]. 2020 [cited 2021 Jul 17].
- Sun L, Yu M, Zhou T, Zhang S, He G, Wang G, *et al.* Current advances in the study of diabetic cardiomyopathy: from clinicopathological features to molecular therapeutics. *Mol Med Rep* 2019; 20(3):2051-62.  
<https://doi.org/10.3892/mmr.2019.10473>
- Travers JG, Kamal FA, Robbins J, Yutzey KE, Blaxall BC. Cardiac fibrosis: the fibroblast awakens. *Circ Res* 2016; 118(6):1021-40.  
<https://doi.org/10.1161/CIRCRESAHA.115.306565>
- Low Wang CC, Hess CN, Hiatt WR, Goldfine AB. Clinical update: cardiovascular disease in diabetes mellitus: atherosclerotic cardiovascular disease and heart failure in type 2 diabetes mellitus - mechanisms, management, and clinical considerations. *Circulation* 2016; 133(24):2459-502.  
<https://doi.org/10.1161/CIRCULATIONAHA.116.022194>
- Paramita AAKY, Saraswati MR, Wiryawan N. Gambaran karakteristik gagal jantung pada pasien diabetes melitus di RSUP Sanglah Denpasar. *Jurnal Penyakit Dalam Udayana* 2021; 5(2):37-45.  
<https://doi.org/10.36216/jpd.v5i2.152>
- Almehmadi F, Joncas SX, Nevis I, Zahrani M, Bokhari M, Stirrat J, *et al.* Prevalence of myocardial fibrosis patterns in patients with systolic dysfunction: prognostic significance for the prediction of sudden cardiac arrest or appropriate implantable cardiac defibrillator therapy. *Circ Cardiovasc Imaging* 2014; 7(4):593-600.  
<https://doi.org/10.1161/CIRCIMAGING.113.001768>

9. Volpe CMO, Villar-Delfino PH, Dos Anjos PMF, Nogueira-Machado JA. Cellular death, reactive oxygen species (ROS) and diabetic complications review-Article. *Cell Death Dis* 2018; 9(2):119. <https://doi.org/10.1038/s41419-017-0135-z>
10. Kong P, Christia P, Frangogiannis NG. The pathogenesis of cardiac fibrosis. *Cell Mol Life Sci* 2014; 71(4):549-74. <https://doi.org/10.1007/s00018-013-1349-6>
11. Walton KL, Johnson KE, Harrison CA. Targeting TGF- $\beta$  mediated SMAD signaling for the prevention of fibrosis. *Front Pharmacol* 2017; 8:461. <https://doi.org/10.3389/fphar.2017.00461>
12. Ighodaro OM, Adeosun AM, Akinloye OA. Alloxan-induced diabetes, a common model for evaluating the glycemic-control potential of therapeutic compounds and plants extracts in experimental studies. *Medicina (Kaunas)* 2018; 53(6):365-74. <https://doi.org/10.1016/j.medic.2018.02.001>
13. Wang Y, Sun H, Zhang J, Xia Z, Chen W. Streptozotocin-induced diabetic cardiomyopathy in rats: ameliorative effect of piperine via Bcl2, Bax/Bcl2, and caspase-3 pathways. *Biosci Biotechnol Biochem* 2020; 84(12):2533-44. <https://doi.org/10.1080/09168451.2020.1815170>
14. Tsutsui H, Matsushima S, Kinugawa S, Ide T, Inoue N, Ohta Y, *et al.* Angiotensin II type 1 receptor blocker attenuates myocardial remodelling and preserves diastolic function in diabetic heart. *Hypertens Res* 2007; 30(5):439-49. <https://doi.org/10.1291/hyres.30.439>
15. Omodanisi EI, Aboua GY, Oguntibeju OO. Therapeutic potentials and pharmacological properties of moringa oleifera lam in the treatment of diabetes mellitus and related complications. *Trop J Pharm Res* 2017; 16(7):1737-46. <https://doi.org/10.4314/tjpr.v16i7.35>
16. Fachriyah E, Kusriani D, Haryanto IB, Wulandari SMB, Lestari WI, Sumariyah. Phytochemical test, determination of total fenol, total flavonoids and antioxidant activity of ethanol extract of moringa leaves (*Moringa oleifera* Lam). *Jurnal Kimia Sains dan Aplikasi* 2020; 23(8):290-4. <https://doi.org/10.14710/jksa.23.8.290-294>
17. Alia F, Putri M, Anggraeni N, Syamsunarno MRAA. The potency of *Moringa oleifera* Lam. as protective agent in cardiac damage and vascular dysfunction. *Front Pharmacol* 2022; 12:724439. <https://doi.org/10.3389/fphar.2021.724439>
18. Panda S. Butanolic fraction of *Moringa oleifera* Lam. (Moringaceae) attenuates isoprotrenol – induced cardiac necrosis and oxidative stress in rats: an EPR study. *EXCLI J* 2015; 14:64-74. <https://doi.org/10.17179/excli2014-43>
19. Gianosa G, Santosa A, Suryono S. Renoprotective activity of *Moringa oleifera* Lam toward kidney injury rats induced by streptozotocin. *NLJ* 2020; 7(1):21-27. <https://doi.org/10.19184/nlj.v7i1.22116>
20. Kumala IN. Gambaran histopatologi hepar tikus putih (*Rattus norvegicus*) yang diinduksi dengan parasetamol dosis toksik pasca pemberian ekstrak etanol daun kelor (*Moringa oleifera*). *Jurnal Kimia Riset* 2017; 2(2):123-130. <https://doi.org/10.20473/jkr.v2i2.6700>
21. Mondal SK. Manual of histological technique. Mumbai: Jaypee Brothers Medical Publisher (P) Ltd. 2019.
22. Li W, Tang R, Ouyang S, Ma F, Liu Z, Wu J. Folic acid prevents cardiac



- dysfunction and reduces myocardial fibrosis in a mouse model of high-fat diet-induced obesity. *Nutr Metab (Lond)* 2017; 14:68.  
<https://doi.org/0.1186/s12986-017-0224-0>
23. Akbarzadeh A, Norouzian D, Mehrabi MR, Jamshidi Sh, Farhangi A, Verdi AA, *et al.* Induction of diabetes by streptozotocin in rats. *Indian J Clin Biochem* 2007; 22(2):60-4.  
<https://doi.org/10.1007/BF02913315>
  24. Jaishree V, Narsimha S. Swertiamarin and quercetin combination ameliorates hyperglycemia, hyperlipidemia and oxidative stress in streptozotocin-induced type 2 diabetes mellitus in wistar rats. *Biomed Pharmacother* 2020; 130:110561.  
<https://doi.org/10.1016/j.biopha.2020.110561>
  25. Lew WYW, Bayna E, Dalle Molle E, Contu R, Condorelli G, Tang T. Myocardial fibrosis induced by exposure to subclinical lipopolysaccharide is associated with decreased miR-29c and enhanced NOX2 expression in mice. *PLoS One* 2014; 9(9):3-10.  
<https://doi.org/10.1371/journal.pone.0107556>
  26. Cheng D, Liang B, Li Y. Antihyperglycemic effect of ginkgo biloba extract in streptozotocin-induced diabetes in rats. *Biomed Res Int* 2013; 2013:162724.  
<https://doi.org/10.1155/2013/162724>
  27. Vique-Sánchez JL, López-Palacios, Miranda-Ozuna, Benítez-Cardoza. Effects of W100E-Leptin in streptozotocin-induced diabetic mice. *Nutrición Clínica y Dietética Hospitalaria* 2020; 40(3):153-61.  
<https://doi.org/10.12873/403vique>
  28. Taher M, Tg Zakaria TMFS, Susanti D, Zakaria ZA. Hypoglycaemic activity of ethanolic extract of *Garcinia mangostana* Linn. in normoglycaemic and streptozotocin-induced diabetic rats. *BMC Complement Altern Med* 2016; 16(1):135.  
<https://doi.org/10.1186/s12906-016-1118-9>
  29. Hinderer S, Schenke-Layland K. Cardiac fibrosis – a short review of causes and therapeutic strategies. *Adv Drug Deliv Rev* 2019; 146:77-82.  
<https://doi.org/10.1016/j.addr.2019.05.011>
  30. Li CJ, Lv L, Li H, Yu DM. Cardiac fibrosis and dysfunction in experimental diabetic cardiomyopathy are ameliorated by alpha-lipoic acid. *Cardiovasc Diabetol* 2012; 11:73.  
<https://doi.org/10.1186/1475-2840-11-73>
  31. Ranjan P, Kumari R, Verma SK. Cardiac fibroblasts and cardiac fibrosis: precise role of exosomes. *Front Cell Dev Biol* 2019; 7:318.  
<https://doi.org/10.3389/fcell.2019.00318>
  32. Raish M, Ahmad A, Bin Jordan YA, Shahid M, Alkharfy KM, Ahad A, *et al.* Sinapic acid ameliorates cardiac dysfunction and cardiomyopathy by modulating NF- $\kappa$ B and Nrf2/HO-1 signaling pathways in streptozocin induced diabetic rats. *Biomed Pharmacother* 2022; 145:112412.  
<https://doi.org/10.1016/j.biopha.2021.112412>
  33. Liu Y, Qi H, Wang Y, Wu M, Cao Y, Huang W, *et al.* Allicin protects against myocardial apoptosis and fibrosis in streptozotocin-induced diabetic rats. *Phytomedicine* 2012; 19(8-9):693-8.  
<https://doi.org/10.1016/j.phymed.2012.04.007>
  34. Zhao Y, Li S, Quan E, Zhang H, Wu Y, Luo Y, *et al.* Trimetazidine inhibits cardiac fibrosis by reducing reactive oxygen species and downregulating connective tissue growth factor in

- streptozotocin-induced diabetic rats. *Exp Ther Med* 2019; 18(2):1477-85.  
<https://doi.org/10.3892/etm.2019.7705>
35. Aju BY, Rajalakshmi R, Mini S. Protective role of *Moringa oleifera* leaf extract on cardiac antioxidant status and lipid peroxidation in streptozotocin induced diabetic rats. *Heliyon* 2019; 5(12):e02935.  
<https://doi.org/10.1016/j.heliyon.2019.e02935>
36. Lin M, Zhang J, Chen X. Bioactive flavonoids in *Moringa oleifera* and their health-promoting properties. *J Funct Foods* 2018; 47:469-79.  
<https://doi.org/10.1016/j.jff.2018.06.011>
37. Trigo C, Castelló ML, Ortolá MD, García-Mares FJ, Desamparados Soriano M. *Moringa oleifera*: an unknown crop in developed countries with great potential for industry and adapted to climate change. *Foods* 2021; 10(1):31.  
<https://doi.org/10.3390/foods10010031>
38. Alegbeleye OO. How functional is *Moringa oleifera*? A review of its nutritive, medicinal, and socioeconomic potential. *Food Nutr Bull* 2018; 39(1):149-70.  
<https://doi.org/10.1177/0379572117749814>
39. Vergara-Jimenez M, Almatrafi MM, Fernandez ML. Bioactive components in *Moringa oleifera* leaves protect against chronic disease. *Antioxidants* 2017; 6(4):91.  
<https://doi.org/10.3390/antiox6040091>
40. Villarruel-López A, López-de la Mora DA, Vázquez-Paulino OD, Puebla-Mora AG, Torres-Vitela MR, Guerrero-Quiroz LA, *et al.* Effect of *Moringa oleifera* consumption on diabetic rats. *BMC Complement Altern Med* 2018; 18(1):127.  
<https://doi.org/10.1186/s12906-018-2180-2>