

On Rainbow k -Connection Number of Special Graphs and It's Sharp Lower Bound

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 855 012003

(<http://iopscience.iop.org/1742-6596/855/1/012003>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 103.241.206.147

This content was downloaded on 13/07/2017 at 02:50

Please note that [terms and conditions apply](#).

You may also be interested in:

[H-Supermagic Labeling on Coronation of Some Classes of Graphs with a Path](#)

H Sandaririra, M Roswitha and T A Kusmayadi

[H-Supermagic Labeling on Shrubs Graph and \$L_m - P_n\$](#)

Risala Ulfatihah, Mania Roswitha and Tri Atmojo Kusmayadi

[Answer to Sharp's letter \(1987\)](#)

H Weisen

[On the total H-irregularity strength of graphs: A new notion](#)

Ika Hesti Agustin, Dafik, Marsidi et al.

[The Kac-Ward combinatorial method](#)

N P Dolbilin, Yu M Zinov'ev, M A Shtan'ko et al.

[Primitive polycycles and helicenes](#)

M Deza and Mikhail I Shtogrin

[On the range of bond percolation thresholds for fully triangulated graphs](#)

John C Wierman

[The One Universal Graph — a free and open graph database](#)

Liang S. Ng and Corbin Champion

[Energy and Laplacian on Hanoi-type fractal quantum graphs](#)

Patricia Alonso-Ruiz, Daniel J Kelleher and Alexander Teplyaev

On Rainbow k -Connection Number of Special Graphs and It's Sharp Lower Bound

Ika Hesti Agustin^{1,2}, Dafik^{1,3}, Gembong A.W.^{1,2}, Ridho Alfarisi^{1,4}

¹ CGANT, University of Jember, Indonesia

² Mathematics Depart. University of Jember, Indonesia

³ Mathematics Edu. Depart. University of Jember, Indonesia

⁴ Department of Mathematics, Sepuluh Nopember Institute of Technology, Surabaya, Indonesia

E-mail: ikahesti.fmipa@unej.ac.id, d.dafik@unej.ac.id

Abstract. Let $G = (V, E)$ be a simple, nontrivial, finite, connected and undirected graph. Let c be a coloring $c : E(G) \rightarrow \{1, 2, \dots, s\}$, $s \in \mathbb{N}$. A path of edge colored graph is said to be a rainbow path if no two edges on the path have the same color. An edge colored graph G is said to be a rainbow connected graph if there exists a rainbow $u - v$ path for every two vertices u and v of G . The rainbow connection number of a graph G , denoted by $rc(G)$, is the smallest number of k colors required to edge color the graph such that the graph is rainbow connected. Furthermore, for an l -connected graph G and an integer k with $1 \leq k \leq l$, the rainbow k -connection number $rc_k(G)$ of G is defined to be the minimum number of colors required to color the edges of G such that every two distinct vertices of G are connected by at least k internally disjoint rainbow paths. In this paper, we determine the exact values of rainbow connection number of some special graphs and obtain a sharp lower bound.

Keywords: Rainbow k -Connection Number, Special Graphs, Sharp Lower Bound

1. Introduction

Suppose G is a simple connected graph with a set of vertices $V(G)$ and edges $E(G)$. For a further reference please see Gross, *et. al.* [6]. Let G be a nontrivial connected graph on which it is defined a coloring $c : E(G) \rightarrow \{1, 2, \dots, s\}$, $s \in \mathbb{N}$, of the edges of G , where adjacent edges may be colored the same. A $u - v$ path P in G is a rainbow path if no two edges of P are colored the same. The graph G is rainbow-connected (with respect to c) if G contains a rainbow $u - v$ path for every two vertices u and v of G . In this case, the coloring c is called a rainbow coloring of G . If k colors are used, then c is a rainbow k -coloring. The minimum k for which there exists a *rainbow k -coloring* of the edges of G is the rainbow connection number $rc(G)$. The completes concept can be found in Chartrand in [4].



A simple observation can be proposed that if G has n vertices then $rc(G) \leq n - 1$ but is not sharp. Since a given spanning tree can be assigned with distinct colors, and color the remaining edges with one of the already used colors then the upper bound of $rc(G) \leq n - 1$, see Caro [1] for detail. It is also easy to understand that $rc(G) \geq diam(G)$, where $diam(G)$ denotes the diameter of G , Caro in [1]. Thus, it gives the following

$$diam(G) \leq rc(G) \leq n - 1$$

There have been some results regarded to rainbow connection numbers. Chandran, *et.al.* in [2] determined rainbow connection number and connected dominating sets, Chakraborty, *et.al.* in [3] considered hardness and algorithms for rainbow connectivity. Furthermore, Li *et.al.* in [7] stated Rainbow connections of graphs - A survey. Also Li *et.al.* in [8] characterized graphs with rainbow connection number and rainbow connection numbers of some graph operations. Schiermeyer in [10] studied rainbow connection in graphs with minimum degree three.

A well-known result shows that in every l -connected graph G with $l \geq 1$, there are k internally disjoint $u - v$ paths connecting any two distinct vertices u and v for every integer k with $1 \leq k \leq l$ [9]. Chartrand *et al.* [5] defined the rainbow k -connectivity $rc_k(G)$ of G to be the minimum integer j for which there exists a j -edge-coloring of G such that for every two distinct vertices u and v of G , there exist at least k internally disjoint $u - v$ rainbow paths.

By the definition of rainbow k -connectivity $rc_k(G)$, we realize that it is almost impossible to derive the exact value or a nice bound of the rainbow k -connectivity for a general graph G [9]. To answer the problem: given that any connected graph G , determine the rainbow connection number $rc_k(G)$ of any graph G ? It tends to be NP-hard problem. Thus, the study of rainbow k -connectivity of some classes of special graphs is still needed. In this paper we will study the rainbow connection number $rc_k(G)$ of Triangular Ladder, Wheel graphs, and edge comb of graph $G = C_n \triangleright TL_m$ and $G = C_n \triangleright K_m$. The edge comb between L and H , denoted by $L \triangleright H$, is a graph obtained by taking one copy of L and $|E(L)|$ copies of H and grafting the i -th copy of H at the i -th edges of L . The result show that all the rainbow k -connection number $rc_k(G)$ of the graph studied in this paper achieve the minimum value.

2. The Results

Before presenting the main results we need to establish the lower bound of $rc_k(G)$ of any graph G such that the graph G is considered to be a k -connected graph. Note that the length of the shortest graph cycle (if any) in a given graph is known as a *girth*, and the length of a longest cycle is known as the *graph circumference*.

Theorem 1. *Let $d(u, v)$ be a distance between u and v , $C(u, v)$ is a shortest cycle that contains the vertices u and v . If G is 2-connected graph then $rc_2(G) \geq \max \{|C(u, v)| - d(u, v), \forall u, v \in V(G)\}$, where $C(u, v)$ and $d(u, v)$ are in one cycle.*

Proof. Let G be a connected cyclical graph. Thus, the length of second alternative internally disjoint rainbow path for any two vertices u and v is $|C(u, v)| - d(u, v)$ where $C(u, v)$ is a girth that contain vertices u and v . The greatest lower bound of

$rc_2(G) \geq \max \{|C(u, v)| - d(u, v)\}$. By contradiction, if we color the edges of G by any value less than $\max \{|C(u, v)| - d(u, v)\}$ then there exist two vertices u and v that do not present two internally disjoint paths. \square

We can extend the theorem for l -connected graph.

Lemma 1. *If G is l -connected graph, $l \geq 2$, then for every two vertices $u, v \in V(G)$, there exist at least $l - 1$ cycles of G containing the vertices u and v .*

Proof. We can prove this theorem by contradiction. Suppose that there exist two vertices $u, v \in V(G)$ that contain one less than $l - 1$ cycles of G . Suppose that the number of cycles that contain $u, v \in V(G)$ is $l - k$ where $k \geq 2$. The set $\{C_i | 1 \leq i \leq l - k\}$ is $l - k$ cycles that contain any two vertices in $V(G)$. One cycle is used to make two internally disjoint paths between u and v . Two cycles are used to make three internally disjoint paths between u and v . Since u and v are on $l - k$ cycles then the number of disjoint paths between u and v is $l - k + 1$. Since $k \geq 2$ and we have two vertices with $l - k + 1$ disjoint paths connecting u and v , then G is $(l - k + 1 < l)$ -connected graph. It is a contradiction. \square

Theorem 2. *Let $d(u, v)$ be a distance between u and v , $C_i(u, v)$ be a shortest cycles that contain vertices u and v . Let C_i be cycles whose their common edge is uv . If G is l -connected graph then $rc_l(G) \geq \max\{\max\{|C_i(u, v)| - d(u, v), 1 \leq i \leq l - 1\}, \forall u, v \in V(G)\}$, where $C(u, v)$ and $d(u, v)$ are in one cycle.*

Proof. If G is l -connected graph, then by Lemma 1 every vertex in $V(G)$ lays on at least $l - 1$ cycles. Suppose the element of $\{C_i(u, v) | 1 \leq i \leq l - 1, u, v \in V(G)\}$ have $l - 1$ cycles containing $u, v \in V(G)$, the $l - 1$ cycles that contain u and v has to be minimum of size $|C_i(u, v)|$. The number of $rc_k(G)$ is at least $\max\{|C_i(u, v)| - d(u, v), 1 \leq i \leq l - 1\}$. Otherwise there exist two vertices u, v that do not give k internally disjoint rainbow path. \square

Now we will present some classes of graphs which can be determined their rainbow k -connection number.

Theorem 3. *Let G be a triangular ladder graph, the rainbow 2-connection number of G is $rc_2(G) = n$.*

Proof. Suppose $G = TL_n$. The graph G has vertex set $V(G) = \{x_i, y_i; 1 \leq i \leq n\}$ and edge set $E(G) = \{x_i x_{i+1}, y_i y_{i+1}, x_i y_{i+1}; 1 \leq i \leq n - 1\} \cup \{x_i y_i; 1 \leq i \leq n\}$. Define a color c of the edges $c : E(G) \rightarrow \{1, 2, \dots, s\}, s \in N$:

$$c(e) = \begin{cases} n - i & , e \in \{x_i x_{i+1}; 1 \leq i \leq n - 1\} \\ i & , e \in \{y_i y_{i+1}; 1 \leq i \leq n - 1\} \\ 1 & , e \in \{x_i y_{i+1}; 1 \leq i \leq n - 1\} \cup \{x_1 y_1\} \\ n & , e \in \{x_i y_i; 2 \leq i \leq n\} \end{cases}$$

It is easy to see that the color $c(e)$ reach a maximum value when $e = x_i y_i$ and $c(e) = n$. Thus, $rc_2(G) \leq n$. Now we will show that $rc_2(G) \geq n$. Consider the vertex $u = y_1$ and $v = x_n$. The vertex u and v lay on the cycle of size $2n$. Since distance, $d(u, v) = n$, then by Theorem 1, we have $rc_2(G) \geq 2n - n = n$. It concludes that $rc_2(G) = n$. \square




Figure 1. Graph $G = TL_5$ with $rc_2(G) = 5$

Theorem 4. Let G be a wheel graph of order $n + 1$, the rainbow 3-connection number G is $rc_3(W_n) = n$.

Proof. Given that $G = W_n$. The graph G has vertex set $V(G) = \{x_i; 1 \leq i \leq n\} \cup \{A\}$ and edge set $E(G) = \{Ax_i; 1 \leq i \leq n\} \cup \{x_i x_{i+1}; 1 \leq i \leq n - 1\} \cup \{x_1 x_n\}$. Define a color c of the edges $c : E(G) \rightarrow \{1, 2, \dots, s\}, s \in \mathbb{N}$:

$$c(e) = \begin{cases} i, & e \in \{x_i x_{i+1}; 1 \leq i \leq n - 1\} \cup \{Ax_i; 1 \leq i \leq n\} \\ n, & e \in \{x_1 x_n\} \end{cases}$$

It is easy to see that the color $c(e)$ reach a maximum value when $e = x_1 x_n$, thus $rc_3(W_n) \leq n$. No we will show that $rc_3(W_n) \geq n$. We will use a contradiction. Suppose that $rc_3(W_n) \leq n - 1$, take $rc_3(W_n) = n - 1$. Consider edge set $E' = \{x_i x_{i+1} | 1 \leq i \leq n - 1\} \cup \{x_1 x_n\}$ and $|E'| = n + 1$. If we color $n + 1$ edges of E' by $n - 1$ colors, then there exist $e_1, e_2 \in E'$ such that $c(e_1) = c(e_2)$, without loss of generality we can choose $e_1 = x_1 x_2$ and $e_2 = x_i x_{i+1}$. Since W_n is 3-connected graph and $rc_3(W_n) = n - 1$ then there must exist three disjoint paths between any two vertices. Consider vertex x_1 and vertex x_{i+1} which give three disjoint paths between x_1 and x_{i+1} . The first possible rainbow path is $x_1 A x_{i+1}$, the second is $x_1 x_n x_{n-1} \dots x_j$, however the third path $x_1 x_2 \dots x_i x_{i+1}$, for x_1 and x_{i+1} is not rainbow path as $c(x_1 x_2) = c(x_i x_{i+1})$. It is a contradiction, thus $rc_3(W_n) \geq n$. It concludes $rc_3(W_n) = n$. \square

Theorem 5. If $G = C_n \triangleright TL_m$ then $rc(G) = \frac{n}{2} + 2m - 2$ for n even and $rc_2(G) = 2m + 1$ for $n = 4$.

Proof. The graph $G = C_n \triangleright TL_m$ is a connected graph with vertex set $V(G) = \{x_i | 1 \leq i \leq n\} \cup \{y_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m - 1\} \cup \{z_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m - 1\}$ and edge set $E(G) = \{x_i x_{i+1} | i \leq i \leq n - 1\} \cup \{x_n x_1\} \cup \{x_i y_{i,1} | 1 \leq i \leq n\} \cup \{x_{i+1} z_{i,1} | 1 \leq i \leq n - 1\} \cup \{x_1 z_{n,1}\} \cup \{y_{i,j} y_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m - 2\} \cup \{z_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m - 2\} \cup \{y_{i,j} z_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m - 1\} \cup \{x_i z_{i,1} | 1 \leq i \leq n\} \cup \{y_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m - 2\}$. The value $|V(G)| = n(2m - 1)$ and $|E(G)| = 3n + 2n(m - 2) + 2n(m - 1)$. The diameter of G , $diam(G) = \frac{n}{2} + 2(m - 1)$. The number $rc(G)$ is given by the following




Figure 2. Graph $G = W_6$ with $rc_3(W_6) = 6$

coloring function:

$$c(e) = \begin{cases} i \bmod \frac{n}{2} & , e \in \{x_i x_{i+1} | 1 \leq i \leq n-1\} \cup \\ & \{y_{i,j} z_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m-1\} \\ n \bmod \frac{n}{2}, & e \in \{x_n x_1\} \\ \frac{n}{2} + 1 & , e \in \{x_i y_{i,1} | 1 \leq i \leq n\} \cup \{x_i z_{i,1} | 1 \leq i \leq n\} \\ \frac{n}{2} + 1 + j & , e \in \{y_{i,j} y_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \\ & \cup \{y_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \\ \frac{n}{2} + m & , e \in \{x_{i+1} z_{i,1} | 1 \leq i \leq n-1\} \cup \{x_1 z_{n,1}\} \\ \frac{n}{2} + m + j & , e \in \{z_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \end{cases}$$

The maximum value of function is $c(e) = \frac{n}{2} + 2m - 2$ so $rc(G) \leq \frac{n}{2} + 2m - 2$. By applying Inequality 1 $rc(G) \geq \frac{n}{2} + 2m - 2$, it implies that $rc(G) = \frac{n}{2} + 2m - 2$.

The number $rc_2(G) \geq 2m + 1$ for $n = 4$ and any m , is obtained by coloring mapping:

$$c(e) = \begin{cases} i \bmod 2 & , e \in \{x_i x_{i+1} | 1 \leq i \leq 3\} \\ 2 & , e \in \{x_4 x_1\} \\ 3 & , e \in \{x_i y_{i,1} | 1 \leq i \leq n\} \cup \{x_i z_{i,1} | 1 \leq i \leq n\} \\ 3 + j & , e \in \{y_{i,j} y_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \\ & \cup \{y_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \\ m + 2 & , e \in \{x_{i+1} z_{i,1} | 1 \leq i \leq n-1\} \cup \{x_1 z_{n,1}\} \\ m + 2 + j & , e \in \{z_{i,j} z_{i,j+1} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \\ 2m + 1 & , e \in \{y_{i,j} z_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m-1\} \end{cases}$$

To prove $rc_2(G) \leq 2m + 1$, consider vertex $y_{2,m-1}$ and vertex $z_{1,m-1}$, the vertex $y_{2,m-1}$ and vertex $z_{1,m-1}$ lay on cycle of size of at least $4m - 1$. The distance between $y_{2,m-1}$ and $z_{1,m-1}$ is $2(m - 1)$ so the length of remaining shortest path between $y_{2,m-1}$ and $z_{1,m-1}$ is $2m + 1$. This path is the shortest alternative path from $y_{2,m-1}$ to $z_{1,m-1}$ to get the second internally disjoint rainbow path. \square




Figure 3. Graph edge comb $G = C_4 \triangleright TL_3$ with $rc_2(G) = 7$.

Theorem 6. If $G = C_n \triangleright K_m$, then the number $rc(G) = \frac{n}{2} + 1$ for n even and $rc_2(G) = 4$, for $n = 4$.

Proof. The graph $G = C_n \triangleright K_m$ is a connected graph with vertex set $V(G) = \{x_i | 1 \leq i \leq n\} \cup \{y_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m-2\}$ and edge set $E(G) = \{x_i x_{i+1} | 1 \leq i \leq n-1\} \cup \{x_n x_1\} \cup \{x_i y_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \cup \{x_{i+1} y_{i,j} | 1 \leq i \leq n-1, 1 \leq j \leq m-2\} \cup \{x_1 y_{n,j} | 1 \leq j \leq m-2\} \cup (\bigcup_{l=1}^{m-3} (\{y_{i,l} y_{i,j+l} | 1 \leq i \leq n, 1 \leq j \leq m-2-l\}))$. The number of vertices and edges of G is $|V(G)| = n + n(m-2)$ and $|E(G)| = n(1 + 2(m-2)) + \frac{(m-2)(m-3)}{2}$. The Diameter of G , $diam(G) = \frac{n}{2} + 1$

The value $rc(G) = \frac{n}{2} + 1$ obtained by the following edge mapping function:

$$c(e) = \begin{cases} i \bmod \frac{n}{2} & , e \in \{x_i x_{i+1} | 1 \leq i \leq n-1\} \cup \\ & \{x_i y_{i,j} | 1 \leq i \leq n, 1 \leq j \leq m-2\} \cup \\ & (\bigcup_{l=1}^{m-3} (\{y_{i,l} y_{i,j+l} | 1 \leq i \leq n, \\ & 1 \leq j \leq m-2-l\})) \\ n \bmod \frac{n}{2} & , e \in \{x_n x_1\} \\ \frac{n}{2} + 1 & , e \in \{x_1 y_{n,j} | 1 \leq j \leq m-2\} \cup \\ & \{x_{i+1} y_{i,j} | 1 \leq i \leq n-1, 1 \leq j \leq m-2\} \end{cases}$$

The maximum value of $c(e)$ is $\frac{n}{2} + 1$ so $rc(G) \leq \frac{n}{2} + 1$, by applying Inequality 1 $rc(G) \geq \frac{n}{2} + 1$ and finally we get $rc(G) = \frac{n}{2} + 1$.




Figure 4. Graph edge comb $C_6 \supseteq K_5$ with $rc(G) = 4$.

The value $rc_2(G) \geq 4$ for $n = 4$ and any m , is obtained by the following

$$c(e) = \begin{cases} i \bmod 2 & , e \in \{x_i x_{i+1} | 1 \leq i \leq 3\} \cup \{x_i y_{i,j} | 1 \leq i \leq 4, \\ & 1 \leq j \leq m-2\} \cup \{y_{i,j} y_{i,j+1} | 1 \leq i \leq 4, \\ & 1 \leq j \leq m-3\} \\ 4 \bmod 2 & , e \in \{x_4 x_1\} \\ 3 & , e \in \{x_1 y_{4,j} | 1 \leq j \leq m-2\} \cup \\ & \{x_{i+1} y_{i,j} | 1 \leq i \leq 3, 1 \leq j \leq m-2\} \\ 4 & , e \in \{x_i y_{i,j} | 1 \leq i \leq 4, 1 \leq j \leq m-2\} \\ & \cup (\bigcup_{l=1}^{m-3} (\{y_{i,l} y_{i,j+l} | 1 \leq i \leq 4, \\ & 1 \leq j \leq m-2-l\} - \{y_{i,j} y_{i,j+1} | 1 \leq i \leq 4, \\ & 1 \leq j \leq m-3\})) \end{cases}$$

To prove $rc_2(G) \leq 4$ consider vertex $y_{1,j}$ and $y_{2,k}$ for $1 \leq j, k \leq m-2$. This vertices is contained on cycle with size at least 6. The distance between $y_{1,j}$ and $y_{2,k}$ is 2 so the lengt of remaining shortest path between $y_{1,j}$ and $y_{2,k}$ is 4. This path is the shortest alternative path from $y_{1,j}$ to $y_{2,k}$ to make second internally disjoint rainbow path. \square

Concluding Remarks

We have studied the rainbow k -connection number of G . The result show that all the rainbow k -connection number $rc_k(G)$ of the graph studied in this paper achieve the minimum value. We have also characterized any graph to have a minimum k -connection number, through the following theorem: If G is l -connected graph then

$rc_l(G) \geq \max \{|C_i(u, v)| - d(u, v), 1 \leq i \leq l-1\}$, where $|C_i(u, v)|$ is a girth that contains the vertices u and v . However, it is just lower bound, we have not found the sharper upper bound of $rc_k(G)$ of any graph. Thus we propose the following open problem.

Open Problem 1. *Given that any connected graph G , determine a sharp upper bound of the rainbow k -connection number $rc_k(G)$ of G .*

Acknowledgement

We gratefully acknowledge the support from FUNDAMENTAL Ristek Dikti Grant 2017 & CGANT - University of Jember.

Reference

- [1] Y. Caro, A. Lev, Y. Roditty, Z. Tuza, R. Yuster, On rainbow connection, *Electron. J. Combin.* **15**, R57, 2008
- [2] L.S. Chandran, A. Das, D. Rajendraprasad, N.M. Varma, Rainbow connection number and connected dominating sets, Arxiv preprint arXiv:1010.2296v1 [math.CO], 2010
- [3] S. Chakraborty, E. Fischer, A. Matsliah, R. Yuster, Hardness and algorithms for rainbow connectivity, 26th International Symposium on Theoretical Aspects of Computer Science STACS 2009, 243-254, 2009
- [4] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang, Rainbow connection in graphs, *Math. Bohem.* **133**, 85-98, 2008
- [5] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang, The rainbow connectivity of a graph, *Networks* **54** (2), 75-81, 2009
- [6] J.L. Gross, J. Yellen and P. Zhang, *Handbook of Graph Theory*, Second Edition, CRC Press, Taylor and Francis Group, 2014
- [7] X. Li, Y. Sun, Rainbow connections of graphs - A survey, arXiv:1101.5747v2 [math.CO], 2011
- [8] X. Li, Y. Sun, Characterize graphs with rainbow connection number $m-2$ and rainbow connection numbers of some graph operations, Preprint, 2010
- [9] Xueliang Li, Yuefang Sun, On the rainbow k -connectivity of complete graphs, *Australian Journal Of Combinatorics*, 217-226, 2011
- [10] I. Schiermeyer, Rainbow connection in graphs with minimum degree three, *IWOCA 2009*, LNCS 5874, 2009