

International Journal on Advanced Science Engineering Information Technology

Available online at
<http://ijaseit.insightsociety.org>



Advanced Science Image processing Image retrieval microprocessor
Back Projection fuzzy Engineering alumina Gait analysis Biometric
Mobile Agent Interpolation Soft Tissue hydrothermal stability Outlier filter
Sampling Theorem Magnetic field porous structure acidity Wireless
Parsimony Database Server Robot Operation Life Cycle Cost
Graphical User Interface Virus Compensate Strength Energy Saving
Current shaping Phylogenetic Analysis Sensor Networks Energy Optimization
Metamorphic Virus neural network Predictive Tools Image matching
parallel port Obfuscation Techniques Admixture Router Image matching
Power factor Boost regulator catalyst Construction Delay Predicted Strength Regression Analysis
breast phantom Effects of Delay Support Vector Machine Current harmonics Surveillance
Information Technology soybean

Editorial Team

Editor in Chief

1. Sri Atmaja Putra (EDITOR IN CHIEF)

Associate Editors

1. Prof Taufik, California Polytechnic State University, USA, United States
2. Moustafa M Eissa, Helwan University, EGYPT
3. Wan Mohtar Wan Yusoff, Universiti Kebangsaan Malaysia, MALAYSIA
4. Halimah Badioze Zaman, Universiti Kebangsaan Malaysia, MALAYSIA
5. Son Radu, Universiti Putra Malaysia, Malaysia, MALAYSIA
6. Mohd Razi Ismail, Universiti Putra Malaysia, MALAYSIA
7. Dr Novizar Nazir, Andalas University, Indonesia
8. Haitham Alali, Amman Arab University, JORDAN
9. Shahrul Azman Mohd Noah, Universiti Kebangsaan Malaysia, MALAYSIA

Managing Editor

1. Rahmat Hidayat

Editors

1. Ifa Puspasari, Fuel Cell Institute, Universiti Kebangsaan Malaysia, Malaysia
2. Mr. Ario Betha Juanssilfero
3. Mr. Zairi Ismael Rizman, Universiti Teknologi MARA, Malaysia
4. Slamet Riyadi
5. Adhi Harmoko Saputro
6. Dr. Gusri Yaldi, Politeknik Negeri Padang
7. Dr Abrar Ismardi, Telkom University, Indonesia



Vol 9, No 3 (2019)

Table of Contents

Articles

- Fuzzy Rule Interpolation and SNMP-MIB for Emerging Network Abnormality
Mohammad Almseidin, Mouhammd Al-kasassbeh, Szilveszter Kovacs 735-744
- Detection of Scratches on Cars by Means of CNN and R-CNN
César Giovany Pachón Suescún, Javier Orlando Pinzón Arenas, Robinson Jiménez Moreno 745-752
- Optimization of Biochemical Systems Production Using Combination of Newton Method and Particle Swarm Optimization
Mohd Arfian Ismail, Vitaliy Mezhuyev, Irfan Darmawan, Shahreen Kasim, Mohd Saberi Mohamad, Ashraf Osman Ibrahim 753-758
- A New Native Video Filtering based on OpenGL ES for Mobile Platform
Sari Wijayanti, Burhan Alfironi Muktamar, Sunu Wibirama, Agus Bejo 759-765
- Enhanced Manhattan-based Clustering using Fuzzy C-Means Algorithm for High Dimensional Datasets
Joven A. Tolentino, Bobby D. Gerardo 766-771
- Static Knowledge-Based Authentication Mechanism for Hadoop Distributed Platform using Kerberos
Thoyazan Sultan Algaradi, Rama B 772-780
- Liquefaction Analysis of Beach Protection Structure: Numerical Simulations
Abdul Hakam, - Junaidi, Bayu M Adji, Shafira R Hape, M Sofian Asmirza 781-786
- The Improvement of Yield and Quality of Soybeans in a Coastal Area Using Low Input Technology Based on Biofertilizers
Yudhy Harini Bertham, Zainal Arifin, Abimanyu Dipo Nusantara 787-791
- Evaluation of Full Interaction Composite Columns Subjected to Static and Dynamic Loading
Mohammed J. Hamood, Zinah Asaad Abdul Husain, Maha Ghalib Ghaddar 792-798
- Effects of Laminate Arrangement on the Failure Behaviour of Hybrid Composite Plates under Transverse Sinusoidal Load
Ahmad Fuad Ab Ghani, Azizul Hakim Samsudin, Shaheera Diana Mohd Sharom, Yongtao Lu, Jamaluddin Mahmud 799-803
- Impact of Interval Irrigation on Amino Acids Composition and Minerals Content of Potato Tubers Granola and Eigenheimer Cultivars
Nurpilihan Bafdal, Condro Wibowo 804-809
- Development of Salt Leached Silk Fibroin Scaffold using Direct Dissolution Techniques for Cartilage Tissue Engineering
Untung Ari Wibowo, Hermawan Judawisastra, Anggraini Barlian, Nayla M. Alfarafisa, Karina F Moegni, Melinda Remelia 810-815
- Study of Lateral Vibration of Drilling String with Mass Imbalance Using Finite Element
Sonny Irawan, Arun Kumar Shanmugam, Mas Irfan P. Hidayat, Totok R Biyanto 816-821
- Use of organic waste as an alternative organic fertilizer and synthetic fertilizer to ameliorate acid soil productivity
- Ermadani, - Hermansah, - Yulnafatmawita, Auzar Syarif, Ismon Lenin 822-828
- Effect of Starter and Biopreservative from Lactic Acid Bacteria and Yeast on Chemical, Physical and Organoleptic Qualities of Mutton Salami

<i>Lilis Suryaningsih, Rahmat Hidayat, Gemilang Lara Utama, Andry Pratama, Roostita L Balia</i>	829-834
The Demarketing of Energy Drinks Using Facebook Media: A Healthcare Perspective <i>Haitham Alali, Raed Wishah, Ghazi Al-Weshah</i>	835-840
Use of Virtual Reality as a Tool to Overcome the Post-Traumatic Stress Disorder of Pensioners <i>Ivan Kovar</i>	841-848
Water Potential Management and Arrangement of River Estuary Area for the Mitigation of the Climate Change in Bali <i>I Gusti Agung Putu Eryani, I Wayan Runa, Made Widya Jayantari</i>	849-854
Isolation and Characterization of Lactic Acid Bacteria Producing GABA from Indigenous West Sumatera Fermented Food <i>Lili Anggraini, Yetti Marlida, - Mirzah, - Wizna, - Jamsari, Nurul Huda</i>	855-860
Evaluation of Scale-Up And Environmental Factors On Microbial Fuel Cell <i>Tun Ahmad Gazali, M. Azizul Moqsud</i>	861-865
Preparation and Characterization of Biopolymer Chitosan Nanofiber from Coconut Crab Shell <i>Hamidin Rasulu, Dinar Praseptiangga, I Made Joni, Ari Handono Ramelan</i>	866-873
Study of Elastic Modulus Determination of Polymers with Ultrasonic Method <i>Firmansyah Sasmita, Tania Zefanya S. Tarigan, Hermawan Judawisastra, Toni Agung Priambodo</i>	874-879
Modeling and Experimental Study of Al-Cu Alloy Sand Casting for Circuit Breaker <i>Dian Mughni Fellicia, M.I.P. Hidayat, Rochman Rochiem, L.B. Aditya Putra, A. T. Wibisono, Mavindra Ramadhani</i>	880-887
In Vitro Regeneration of Sugar Palm (Arenga pinnata Wurmb Merr.) <i>Nazatul Asikin Muda, Asmah Awal, Mohd Yusoff Abdullah, Nor Azma Yusuf, Nor Azma Yusuf</i>	888-894
Kinetics of CO ₂ Absorption into Aqueous MDEA Solution Promoted by Mixture of Potassium Salt of L-Arginine and L-Glutamic Acid <i>Toto Iswanto, Ali Altway, Maya Shovitri, Muhammad Haikal, Septiani Ayustiningrum, Tri Widjaja</i>	895-903
Physicochemical Properties of Pectin Extracted from Selected Local Fruits By-Product <i>Aida FMNA, Norasmanizan Zailan</i>	904-910
The Influence of Type of Solvent and Extraction Temperature of Corn Silk Extracts <i>- Haslina, Novizar Nazir, Sri Budi Wahjuningsih, Dewi Larasati</i>	911-915
Effects of Leucaena leucocephala Supplementation to Total Mixed Ration Based on Ammoniated Rice Straw Onfiber Digestibility and Rumen Fermentation Characteristics in vitro <i>Rusmana Wijaya Setia Ningrat, Mardiaty Zain, - Erpomen, Ezi Masdia Putri, Malik Makmur</i>	916-921
Overtopping wave reduction capability of Rakuna IV armor block used for Rubblemound Breakwater <i>Giang. Le Thi Huong</i>	922-928
Experimental Test of Masonry Wallets Retrofitted by ABACA Fiber Reinforced Mortar <i>Etri Suhelmidawati, Fauna Adibroto, - Mukhlis, Suhendrik Hanwar, Muneyoshi Numada, Kimiro Meguro</i>	929-935

From Parking to Pollution: Investigating the Effect of 45' Parking on Environment - <i>Anugrah, - Soemarno, Achmad Wicaksono, Lambang Basri Said</i>	936-945
Detection of Changes in Surface Water Bodies Urban Area with NDWI and MNDWI Methods <i>Muhammad Ichsan Ali, Gufran Darma Dirawan, Abdul Hafid Hasim, Muh Rais Abidin</i>	946-951
Combining ABC and VED Analysis for Managing Medicine Inventory (Case Study at Community Development Elderly in Indonesia) <i>Fitra Lestari, - Ulfah, - Suherman, Budi Azwar, Prima Fithri</i>	952-959
Automatic Rule Generator via FP-Growth for Eye Diseases Diagnosis <i>Rahmad Kurniawan, Mohd Zakree Ahmad Nazri, Siti Norul Huda Abdullah, Jemaima Che Hamzah, Rado Yendra, Westi Oktaviana</i>	960-966
Analysis of Attribute Selection and Classification Algorithm Applied to Hepatitis Patients <i>Sherylaidah Samsuddin, Zuraini Ali Shah, RD Rohmat Saedudin, Shahreen Kasim, Choon Sen Seah</i>	967-971
Application of The KNN Algorithm in Determining the Orientation of The Probability Area Containing The Ship Position by GPS Systems on Hai Phong Coastal Area <i>Thai Duong Nguyen, Trong Duc Nguyen</i>	972-978
Comparison of Nelder Mead and BFGS Algorithms on Geographically Weighted Multivariate Negative Binomial <i>Yuliani .S. Dewi, - Purhadi, - Sutikno, Santi. W. Purnami</i>	979-987
Implementation of The Variable Data Transmission System of Abnormal ECG with Activity State <i>Yun-Hong Noh, Ji-Yun Seo, Hosoo Nam, Do-Un Jeong</i>	988-992
Generating and Validating DSA Private Keys from Online Face Images for Digital Signatures <i>Asraa Safaa Ahmed, Firas A. Abdullatif, Taha Mohammad Hasan</i>	993-998
A New Feature Extraction Algorithm to Extract Differentiate Information and Improve KNN-based Model Accuracy on Aquaculture Dataset <i>Oskar Natan, Agus Indra Gunawan, Bima Sena Bayu Dewantara</i>	999-1007
Critical Information for Vietnamese Economy Aiming at a Strategic Breakthrough as Approaching the Industry 4.0 <i>Van Tai Pham</i>	1008-1016
Integration Training Information in Vietnam Maritime University Based on the Conceive-Design-Implement-Operate <i>Thi Anh Thu Do, Quoc Tien Le, Thi Ngoc Diep Hoang</i>	1017-1024
Local Descriptor Approach to Wrist Vein Recognition with DVH-LBP Domain Feature Selection Scheme <i>Dini Fronitasari, - Basari, Dadang Gunawan</i>	1025-1032
Designing Portable Shopping Trolley with Scooter Using Kansei Engineering Approach <i>Alex Kisanjani, Hari Purnomo</i>	1033-1038
Effects of Frequency and Mass of Eccentric Balls on Picking Force of The Coffee Fruit for The As-Fabricated Harvesting Machines <i>Minh Quang Chau, Vo Tien Nguyen</i>	1039-1045
Genetic Diversity of Black and Aromatic Rice Cultivar (<i>Oryza sativa</i> L.) from Various Regions in Indonesia using Random Amplified Polymorphic DNA Markers (RAPD) <i>Ummi Sholikhah, - Parjanto, Tri Handoyo, Ahmad Yunus</i>	1046-1051
Effect of Temperature on Synbiotic Sorghum Drinks Characteristics <i>Een Sukarminah, Endah Wulandari, Elazmanawati Lembong, Nurul Annisa</i>	1052-1057

Digital Repository Universitas Jember

The Glycemic Index (GI) of Adlay (*Coix lachryma-jobi var-mayuen*) on Processed Products

- *Tensiska, Imas Siti Setiasih, O. Suprijana, Warid Ali Qosim, Yana Cahyana*

1058-1062

Convolutional-NN and Word Embedding for Making an Effective Product Recommendation Based on Enhanced Contextual Understanding of a Product Review

- *Hanafi, Nanna Suryana, Abd Samad Bin Hasan Basari*

1063-1070

Open Data for Evidence-based Decision-making: Data-driven Government Resulting in Uncertainty and Polarization

Ahmad Luthfi, Marijn Janssen

1071-1078

Evaluating Services Computing Systems Engineering Framework using An Acceptance Model

Novianto Budi Kurniawan, - Suhardi, Yoanes Bandung, Yuli Adam Prasetyo, Purnomo Yustianto

1079-1085



Genetic Diversity of Black and Aromatic Rice Cultivar (*Oryza sativa* L.) from Various Regions in Indonesia using Random Amplified Polymorphic DNA Markers (RAPD)

Ummi Sholikhah^{#*}, Parjanto^{*}, Tri Handoyo^{*}, Ahmad Yunus^{*}

[#] Department of Agricultural Science, Graduate School of Sebelas Maret University (UNS), Surakarta 57126, Indonesia

^{*} Department of Agrotechnology, Faculty of Agriculture, Jember University (UNEJ), Jember 68121, Indonesia
E-mail: ummisholikhah.faperta@unej.ac.id, trihandoyo.faperta@unej.ac.id

^{*} Department of Agrotechnology, Faculty of Agriculture, Sebelas Maret University (UNS), Surakarta 57126, Indonesia
E-mail: parjanto_uns@yahoo.co.id, yunusuns17@gmail.com

Abstract— Indonesia is one of many countries with a high potential for genetic diversity resource of rice plants, both black rice, and aromatic rice. Therefore, the kinship relationship in each cultivar of which results are not influenced by the environment can be determined by molecular methods, one of which is by using RAPD analysis. This research employed 20 RAPD primers and DNA genome materials used in isolation from ten black rice cultivars, and 15 aromatic rice cultivars. The research stage began with a collection of black rice and aromatic rice germ plasma in several regions in Indonesia, seed nursery, DNA isolation, DNA quantity, and quality testing, primary selection, RAPD, running RAPD, and phylogenetic analysis. The objectives of this research were to obtain phylogenetic diversity of black and aromatic rice in this analysis. Twenty RAPD primers used to amplify 10 black rice cultivars and 15 aromatic rice cultivars obtained 180 polymorphic DNA Ribbons, with an average percentage of 89.2%. The genetic diversity of all cultivars analyzed was considerably high. At 84% coefficient, two major groups were obtained: black rice and aromatic rice, divided into eight small groups. The results of the phylogenetic analysis demonstrate high diversity between black rice and aromatic rice. Such a thing can be a first step for selecting rice plants which will be taken as plant parent in the crossbreeding.

Keywords— RAPD; cultivar; black rice; aromatic rice.

I. INTRODUCTION

Considerably-vast Indonesian territory with the characteristic of the agricultural country has enormous potential in terms of crops' genetic resources, especially rice plants. In the view of [1], Indonesia is one of the producers of black rice and aromatic rice in the world, where Indonesia ranks third in the distribution source of black rice accession in the world, with a distribution percentage of 7.2%. China ranks the highest in terms of access distribution at 62%, followed by Sri Lanka at 8.2%, India at 5.1%, and the Philippines at 4.3%, while Malaysia and Thailand trailing behind. With this potential, black rice and aromatic rice cultivars can be further developed as raw material for assembling superior varieties of aromatic black rice.

The need for rice, both in Indonesia and the world, is very extensive since rice is included as one of the staple food sources; and dependence on rice consumption nearly reaches 60%, while dominating the intake and consumption pattern

of staple foods at the present time. The community in general starts to selectively choose rice with health benefits, one of which is black rice. Black rice has high anthocyanin content, making its antioxidant activity useful as an anti-cancer [2], hypoglycemia [3], anti-inflammatory effects [4] It is also effective in reducing cholesterol level [5]-[10], As for aromatic rice, it has the advantage of which aroma is considerably fragrant, and the rice is relatively chewier or fluffier when cooked, thus enhancing the appetite of the people. Aromatic rice is rice that gives off aroma and it comes from 1-Proline volatile compound (2AP) Compound 2 AP in aromatic rice is 10 times higher, compared to non-aromatic rice [11]-[13].

From the advantages of each black rice and aromatic rice, it is highly important to examine genetic diversity further, via molecular and plant morphology, so that for a breeder can utilizes this into usable initial selection action in selecting the parent elders plant for crossbreeding and propagating superior varieties resulted from a combination

of both, [14] one of the important standard tools commonly used by biologists is a phylogenetic analysis; in which this analysis is capable of analyzing evolutionary relationship between organism, molecules, and phenotypes, allowing the results of phylogenetic analysis to produce kinship information. Genetic distance based on the similarity of genotypes is extremely necessary for a breeder to determine the parent plants and the evolutionary process occurred [15]. At the present time, the most widely used is the phylogenetic analysis of molecular results. Such thing is caused by molecular, thus allowing more characters to observe in concluding more accurate data and is free from environmental influences. This is far different from biochemical and morphological analysis, where the environment still influences the results of the analysis [16].

Molecular use [17,18], in phylogenetic analysis can be conducted using genetic markers. The function of markers is to select the properties to examine from one individual or to trace one genotype from a plant [19], RAPD is one of the markers widely used in the molecular analysis. In its use for a molecular genetic mapping application, the characteristic of RAPD is neutral and non-biased, and certain information on genome sequences is unnecessary. Based on the description above, the author conducted a series of research entitled Random Amplified Polymorphic DNA (RAPD) Genetic Diversity of Ten Black Rice Cultivars and Fifteen Aromatic Rice Cultivators (*Oryza sativa* L.) from Various Regions in Indonesia using RAPD. This research aimed to explore a genetic diversity of all cultivars of black rice and aromatic rice.

II. MATERIAL AND METHOD

A. Materials

1) Germ plasma collection from various locations in Indonesia

Germ plasma in the form of seeds was collected from various locations in Indonesia: 10 black rice cultivars and 15 aromatic rice cultivars as listed in Table 1.

2) Nursery and Planting of Rice Seeds (*Oryza sativa* L.)

The plant material of which leaves would be taken for extraction is prepared in advance via seed nursery and planting. The seedling was previously soaked in water for 4-5 days, and then after germination, the seeds were ready to be placed right on the ray pot, and after 12 days, it could be moved to a bucket filled with soil media.

3) Preparation for DNA Extract

DNA samples were taken from leaves of rice plants aged 2 weeks [20]. Then the measurement of DNA purity and concentration was done using spectrophotometry at 260 nm and 260/280 nm. The next step was DNA isolation by taking samples of young leaves of 0.5 grams which was then crushed using Liquid Nitrogen (N₂) and inserted into the Eppendorf filled with 500µl of complete extraction buffer (500µl Buffer Extraction + 25µl SDS 20% + 1, 25 β-mercapto). It was subsequently unsorted until reached homogeneity and was then incubated at 65°C for 10 minutes, swirled to taste, added with Potassium Acetate 5M 500µl, and incubated on ice for 10 minutes. Next, it was centrifuged

at 12,000 rpm for 10 minutes at 4°C. The supernatant was discarded and added with 500µl TE+15 µl RNase and was incubated for 1 hour. 500µl of PCI was added and then centrifuged of 12,000 rpm for 10 minutes at 4°C. The supernatant was transferred to the new Eppendorf and added with Chloroform Equal Volume, vortexed and then centrifuged at 12,000 rpm for 10 minutes at 4°C. The supernatant was transferred to the new Eppendorf, added with 0.8 isopropanol and 0.2 NaAc/ Sodium Acetate, swirled and incubated for 1 hour at -20°C, centrifuged of 12,000 rpm for 10 minutes at 4°C. The supernatant was discarded and added with 600µl of Ethanol PA 70% to the pellet. Centrifuge at 12,000 rpm 10 minutes at 4°C. The supernatant was discarded, the pellets were dried and added with 20µl of Buffer TE and 1µl of RNase, incubation of 37°C for 1 hour, and the results of the isolation were stored at -20°C.

TABLE I
RICE VARIETIES AND THEIR ORIGIN.

No	Name of Cultivar	Origin
1	Banjarnegara Black Rice	Banjarnegara, Central Java
2	Purbalingga Black Rice	Purbalingga, Central Java
3	Pari Ireng	Central Java
4	Purwokerto Black Rice	Purwokerto Central Java
5	Hare Lahok	Timor Leste
6	Bantul Black Rice	Bantul, Central Java
7	Lumajang Black Rice	Lumajang, East Java
8	Toraja	South Sulawesi
9	Melik Black Rice	Central Java
10	Blitar Black Rice	Blitar, East Java
11	Mentik Wangi	Pakuncen, Jogjakarta
12	Mentik Wangi Banjarnegara	Banjarnegara, Central Java
13	Mapan 05 Banjarnegara	Banjarnegara, Central Java
14	Situpatenggang	Kartuna/TB47H-MR-10
15	Pare Pulu Mandoti	Enrekang, South Sulawesi
16	Batang Gadis	IR 64/NDR 308//IR 64
17	Radah Putih Karanganyar	Karanganyar, Central Java
18	Mentik Susu Karanganyar	Karanganyar, Central Java
19	Situbagendit (non aromatic)	Batur/S2823-7d-8-1-A//S283-7d-8-1-A
20	Celebes	Tetep/IR2415-90-4-3-2//IR19661-131-1-2
21	Rojo Lele Delanggu	Delanggu, Klaten Central Java
22	Pendok	Tuban, East Java
23	Gilirang	B6672/ Membrano
24	Kurik Kusut Karanganyar	Karanganyar, Central Java
25	Genjah Arum	Banyuwangi, East Java

4) Quality Testing and DNA Quantity

DNA quality testing used electrophoresis method [14], prepared agarose gel 1% (w/v) by dissolving 0.25 g agarose into 25 ml TAE buffer on the Erlenmeyer tube and was heated on the microwave for 1 minute until the solution seemed clear, and then added with EtBr (Stadium Bromide) 1.5 µl. furthermore, the solution was poured into a mold equipped with a comb and was left to harden. Then, the print was put into an electrophoresis device containing the TAE buffer solution and was subsequently mixed with 5 µl of genomic DNA in 1 µl loading dye using the pipetting technique. After mixing it up, then it was put into agarose wells placed in an electrophoresis device. For

electrophoresis, it was set with a voltage of 75 Volts for 60 minutes. After the electrophoresis process was completed, and then the electrophoresis results were visualized using the Gel Document tool.

DNA Quantity Testing used a spectrophotometer. Before using the tool, it must be calibrated first, which was done by putting 400 μ l dd H₂O into the cuvette and pressing the blank button. Afterward, the 2 μ l DNA sample was put into the cuvette and 398 μ l dd H₂O was added. And the measurement of DNA concentration was at a wavelength of 260/ 280 nm. [21] At a wavelength of 260 nm, the RNA and DNA Ribbons can be absorbed optimally, and at a wavelength of 280 nm, protein can be absorbed. DNA purity will occur if the value is higher than 1.8, while the DNA indication is contaminated if the value is lower than 1.8.

5) PCR

The master mix used in the PCR reaction was Bioline with a total of 10 μ l tubes, where per tube contained 1 μ l DNA, 3 μ l ddH₂O, 5 μ l Master mix and 1 μ l primer and then running PCR with pre-denaturation at 95°C for 5 minutes, and then being denatured for 40 cycles for 30 seconds at 94°C. Annealing was conducted for 30 seconds at 30°C and extension was done for 30 seconds at 72°C. Final extension was done for 5 minutes at 72°C and soaked for 15 minutes at 16°C. [22]

6) Electrophoresis of PCR Results

The PCR amplification results from the targeted genes can be seen through DNA electrophoresis, by preparing 1 gram of agarose and were dissolved in TAE buffer, heated to clear and added with 2 μ l of ethidium bromide and was then poured into the mold. After solidifying, it was moved into the tank electrophoresis. Furthermore, DNA product of PCR results was inserted into the gel well, and electrophoresis ran for 40 minutes at 75V.

7) Primary and Phylogenetic Analysis

Analysis of PCR electrophoresis results began by marking DNA Ribbons using binary data, namely 1 for the DNA Ribbon that appeared and 0 for DNA Ribbon that did not appear at the same marker size position. It was then run using the NT SYS -pc, [23] resulting phylogenetic tree from PCR results. And the percentage of polymorphism allele on each primer can be calculated using the formula:

$$\% \text{ Polymorphism} = \frac{\text{Number of polymorphism alleles}}{\text{Total number of alleles}} \quad (1)$$

The information of a primary can be determined by calculating the PIC value (Polymorphic Information Content). From here on, the accuracy of genetic markers in kinship can be determined.

III. RESULT AND DISCUSSION

A. RAPD Analysis of Black and Aromatic Rice Samples

A total of 25 genotype samples consisting of 10 black rice genotypes and 14 aromatic rice samples and 1 non-aromatic

rice sample were analyzed using 20 RAPD primers with an average primary number 9.85 Table 2. OPA 6 and OPA 11 primers produce a number DNA bands that are the least compared to other primers and the highest number of DNA bands is produced from OPA 12 and OPA 19 primers with 14 bands of DNA each. The total number of alleles produced 202 with details of 180 alleles of polymorphism and 22 alleles of monomorphism. Some primers that produce 100% polymorphisms are OPA 04, OPA 07, OPA 11, OPA 12, OPA 16, OPA 17, OPA 19, and each produces polymorphisms 7,12,6,14,13,13,14 , 10. In Figure 2, OPA 16 is shown with 100% polymorphism and Figure 1, OPA 08 is 88.9% polymorphism, while for the lowest polymorphic is generated from OPA 15 primer with 4 band bands and only 2 bands of polymorphic bands.

The analysis using 20 RAPD primers were quite effective in amplifying the DNA of 10 black rice cultivars and 15 aromatic rice cultivars. It can be seen from the three replications of running PCR with the same primer showing the same results. The role of sample DNA purity of each cultivar will greatly affect the results of fragment analysis during separation with gel electrophoresis. DNA purity and diversity result from RAPD analysis will be more apparent when DNA isolation is done in large amount, so that at the time of extraction it does not affect the breakdown of the genomic DNA chain [24] And oftentimes, overly small DNA concentrations will produce a less clear DNA ribbon and DNA prints containing phenolic compounds and polysaccharides that can also affect the DNA ribbon results.

The results of each primer show different differences in the number of DNA ribbons, as shown in Table 2. In the primer OPA 08 produces a polymorphic band 8, and in the OPA 16 primer produces 13 polymorphic ribbons.

Reading results of polymorphism DNA ribbons were then converted into binary data, where if there was a DNA band, it would be marked 1 and if there was no DNA band then it will be marked with 0, and was then analyzed using NTSYSpc (Numerical Taxonomy and Multivariate Analysis System) program version 2.01.

The total number of polymorphic DNA ribbons produced from these 20 primers was 180 so that above average, each primer had at least 9 polymorphic ribbons with an average percentage of polymorphism of 89.2%.

Amplified DNA ribbons that are produced in PCR [25], both in terms of quantity and intensity, depending also on the primary ability to recognize the sequence of complementary DNA to the DNA mold. And for the distribution of primer attachment in the mold will bring up fragments in a large number of primers and a small number of fragments in other primers. And the selectivity of the [26] RAPD primer selection for use will also have an effect on polymorphism. This is caused by each primer having a different site attachment, which can also lead to a difference in polymorphic DNA ribbons. This difference also appears in the large number of base pairs & number of ribbons and their size as shown in Figure 1 OPA 8 primers and Figure 2 OPA 16 primers. The ribbons are scattered at different marker sizes ranging from 100 to 3000 bp.

TABLE II
 TWENTY AMPLIFIED RAPD PRIMERS AND POLYMORPHISMS IN 25 RICE CULTIVARS

No	Primary	Base Order (5' -3')	DNA Ribbon	Monomorphic amplicons	Polymorphic amplicons	(%) of polymorphism
1	OPA-01	CAGGCCCTTC	10	1	9	90
2	OPA-02	TGCCGAGCTG	7	1	6	85.7
3	OPA-03	AGTCAGCCAC	10	6	9	90
4	OPA-04	AATCGGGCTG	7	-	7	100
5	OPA-05	AGGGGTCTTG	10	1	9	90
6	OPA-07	GAAACGGGTG	12	-	12	100
7	OPA-08	GTGACGTAGG	9	1	8	88.9
8	OPA-09	GGGTAACGCC	11	1	10	90.9
9	OPA-10	GTGATCGCAG	9	4	5	55.6
10	OPA-11	CAATCGCCGT	6	-	6	100
11	OPA-12	TCGGCGATAG	14	-	14	100
12	OPA-13	CAGCACCCAC	8	2	6	75
13	OPA-14	TCTGTGCTGG	10	1	9	90
14	OPA-15	TTCCGAACCC	4	2	2	50
15	OPA-16	AGCCAGCGAA	13	-	13	100
16	OPA-17	GACCGCTTGT	13	-	13	100
17	OPA-18	AGGTGACCGT	13	1	12	92.3
18	OPA-19	CAAACGTCGG	14	-	14	100
19	OPA-20	GTTGCGATCC	10	-	10	100
20	OPC-07	GTCCCGACGA	7	1	6	85.7
Total			197	22	180	1784.1
Average			9.85	1,1	9	89.205



Fig 1. DNA ribbon profile of 10 black rice cultivars (1-10) and 15 aromatic rice cultivars (11-15) using RAPD OPA 08 primers with a 100-bp DNA marker size.




Fig 2. DNA band profiles of 10 black rice cultivars (1-10) and 15 aromatic rice cultivars (11-15) using RAPD OPA 16 primers with a 100-bp DNA marker size.

**TABLE III
SIMILARITY MATRIX**

	H.Banjarnegara	H.Purbalingga	Pariireng	H.Purwokerto	Harelahok	H.Bantul	H.Lumajang	H.Toraja	H.Melik	H.Blitar	Mentikwangi	MWB	M.05banjarnegara	Situpatenggang	PPM	Batanggadis	RPK	MSK	Situbagendit	Celebes	Rojoleledelelangu	Pendok	Gilirang	KKK	Genjaharum				
H.Banjarnegara	100																												
H.Purbalingga	81.97	100																											
Pariireng	75.43	85.56	100																										
H.Purwokerto	83.80	83.70	86.36	100																									
Harelahok	72.52	76.14	76.19	75.58	100																								
H.Bantul	80.93	83.15	83.53	87.36	74.70	100																							
H.Lumajang	87.50	81.77	79.77	88.14	76.92	90.06	100																						
H.Toraja	80.00	87.00	81.25	85.71	74.47	83.16	84.97	100																					
H.Melik	77.38	76.30	83.64	82.84	77.02	88.34	84.34	83.24	100																				
H.Blitar	75.61	72.19	75.78	77.58	71.34	77.99	82.72	79.56	85.71	100																			
Mentikwangi	57.75	58.33	57.61	59.57	55.56	56.04	58.38	58.82	59.89	57.80	100																		
MWB	53.26	53.97	57.46	55.14	54.24	53.63	57.14	55.72	56.32	56.47	78.76	100																	
M.05banjarnegara	58.95	58.46	59.89	59.69	59.02	59.46	61.70	61.84	63.33	60.23	81.41	88.78	100																
Situpatenggang	64.92	58.16	60.64	63.54	58.70	61.29	64.55	63.46	64.09	62.15	82.00	82.23	87.69	100															
PPM	50.00	47.27	44.59	47.21	50.98	46.45	46.84	45.20	46.67	46.58	66.27	65.06	67.44	68.21	100														
Batanggadis	55.32	55.96	55.14	56.09	56.35	52.46	56.99	54.63	53.93	55.17	82.23	78.35	82.00	78.61	65.88	100													
RPK	56.68	52.08	52.17	54.26	50.00	54.95	58.38	57.84	57.63	58.96	79.59	77.72	82.41	81.00	66.27	81.22	100												
MSK	57.14	53.61	54.84	57.90	54.95	55.44	57.75	56.31	58.10	54.86	83.84	80.00	84.58	84.16	64.33	82.41	84.85	100											
Situbagendit	54.14	54.84	56.18	56.04	54.02	56.82	56.98	55.56	59.65	57.49	76.84	79.14	80.83	77.32	63.80	77.49	82.11	84.38	100										
Celebes	56.00	57.78	59.30	57.96	59.52	57.65	60.12	57.29	59.39	58.39	77.17	74.03	78.08	76.60	58.60	80.00	73.91	76.34	80.90	100									
Rojoleledelelangu	55.62	56.25	55.44	57.45	56.67	56.04	56.22	54.90	56.50	54.34	74.49	70.47	74.37	75.00	68.64	73.10	71.43	74.75	76.84	72.83	100								
Pendok	56.32	51.40	49.12	53.71	57.49	53.25	54.65	48.17	51.22	50.00	66.67	67.78	66.67	70.59	71.80	66.30	72.13	72.43	74.58	66.67	79.78	100							
Gilirang	56.18	52.46	52.57	55.87	54.97	54.34	57.96	55.39	57.14	58.54	78.08	78.26	81.05	79.58	60.00	75.53	77.01	81.48	85.08	75.43	74.87	72.41	100						
KKK	57.78	55.14	55.37	56.35	54.34	54.86	58.43	54.82	55.29	57.83	75.13	75.27	78.13	76.68	58.03	74.74	76.19	77.49	80.87	74.58	75.13	71.59	90.00	100					
Genjaharum	54.34	55.06	55.29	56.32	61.45	54.76	56.14	52.63	57.67	55.35	70.33	71.51	73.51	70.97	64.52	66.67	69.23	71.74	77.27	69.41	79.12	76.92	77.46	77.71	100				




Fig 3. Phylogenetic relationship analysis of black and aromatic rice cultivars

B. Similarity and Phylogenetic Analysis

Dendograms of black and aromatic rice clusters can be obtained after knowing the genetic distance from the similarity values of 25 black and aromatic rice varieties, in Figure 3 it is clearly shown that the dendrogram obtained with two large clusters. Table 3 (in appendix) describes the similarity level between one variety and another, where a high degree of similarity can be used as information to select parent plant of crossbreeding so that the success of crossbreeding has a greater chance. If viewed from the two major groups, namely black rice and aromatic rice, Lumajang black rice, and Situpatenggang aromatic rice have a similarity level of 0.649%. With a high degree of similarity, the kinship between the two cultivars is also high.

The results of the phylogenetic kinship analysis of 10 black rice cultivars and 15 aromatic rice cultivars, at a coefficient of 0.84%, show high genetic variation as shown in Figure 3.

The phylogenetic tree produced shows that there were two major groups: 10 black rice cultivars (Herelahok, Hitam Blitar, Hitam Melik, Pari Ireng, Hitam Toraja, Hitam

Purbalingga, Hitam Lumajang, Hitam Bantul, Hitam Purwokerto, Hitam Banjarnegara) and 15 cultivars aromatic rice (PPM, PPM, Genjah Arum, Pondok, Rojolele Delanggu, Celebes, KKK, Gilirang, Situbagendit, Mentik Susu Karanganyar, Mentik Wangi Banjarnegara, RPK, Situpatenggang, Batang Gadis dan Mentik Wangi Karanganyar) are further divided into eight small groups, and the small group will still also be subdivided into small groups. This shows a kinship or genetic variation between the groups. The more groups being formed in phylogenetic trees, the larger diversity between cultivars, thus showing that it can be used as the first step for selecting cultivar collection to be made as parent plant in crossbreeding to obtain superior varieties.

IV. CONCLUSIONS

Twenty RAPD primers used to amplify 10 black rice cultivars and 15 aromatic rice cultivars were obtained. The total number of polymorphic DNA ribbons was 180; so that on average, in each primer, there were at least 9 polymorphic ribbons with an average polymorphism percentage of 89.2%.

The genetic diversity of all cultivars being analyzed was also seen to be quite high at 84% coefficient, where there were two major groups, i.e. black rice and aromatic rice, which was then divided into eight small groups. The results of the phylogenetic analysis demonstrate that the high diversity between black rice and aromatic rice can be made as a first step for selecting rice plants like the parent plant in crossbreeding.

ACKNOWLEDGMENT

This research is fully supported by using the scholarship funds of the Government of the Republic of Indonesia through the Indonesian Institute of Education Scholarship Fund Management Institution (LPDP BUDI-DN) Indonesia. Number: PRJ-4951/LPDP.3/2016.

REFERENCES

- [1] Chaudary, R.C. and D.V. Tran. 2001. Speciality Rices of the World: A Prologue. In *Speciality Rices of the World; Breeding, Production and Marketing*. Enfield, N.H. (USA) :Science Publishers. Inc. and FAO. p. 3-12.
- [2] Murdifin M, Pakki E, Rahim A, Syaiful SA, Ismail, EvaryYM and Baha, 2015. Physicochemical properties of Indonesian pigmented Rice (*Oryza sativa* Linn.)varieties from South Sulawesi. *Asian JPlant Sci*,14(2): 59-65.18.
- [3] Hu, C.; Kitts, D. D., 2001, Evaluation of antioxidant activity of epigallocatechin gallate in biphasic model systems in Vitro. *Mol. Cell. Biochem*, 218, 147-155.
- [4] Hu, C.; Zhang, Y.; Kitts, D. D.,2000, Evaluation of antioxidant and prooxidant activities of bamboo Phyllostachys nigra var. Henonisleaf extract in Vitro. *J. Agric. Food Chem*, 48, 3170 - 3176.
- [5] Tsuda, T.; Horio, F.; Osawa, T, 1999 . Absorption and metabolism of cyanidin-3-O- β -glucoside in rats. *FEBS Lett.*, 449, 179-182.
- [6] Oikawa T, Maeda H, Oguchi T, Yamaguchi T, Tanabe N, Ebana K, Yano M, Ebitani T, and Izawa T, 2015 .The birth of a black rice gene and its local spreadby introgression. *Plant Cell* ,27(9): 2401-14
- [7] Hu, C.; Kitts, D. D, 2001. Evaluation of antioxidant activity of epigallocatechin gallate in biphasic model systems in Vitro. *Mol. Cell. Biochem*, 218, 147-155.
- [8] Lee, J. C., Kim, J. D., Hsieh, F. H., & Eun, J. B. 2008. Production of black rice cake using ground black rice and medium - grain brown rice. *International journal of food science & technology*, 43(6), 1078-1082
- [9] Pratiwi, R., Tunjung, W. A. S., Rumiati, R., & Amalia, A. R. 2016. Black Rice Bran Extracts and Fractions Containing Cyanidin 3-glucoside and Peonidin 3-glucoside Induce Apoptosis in Human Cervical Cancer Cells. *Indonesian Journal of Biotechnology*, 20(1), 69-76.
- [10] Sui, X., Zhang, Y., & Zhou, W. 2016. Bread fortified with anthocyanin-rich extract from black rice as nutraceutical sources: Its quality attributes and in vitro digestibility. *Food chemistry*, 196, 910-916.
- [11] Buttery, R.G., Juliano, B.O., Ling, L.C. 1983. Identification of rice aroma compound 2-acetyl-1-pyrroline in pandan leaves. *Chemical Industries (London)* 20: 478.
- [12] Buttery, R.G., Ling, L.C., Juliano, 1982. B.O. 2- Acetyl-1-pyrroline: an important aroma component of cooked rice. *Chemical Industries (London)* 19: 958.
- [13] Chen, S., Yang, Y., Shi, W., Ji, Q., He, F., Zhang, Z., Cheng, Z., Liu, X., Xua, M. Badh2, 2008. encoding betaine aldehyde dehydrogenase, inhibits the biosynthesis of 2-acetyl-1-pyrroline, a major component in rice fragrance. *Plant Cell*.
- [14] Rokas, Antonis, 2011 phylogenetic Analysis of Proten Sequence Data Using the Randomized Axelerated Maximum Likelihood (RAXML) Program. Wiley Online Library. UNIT 19.11
- [15] Arif S. A., Ma, N.L., Mahmoud, A, Rahmat, Z., 2016. Molekuler Phylogeny and Structure Prediction Of Rice RFTI Protein. *Jurnal Tekhnologi*. Vol. 78 (2)
- [16] Jonah, P. M. Bello, L. Lucky, O., Midau A., Moruppa, S.M. 2011, Review The Importance of Molecular Markers in Plant Breeding Programmes *International Research Journal* Vol. 11 (5)
- [17] Sing, R., K.. 2000. *Aromatic Rices.*, Calcuta: Oxford & IBH Publishing Co. Pvt.
- [18] Horrike Tokumasa, 2016. An Introduction To Molecular Phylogenetic Analysis *Reviews In Agriculture Science*, Vol. 4, 34 - 45.
- [19] Abbas, Y. A dan Radhi, G. F, 2017, Genetic Variation of Enterobacter coaceae Using Random Amplified Polymorphic DNA RAPD-PCR Technique, *Donnish Journal of Microbiology and Biotechnology Research* Vol 4 (3) pp. 010-123.
- [20] Doyle, JJ and Doyle, J, L, 1987, A Rapid DNA Isolation Prosedure For Small Quantities of Fresh leaf Tissue ,*Phytochemical Bulletin*, Vol 19, no. 1, p. 11-15
- [21] Mordechai, E, 1999. Application of PCR The Methodologies in Molecular Diagnostic, Burlington Country, USA
- [22] Saiki R.K, 1989, The Desain and Optimization of the PCR, Pages: 7-16 in Erlich, H.A (ed) *PCR Technology. Principle and Appplication for DNA Amplification*.
- [23] Rohlf, F J. 2000, *NT SYS-pc: Numerical Taxonomy and Multivariate Analysis System Version 2.1 User Guide*. Departemen Of Ecology and Evaluation State Unversyty of New York
- [24] Widden B and Svvenson L, 1992 Conservation of Genetic Variation in Plant- The Importan of Population Size and Gene Flow. In: *Ecological Principle Of Nature Conservation*. Hansson L (ed) Elsevier Applied Science, London, pp 113 – 161
- [25] Tingey, S.V., JA Rafalski, and M. K. Hanafey, 1994. Genetic Analysis with RAPD marker In: Corruzi, C and P Puidormenech (eds) *Plant Molecular Biology Beli*: Springer- Verlag.
- [26] Vi, K. 2017 Using RAPD PCR Method for Studying DNA Polymorphism of Agriculture Importance Arthropods. *Agriculture Research And Technology*. Vol 9 (4).