

7.1.3

“CROSSING BOUNDARIES TOV. ARE  
Digital Repository Univ MODERN AGRICULTURE :  
Science and Technology, Economic  
and Social Welfare in Developing  
Countries”

# INTERNATIONAL CONFERENCE ON AGRICULTURE

Proceedings

International  
Conference  
on  
Agriculture  
**I C A** 2013

*Surabaya, October 10-11, 2013*

Published by UPN "Veteran" JAWA TIMUR  
ISBN: 978-602-9372-57-1





23	Farmers' Adaptation Strategy on Climate Change Impacts and Technical Efficiency of Rainfed Lowland in Organic Rice in Tanggamus Regency, Lampung Province <i>Ktut Murniati, Jangkung Handoyo Mulyo, Irfham, and Slamet Hartono</i>	213
24	Cocoa Farming and Analysis of Economic Community Farmers State in East Java <i>Pawana Nur Indah</i>	224
26	Market Institutional Roles on Rice Farmer's Income <i>Raharto, S.</i>	235
27	Contract Farming Syariah as a Social Business Model: Lesson From Local Wisdom Experience <i>Sumartono</i>	250
28	Relative Economy Efficiency of Coffee Farmer in Belantih Village Kintamani Sub-District Bangli Regency <i>Hariyati, Y.</i>	260
29	Analysis of Coffee Secondary Agro-Industry as Superior Products in Bondowoso Regency <i>Mustapit and Evita Soliha Hani</i>	274
30	The Effectiveness of Gene Transformation Sosps 1 Using Plasmid Psmab-Sosps 1 Construct in Sugarcane ( <i>Saccharum Officinarum</i> L.) <i>Parawita Dewanti, Pumama Okviandari, Yunianzi Tiara Prima and Bambang Sugiharto</i>	281
31	Analysis Of Factors Influencing Beef Supply Chain in Jember District <i>Joni Murti Mulyo Aji, Titin Agustiana and Annona Emhar</i>	291
32	Agriculture Sector Human Resource Overlook in Indonesia (Overview of The Agricultural Sector Human Resource Alignment in Merauke – Papua) <i>Aldon Sinaga, Umi Rofiatin, Asnah, Ana Arifatus S. and Taufik Iskandar</i>	313
33	The Involvement of The Institution on The Dynamics of Revenue Resources Based on P3A Farmers Members Opinion in Psetk Activity (A Survey On Citaal Irrigation Regency Kecamatan Ciwaru Kabupaten Kuningan) <i>Dedi Sufyadi</i>	332

**THE EFFECTIVENES OF GENE TRANSFORMATION SoSPS1  
USING PLASMID pSMAB-SoSPS1 CONSTRUCT  
IN SUGARCANE (*Saccharum officinarum* L.)**

Parawita Dewanti,<sup>1,3)</sup> Purnama Okviandari<sup>2,3)</sup>, Yunianzi Tiara Prima<sup>2,3)</sup> dan  
Bambang Sugiharto<sup>2,3)</sup>

<sup>1)</sup> Faculty of Agriculture University of Jember, <sup>2)</sup> Faculty of Mathematics and Natural Sciences  
University of Jember, <sup>3)</sup> CDAST (Center for Development Advance of Sciences Technology)  
University of Jember

e-mail : [parawita@yahoo.co.id](mailto:parawita@yahoo.co.id), Hp 08123463272

### ABSTRACT

SoSPS1 gene is a gene isolated from sugarcane plants. This gene plays an important role in the biosynthesis of sucrose and sucrose accumulation-effect on the crop. Efforts to increase the sucrose content of the sugarcane can be done by using the transformation SoSPS1 gene of plasmid constructs pSMAB-SoSPS1 through *Agrobacterium tumefaciens* vector. This study aims to determine the effectiveness of SoSPS1 gene transformation using plasmid constructs pSMAB-SoSPS1 on sugarcane (*Saccharum officinarum* L.). The results showed that the effectiveness of SoSPS1 gene transformation using plasmid constructs pSMAB-SoSPS1 about 2% and PCR analysis amplified by the size of 462 bp.

Keywords: Gen SoSPS1, pSMAB-SoSPS1, gene transformation, Sugarcane (*Saccharum officinarum* L.)

### INTRODUCTION

Sugarcane (*Saccharum officinarum* L.) is a major sugar-producing plant in Indonesia, but sugar production is still insufficient national. Based on data from Market & Trade Sugar Word 2004 / 2005, imports of sugar in the last 20 years to more than 50%. The development of techniques gene insertion of genetic material such as SPS, is expected to produce new varieties with high productivity and yield in a relatively quick time.

Sucrose phosphate synthase (SPS) plays an important role in the biosynthesis of sucrose and the affect to increased sucrose content (Huber and Huber, 1996; Laporte *et al.*, 2001). Several studies have shown that overexpression of SPS gene can increase the sucrose content in tomatoes (Worrell *et al.*, 1991; Nguyen-quoc *et al.*, 1999), *Arabidopsis thaliana* (Signora *et al.*, 1998), tobacco (Miswar *et al.*, 2005) and sugarcane (Miswar *et al.*, 2007), but the results still vary.

Some factors influencing the success of insertion of genetic material, among others, is a binary vector (An *et al.*, 1986), varieties (Ling *et al.*, 1998; Ellul *et al.*, 2003; Santoso *et al.*, 2009), explant type (Fillati *et al.*, 1987), plant growth



regulators (Pfitzner *et al.*, 1998; Cortina *et al.*, 2004), the concentration of bacteria (Shahin *et al.*, 1986), *Agrobacterium* strains (Roekel *et al.*, 1993), concentration antibiotics (Hu and Phillips, 2001; Qiu *et al.*, 2007) and promoter (Liu *et al.*, 2003; Yoo *et al.*, 2005).

Promoter is a DNA regulator that aims to regulate gene expression patterns so that it can be seen that the presence and expression of the gene in plants. Promoter CaMV (*cauliflower mosaic virus*) are often used in plant transformation, but if used in the transformation of sugarcane (monocotyl) is low success rate (Chowdhury and Vasil., 1992). Promoter RUBQ2 (*rice polyubiquitin*) is a new constitutive promoter in rice plants and monocotyl plant groups (Wang and Oard, 2003), but for the transformation of the SPS gene in sugarcane effectiveness to be investigated.

Plasmid Constructs pSMAB-SoSPS1 (Sugiharto, 2009) which is controlled by the promoter RUBQ2, never been tested in sugarcane. Therefore, genetic transformation using plasmid constructs pSMAB-SoSPS1 controlled by promoter RUBQ2 with lateral buds as explants needs to be done. According Manickavasagam *et al.* (2004), the use of lateral buds as explants to avoid somaclonal variation, faster regeneration time and can increase 50% the transformation efficiency.

## EXPERIMENTAL DETAILS

### Plant Material

Explants from axillary buds of sugarcane var. BL soaked in alcohol 70% for 5 seconds, then in 0.05% HgCl<sub>2</sub> solution for 60 seconds, rinsed 3 times with sterile distilled water and drained on a filter paper for 30 minutes. Explants grown on MS medium (Murashige and Skoog + 1.5 mg l<sup>-1</sup> BAP and 0.1 mg l<sup>-1</sup> GA<sub>3</sub> at 25°C under 16-h and 8-h cycle with a light intensity of 1600 lux illumination. Explant were sub cultured every 2 weeks and form plantlets for 10 weeks. Basal part of plantlet is used as a source for transformation.

### *Agrobacterium* strain and plasmid vector

*Agrobacterium tumefaciens* strain LBA4404, harboring plasmid pSMAB, which contained bar gene as selectable marker, and *SoSPS1* gene construct, was utilized for *Agrobacterium*-mediated transformation. In this construct, the expression of *phosphinotricyn* gene was under the control of the nopaline synthase promoter (pNOS), pArbcs terminator and *SoSPS1* gene was driven by RUBQ2 (Figure 1).




Figure 1. Map of pSMAB-SoSPS1 construct. cDNA-SoSPS1 under the control of RUBQ2 promoter and resistance gene (*bar*) for phosphinotricyn (Sugiharto *et al.*, 2009).

#### ***Agrobacterium tumefaciens* preparation for Transformation**

Single colonies of *Agrobacterium tumefaciens* LBA4404, harboring plasmid pSMAB-SoSPS1 on Luria Bertani (LB) medium plate (10 g<sup>l</sup><sup>-1</sup> pepton, 10 g<sup>l</sup><sup>-1</sup> yeast extract, 5 g<sup>l</sup><sup>-1</sup> NaCl, 14 g<sup>l</sup><sup>-1</sup> agar, pH 7.0) were inoculated in 2 ml liquid Yeast Extract Peptone (YEP) medium supplemented with 50mg<sup>l</sup><sup>-1</sup> kanamycin, 30 mg<sup>l</sup><sup>-1</sup> Streptomycin and 50 mg<sup>l</sup><sup>-1</sup> rifamycin and incubated overnight at 28°C with continuous shaking at 150 rpm. *Agrobacterium* resubcultured into 50 ml liquid YEP medium containing 50 mg<sup>l</sup><sup>-1</sup>kanamycin, 30 mg<sup>l</sup><sup>-1</sup> Streptomycin and 50 mg<sup>l</sup><sup>-1</sup> rifamycin and incubated 4-h at 28°C to a final density of 0.3-0.4 OD<sub>600</sub>. *Agrobacterium* cells were harvested by centrifugation at 5.000 rpm for 10 min and resuspended in liquid MS medium.

#### **Transformation mediated by *Agrobacterium tumefaciens*.**

The explants were infected by immersing in *Agrobacterium* suspension for 30 min with gentle shaking three to five times during the infection process. Subsequently, the infected explants were dried on a sterile filter paper and transferred onto co cultivation medium (C) for 2-d in dark at 25°C. After co cultivation, the infected explants were washed with 50 ml of liquid MS containing cefotaxime 500 mg<sup>l</sup><sup>-1</sup>, drained on sterile filter paper and transferred onto Elimination medium (E) at 25°C under 16-h light, 8-h dark cycle for 5 days. Explants were cultured subsequently on Selection medium (S) 2 weeks for shoot induction with 3 times subculture, elongation medium (L) 2 weeks for shoot enlargement and rooting



medium (R) 2 weeks for rooting induction. Medium for transformation shown in Table 1.

Table 1. Kinds of medium are used for the process of transformation

Chemical Material	Co cultivation Medium (C)	Elimination Medium (E)	Selection Medium (S)	Elongation Medium (L)	Rooting Medium (R)
MS salts	√	√	√	√	√
IAA ( $\text{mg l}^{-1}$ )	0.2	0.2	0.2	-	-
BAP ( $\text{mg l}^{-1}$ )	2	2	2	0.25	-
GA3 ( $\text{mg l}^{-1}$ )	-	-	-	0.25	-
NAA ( $\text{mg l}^{-1}$ )	-	-	-	-	0.15
Acetosyringone ( $\text{mg l}^{-1}$ )	50	-	-	-	-
Cefotaxim ( $\text{mg l}^{-1}$ )	-	500	500	500	500
Phosphinotricyn ( $\text{mg l}^{-1}$ )	-	-	5	5	5
Phytigel ( $\text{g l}^{-1}$ )	3	3	3	3	3
Sucrose ( $\text{g l}^{-1}$ )	30	30	30	30	30
Ph	5.8	5.8	5.8	5.8	5.8

#### DNA isolation and PCR analysis

Genomic DNA was isolated from non-transformed (control) and transformed leaves. PCR analysis were carried out to detect the presence of the *bar* genes, using primer bar-F/R, 462 bp: 5-ATC GTC AAC CAC TAC ATC GAG AC-3 dan 5-CCA GCT GCC AGA AAC CCA CGT C-3. Linkage analysis of these genes in transgenic plants was conducted using PCR amplification.

## RESULTS AND DISCUSSION

Sugarcane var. BL has superior properties, among others: pest-resistant, high productivity and easily cultured (P3GI, 2009). Explant used is the lateral buds of sugarcane as shown in (Figure 2). The use of lateral buds explants have several advantages such as rapid regeneration and can produce a number of plants in a relatively short time (Manickavasagam *et al.*, 2004). Explants grown on MS medium containing  $1.5 \text{ mg l}^{-1}$  BAP and  $0.1 \text{ mg l}^{-1}$  GA<sub>3</sub> (L Medium). The addition of BAP can induce shoot formation (Lee, 1987), while GA<sub>3</sub> used for stem elongation. Husnah (2007) reported that, a concentration of  $0.1 \text{ mg l}^{-1}$  GA<sub>3</sub> provide optimal effect on the extension of the sugarcane stem in vitro. Figure 2 is an explant preparation process to form sugarcane plantlets. Plantlets age of 10 weeks (Figure 2E) ready to be used for the transformation process.



Figure 2. Culture of sugarcane lateral shoots. Lateral shoots of sugarcane (A), lateral bud explant isolation (B), the growth of shoots (C), 8 weeks (D), 10 weeks (E).

SoSPS1 gene transformation in sugarcane explants with *A. tumefaciens* lateral buds have several stages of co-cultivation, elimination and selection. The result of the transformation that has been done using pSMAB-SoSPS1 shown in Figure 2). In Table 2 it can be seen the number of explants of co-cultivation and the elimination of 50 explants. At the time of selection 1 (Figure 2 C) decrease the number of explants that are resistant to the selection agent. Explants were resistant to the selection agent for 31 explants. End of Selection 2 (Figure 2 D), from 31 explants who passed 1, only 23 explants that pass the selection. At the time of selection 3 (Figure 2 E), a putative transformed plant obtained a total of 18 plants. The results of the final stage of the selection of 18 plants indicated on the plant



resistance *phosphinotricyn* (PPT) media (S Medium ). Miki and McHugh, 2003 states transformed plants will have a tolerance to PPT, PPT is an inhibitor of glutamine synthetase. Glutamine synthetase is the only enzyme that can catalyze the assimilation of ammonia to glutamic acid. Therefore, the existence of this inhibition in the end the accumulation of ammonia which can lead to cell death (Toki et al., 1992). It is characterized by non-transformed plants will experience death (Song and Douches, 2009).




Figure 3. Growth of explants on co-cultivation medium (100 mgL<sup>-1</sup> acetosyringone) (A), media eliminations (500 mgL<sup>-1</sup> cefotaxime) (B), media selection (500mgL<sup>-1</sup> cefotaxime 5 mgL<sup>-1</sup> PPT) 21 days (C), 42 days (D) and 63 days (E).

After the end of the third selection, putative transformed plants moved into MS0 media with the addition of 500 mgL<sup>-1</sup> cefotaxim and 0.5 mgL<sup>-1</sup> Naphtalene cefotaxime Acetic Acid (NAA) which serves to induce root. After the complete plant morphology, plantlets acclimatized to genomic DNA isolated from the leaves of plants.





resistance *phosphinotricyn* (PPT) media (S Medium ). Miki and McHugh, 2003 states transformed plants will have a tolerance to PPT, PPT is an inhibitor of glutamine synthetase. Glutamine synthetase is the only enzyme that can catalyze the assimilation of ammonia to glutamic acid. Therefore, the existence of this inhibition in the end the accumulation of ammonia which can lead to cell death (Toki et al., 1992). It is characterized by non-transformed plants will experience death (Song and Douches, 2009).




Figure 3. Growth of explants on co-cultivation medium (100 mgL<sup>-1</sup> acetosyringone) (A), media eliminations (500 mgL<sup>-1</sup> cefotaxime) (B), media selection (500mgL<sup>-1</sup> cefotaxime 5 mgL<sup>-1</sup> PPT) 21 days (C), 42 days (D) and 63 days (E).

After the end of the third selection, putative transformed plants moved into MS0 media with the addition of 500 mgL<sup>-1</sup> cefotaxim and 0.5 mgL<sup>-1</sup> Naphtalene cefotaxime Acetic Acid (NAA) which serves to induce root. After the complete plant morphology, plantlets acclimatized to genomic DNA isolated from the leaves of plants.

Table 2. number of putative transformed plant in co cultivation, selection and rooting medium

	Number of explant
Cocultivation	50 (100%)
Elimination	50 (100%)
Selection 1	31 (62%)
Selection 2	23 (46%)
Selection 3	18 (36%)
Rooting	18 (36%)
Acclimatization	12 (24%)
Effectiveness Transformation	1 (2%)

The success of acclimatization by 70% because of the 18 plants were successfully acclimatized only 12 plants that can be isolated DNA genome. PCR is used to see the successful integration of the target gene into the plant genome SoSPS1. In (Figure 3) can be seen the results of electrophoresis of genomic DNA are amplified. Of the 12 putative transformed plants that are resistant to PPT analyzed using PCR, only 1 positive transformed plants. Amplified DNA indicated the presence of DNA bands or band size 462 bp, the presence of DNA bands were seen indicates that SoSPS1 gene has been integrated into the plant genome. While 11 other plants are not a band that signifies not the integration of T-DNA into the plant genome.



Figure 3. Analysis PCR of plant genomic results with primer F / R. K +: pSMAB plasmid-SoSPS1, K-: control plants (wild type), M: 1 kb DNA ladder, E1-12: plant 1-12.



Percentage of successful transformation obtained by 2% (1 of 50 explant transformed plant) indicates the efficiency is still low compared to the use of plasmids and different promoters, because SoSPS1 gene transformation research has been done using the plasmid with the promoter CaMV 35S pKYS (Miswar *et al.* 2007) and plasmid pCI4 with promoter RUBQ2 (Baskoro, 2012) demonstrated the success of 4% and 6%.

## CONCLUSION

The results showed that the effectiveness of SoSPS1 gene transformation using plasmid constructs pSMAB-SoSPS1 about 2% and PCR analysis amplified by the size of 462 bp.

## ACKNOWLEDGEMENTS

We are thankful to Prof. Dr. Ir. Bambang Sugiharto, M.Sc., Research Center for Molecular Biology University of Jember and research grant from 'Hibah Kompetensi' for providing the necessary facilities.


## REFERENCES

- An, G. 1985. High efficiency transformation of cultured tobacco cells. *Plant Physiol.* 79:568-570
- Ellul P, Garcia-Sogo B, Pineda B, Rios G, Roig LA & Moreno V. 2003. The ploidy level of transgenic plants in *Agrobacterium* mediated transformation of tomato cotyledons (*Lycopersicon esculentum* L. Mill.) is genotype and procedure dependent. *Theor. Appl. Gen.* 106 (2): 231-238
- Huber, S.C. and J.L. Huber. 1996. Role and regulation of sucrose-phosphate synthase in higher plant. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47 : 431 - 444
- Husnah, R. 2007. Mikropropagasi Tanaman *in Vitro* Menggunakan *Axillary Buds* Pada Tanaman Tebu. Tidak Diterbitkan. Skripsi. Jember. Universitas Jember.
- HU W. and G. C. PHILLIPS. 2001. A combination of overgrowth-control antibiotics improves *Agrobacterium tumefaciens*-mediated transformation efficiency for cultivated tomato (*L. esculentum*). *In Vitro Cell. Dev. Biol. Plant* 37:12-18
- Lee, T.S.G. 1987. Micropropagation of Sugarcane (*Saccharum spp.*). *Plant cell, Tissue, Organ and Culture*, 10: 47-55.

- Nguyen-Quoc, B., N'Tchobô, H., Foyer, C.H and Yelle, S. 1999. Overexpression of sucrose phosphate synthase increases sucrose unloading in transformed tomato fruit. *Journal of Experimental Botany*, **50**: 785-791.
- Manickavasagam, M., Ganapathi, A., Anbazhagan, V.R., Sudhakar, B., Selvaraj, N., Vasudevan, A. and Kasthuriangan, S. 2004. *Agrobacterium*-mediated Genetic Transformation and Development of Herbicide-resistant Sugarcane (*Saccharum* species hybrid) Using Axillary Buds. *Plant Cell*, **23**: 134-143.
- Miki, B and McHugh, S. 2003. Selectable Marker Genes in Transgenic Plants: Application, alternatives and biosafety. *Journal of Biotechnology*, **107**: 193-232.
- Miswar., Sugiharto, B., Soedarsono, J. dan Moeljopawiro, S. 2005. Transformasi Gen Sucrose Phosphate Synthase (SoSPS1) Tebu (*Saccharum officinarum* L.) Untuk Meningkatkan Sintesis Sukrosa pada Daun Tembakau (*Nicotiana tabacum* L.). *Biologi*, **4**: 337 - 347.
- Miswar., Sugiharto, B., Soedarsono, J., dan Moeljapawiro, S. 2007. Transformasi Gen Sucrose Phosphate Synthase (SoSPS1) Menggunakan *Agrobacterium tumefaciens* untuk Meningkatkan Sintesis Sukrosa pada Tanaman Tebu (*Saccharum officinarum* L.) *Berk. Penel. Hayati*, **12**: 137-143.
- Qiu D., G. Diretto, R. Tavarza, G. Giuliano. 2007. Improved protocol for *Agrobacterium* mediated transformation of tomato and production of transgenic plants containing carotenoid biosynthetic gene CsZCD. *Scientia Horticulturae* **112** : 172-175.
- Roekel J. S. C. van, B. Damm, L. S. Melchers, and A. Hoekema. 1993. Factors influencing transformation frequency of tomato (*Lycopersicon esculentum*). *Plant Cell Reports* **12**:644-647
- Santoso T.J., A. Sisharmini dan M. Herman.2009. Respon Regenerasi Beberapa Genotipe Dan Studi Transformasi Genetik Tomat (*Lycopersicon esculentum* Mill.) Melalui Vektor *Agrobacterium tumefaciens*. Balai Besar Penelitian dan Pengembangan Bioteknologi dan Sumberdaya Genetik Pertanian.
- Signora, L., Galtier, N., Skot, L., Lucas, H and Foyer, C.H. 1998. Over-expression of sucrose phosphate synthase in *Arabidopsis thaliana* results in increased foliar sucrose/starch ratios and favours decreased foliar carbohydrate accumulation in plants after prolonged growth with CO<sub>2</sub> enrichment. *Journal of Experimental Botany*, **49**: 669-680.
- Song, M and Douches, D. 2009. Selectable Marker & Marker for Screening. [http://www.msu.edu/course/css/451/Lecture/WK3-PT-SMG %2822009 %29.pdf](http://www.msu.edu/course/css/451/Lecture/WK3-PT-SMG%2822009%29.pdf). Diakses tanggal 3 Juli 2011.



- Sugiharto, B., Slameto dan Dewanti, P. 2009. Peningkatan Produksi Gula Melalui Overekspresi Gen untuk Sucrose-phosphate synthase dan Sucrose Transporter Protein pada Tanaman Tebu. Laporan Penelitian Hibah Kompetensi tahun 2009. Tidak dipublikasikan. Direktorat Jendral Pendidikan Tinggi Departemen Pendidikan Nasional.
- Toki, S., Takamatsu, S., Nojiri, C., Ooba, S., Anzai, H., Iwata, M., Christensen, A.H., Quail, P.H and Uchimiya, H. 1992. Expression of Maize Ubiquitin Gen Promoter-bar Chimeric Gene in Transgenic Rice Plants. *Plant Physiol*, **100**: 1503-1507.
- Worrell, A.C., Bruneau, J-W., Summerfelt, K., Boersig, M and Voelker, T.A. 1991. Expression of a Maize Sucrose Phosphate Synthase in Tomato Alters Leaf Carbohydrate Partitioning. *The Plant Cell*, **3**: 1121-1130.



# CERTIFICATE of APPRECIATION

This certificate is awarded to

*Dr.Ir. Parawita Dewanti, MP*

for participating in **International Conference on Agriculture - ICA 2013**  
University of Pembangunan Nasional "Veteran" Jawa Timur, Surabaya - Indonesia  
October 10-11, 2013

as

**PRESENTER**

title of paper

*The Effectiveness of Gene Transformation SoSPS1 Using Plasmid  
pSMAB-SoSPS1 Construct in Sugarcane (Saccharum officinarum L.)*

Chairman



Dr.Ir. Endang Yektiningsih, M.P.

Rector



Prof. Dr. Ir. Teguh Soedarto, M.P.

