Sequencing, genome analysis and ber host range of a novel Ralstonia phage, RsoP1EGY, isolated in Egypt

Abdelmonim Ali Ahmad, Kamel M. Elhalag, Hardian Susilo Addy, Mohamed A. Nasr-Eldin, Ahmed S. Hussien & Qi Huang

Archives of Virology Official Journal of the Virology Division of the International Union of Microbiological Societies

ISSN 0304-8608

Arch Virol DOI 10.1007/s00705-018-3844-4

ARCHIVES OF VIROLOGY

Official Journal of the Virology Division of the International Union of Microbiological Societies





Digital Repository Universitas Jember Authors & Editors • My account



Search **Q**

Menu 🕶

You're seeing our new journal sites and we'd like your opinion, please send feedback



Archives of Virology

Official Journal of the Virology Division of the International Union of **Microbiological Societies**



Journal home , Digital Repository Universitas Jember

Editors

Editor-in-Chief

Tim Skern, Vienna, Austria Fields of interest: RNA viruses, structure of viruses and viral proteins, virus-host interactions Reviews in medical virology ORCID iD: 0000-0001-8865-5500

Editorial Board

Patricia V. Aguilar, Galveston, TX, USA Fields of interest: bunyavirus, emerging virus diseases, dengue virus, vector-borne viruses, flaviviruses, influenza, orthomyxoviruses, Togaviruses, virus-host cell interaction

Graciela Andrei, Leuven, Belgium Fields of interest: Animal Virus Pathogenesis, Animal Virus-Host Cell Interactions, Cytomegalovirus, Epstein Barr virus, Herpesviruses, Large DNA viruses, Papillomaviruses, Polyomavirus, Poxviruses, Varicella-Zoster Virus, Virus-Host Cell Interactions

Tatjana Avšič-Županc<mark>, Ljubljana</mark>, Slovenia Fields of interest: Bunyaviruses, Flaviviruses, Zoonoses

Ana C. Bratanich, Buenos Aires, Argentina

Fields of interest: Circoviruses, Diagnostics, Herpesviruses, Molecular Biologyof Animal Viruses, Paramyxoviruses, Parvoviruses, Picornaviruses, Veterinary viruses, Viral molecular pathogenesis, Viral pathogenesis, Virus Vectors, Virus-Host Cell Interactions

Michael Carpenter, Winnipeg, MB, Canada Fields of interest: diagnostics of human/animal viruses, hepatitis viruses, molecular biology of animal viruses, virus-host cell interactions

ERS

Publish with us

☑ Springer

Authors & Editors Journal authors Publishing ethics Open Access & Springer

Discover content

SpringerLink Books A-Z Journals A-Z Video Other services Instructors Librarians (Springer Nature)

Societies and Publishing Partners Advertisers Shop on Springer.com

About Springer

About us Help & support Contact us Press releases Impressum

Legal

General term & conditions Rights & permissions Privacy

How we use cookies

Skip to main content

🙆 Springer Link

Archives of Virology All Volumes & Issues

Special Topic-Section to honor Marc van Regenmortel (Pages 2017 - 2054)

ISSN: 0304-8608 (Print) 1432-8798 (Online)

In this issue (37 articles)

1. In honor of Marc van Regenmortel - Editorial

Editorial

Tim Skern Pages 2017-2018

2. In honor of Marc van Regenmortel

Virus taxonomy—a taxing task

Ulrich Desselberger Pages 2019-2020

3. In honor of Marc van Regenmortel

MARC VAN REGENMORTEL: a personal reminiscence

<u>Donald A. Hendry</u> Pages 2021-2022 4. In honor of Marc van Regenmortel

Marc H. van Regenmortel

Dion du Plessis Pages 2023-2024

5. In honor of Marc van Regenmortel

A 30-year journey of trial and error towards a tolerogenic AIDS vaccine

<u>Jean-Marie Andrieu, Wei Lu Pages 2025-2031</u>

6. In honor of Marc van R<mark>egenmortel</mark>

Viruses do not h<mark>ave polythetic properties; spe</mark>cies are polythe<mark>tic classes and do no</mark>t have any properties

Charles H. Calisher Pages 2033-2036

7. In honor of Marc van Regenmortel

<u>Virus classificat<mark>ion – where do you dra</mark>w the line?</mark></u>

Peter Simmonds, Pakorn Aiewsakun Pages 2037-2046

8. In honor of Marc van R<mark>egenmortel</mark>

Viral species, vira<mark>l genomes and HIV vacc</mark>ine design: is the ratio<mark>nal design of biological complexity a</mark> utopia?

Marc H. V. van Regenmortel Pages 2047-2054

9. Original Article

Host serum microRNA profiling during the early stage of foot-and-mouth disease virus infection

Suresh H. Basagoudanavar, Madhusudan Hosamani, R. P. Tamil Selvan... Pages 2055-2063

10. Original Article

PCR array profiling of antiv<mark>iral genes in human embryonic kid</mark>ney cells expressing human coronavirus OC43 structural and

accessory proteins

Meshal Beidas, Wassim Chehadeh Pages 2065-2072

11. Original Article

Inactivated enterovirus 71 with poly-y-glutamic acid/Chitosan nano particles (PC NPs) induces high cellular and humoral immune responses in BALB/c mice

Prabuddha S. Pathinayake, W. A. Gayan Chathuranga, Hyun-Cheol Lee... Pages 2073-2083

12. Original Article

Bursal immunopathology responses of specific-pathogen-free chickens and red jungle fowl infected with very virulent infectious

bursal disease virus

Mohd Isa Farhanah, Abdul Rahaman Yasmin, Nguyen Phuc Khanh... Pages 2085-2097

13. Original Article

Functional growth inhibition of influenza A and B viruses by liquid and powder components of leaves from the subtropical plant Melia azedarach L.

Kuniaki Nerome, Kazufumi Shimizu, Shiori Zukeran, Yasuhiro Igarashi... Pages 2099-2109

14. Original Article

Surveillance of HIV-1 drug resistance in Xinjiang: high prevalence of K103N in treatment-naïve individuals

Zhanjun Shu, Yuying Chen, Aihemaiti Abudureyimu, Tingting Li... Pages 2111-2119

15. Original Article

Antiviral activity of Embelia ribes Burm. f. against influenza virus in vitro

Md Shahadat Hossan, Ayesha Fatima, Mohammed Rahmatullah... Pages 2121-2131

16. Original Article

A multiplex PCR method for the simultaneous detection of three viruses associated with canine viral enteric infections

Xiaoyu Deng, Jiali Zhang, Jiazi Su, Hao Liu, Yanlong Cong... Pages 2133-2138

17. Original Article

Comparative genomics of Cp8viruses with special reference to Campylobacter phage vB_CjeM_los1, isolated from a slaughterhouse in

Ireland

Lisa O'Sullivan, Alan Lucid, Horst Neve, Charles M. A. P. Franz... Pages 2139-2154

18. Original Article

Different susceptibility to porcine reproductive and respiratory syndrome virus infection among Chinese native pig breeds

Chunhua Meng, Lei Su, Yinxia Li, Qianming Zhu, Jingxin Li... Pages 2155-2164

19. Original Article

Increasing predominance of G8P[8] species A rotaviruses in children admitted to hospital with acute gastroenteritis in Thailand,

2010-2013

Arpaporn Yodmeeklin, Pattara Khamrin, Kattareeya Kumthip... Pages 2165-2178

20. Original Article

Support

Genomic and biological characterization of Newcastle disease viruses isolated from migratory mallards (Anas platyrhynchos)

Momena Habib, Tahir Yaqub, Jawad Nazir, Wasim Shehzad... Pages 2179-2188



Skip to main content

🙆 Springer Link

Archives of Virology All Volumes & Issues

Special Topic-Section to honor Marc van Regenmortel (Pages 2017 - 2054)

ISSN: 0304-8608 (Print) 1432-8798 (Online)

In this issue (37 articles)

1. Original Article

Isolation and characterization of a group of new Proteus bacteriophages

V. Morozova, Yu. Kozlova, E. Shedko, I. Babkin, A. Kurilshikov... Pages 2189-2197

2. Original Article

Characterisation of the antigenic epitopes in the subunit 2 haemagglutinin of avian influenza virus H5N1

Khrisdiana Putri, Nadeeka Wawegama, Jagoda Ignjatovic... Pages 2199-2212

3. Brief Report

Identification and molecular characterization of a novel varicosa-like virus from red clover

Igor Koloniuk, Jana Fránová, Tatiana Sarkisova, Jaroslava Přibylová... Pages 2213-2218

4. Brief Report

<u>Microevolution and independent incursions as main forces shaping H5 Hemagglutinin diversity during a H5N8/H5N5 highly</u> pathogenic avian influenza outbrea<mark>k in Czech Re</mark>public in 2017

<u>Alexander Nagy, Ádám Dán, Lenka Černíková, Eliška Vitásková... Pages 2219-2224</u>

5. Brief Report

Molecular detection and phylogenetic analysis of megalocytivirus in Brazilian ornamental fish

Samara Rita de Lucca M<mark>aganha, Pedro Henrique</mark> Mag<mark>alhães Cardoso</mark>... Pages 2225-2231

6. Brief Report

Characterization of HIV-1 subtypes and transmitted drug resistance among treatment-naive HIV-infected individuals in Zhejiang,

<u>China, 2014-2017</u>

Yufan Xu, Xiaorong Peng, Xiuming Peng, Shujing Ji, Bin Chen... Pages 2233-2237

7. Brief Report

Screening of som<mark>e Croatian autochthon</mark>ous gra<mark>pev</mark>ine varieties reveals a multitude of virus<mark>es, includ</mark>ing novel ones

<u>Darko Vončina, Rodrigo P. P. Almeida</u> Pages 2239-2243

8. Brief Report

Molecular characterization of avipoxviruses circulating in Mozambique, 2016-2018

Lourenço P. Mapaco, Zeiss Lacerda, Iolanda V. A. Monjane... Pages 2245-2251

9. Annotated Sequence Record

Complete nucleotide sequences of the large RNA genome segments of Main Drain and Northway viruses (family Peribunyaviridae)

Jermilia Charles, Chandra S. Tang<mark>udu, Bradley J. Blitvich</mark> Pages 2253-2255

10. Annotated Sequence Record

Full genotype constellations o<mark>f six feline Rotavirus A strains</mark> isolated in Japan in the 1990s including a rare A15 NSP1 genotype

Toyoko Nakagomi, Chantal Ama Agbemabi<mark>ese, Osamu Nakagomi</mark> Pages 2257-2260

11. Annotated Sequence Record

Porcine-like G3P[6] and G4P[6] rotav<mark>irus A strains detecte</mark>d from children with d<mark>iarrh</mark>oea in V<mark>ietnam</mark>

<u>Miho Kaneko, Loan Phuong Do, Yen Hai Doan, Toyoko Nakagomi... Pages 2261-2263</u>

12. Annotated Sequence Record

Complete genome sequence of peanut virus C, a putative novel ilarvirus

Seungmo Lim, Yun-Woo Jang, Jin Woo Bae, Yeong-Hoon Lee ... Pages 2265-2269

13. Annotated Sequence Record

Sequencing, genome analysis and host range of a novel Ralstonia phage, RsoP1EGY, isolated in Egypt

Abdelmonim Ali Ahmad, Kamel M. Elhalag, Hardian Susilo Addy... Pages 2271-2274

14. Annotated Sequence Record

Complete genome sequence of yam chlorotic necrosis virus, a novel macluravirus infecting yam

Pingxiu Lan, Yu Meng, Pan Shen, Ruhui Li, Yan Ma, Songtao Tan... Pages 2275-2278

15. Annotated Sequence Record

A novel endornavirus isolated from cluster bean (Cyamopsis tetragonoloba)

Ricardo Iván Alcalá-Briseño, Ryo Okada, Favio Herrera... Pages 2279-2282

16. Virology Division News

Taxonomy of the order Mononegavirales: update 2018

Gaya K. Amarasinghe, Nidia G. Aréchiga Ceballos, Ashley C. Banyard... Pages 2283-2294

17. Virology Division News

Taxonomy of the family Arenaviridae and the order Bunyavirales: update 2018

Piet Maes, Sergey V. Alkhovsky, Yimíng Bào, Martin Beer... Pages 2295-2310

Support



https://doi.org/10.1007/s00705-018-3844-4

ANNOTATED SEQUENCE RECORD



Sequencing, genome analysis and host range of a novel *Ralstonia* phage, RsoP1EGY, isolated in Egypt

Abdelmonim Ali Ahmad^{1,2} · Kamel M. Elhalag^{3,4} · Hardian Susilo Addy^{1,5} · Mohamed A. Nasr-Eldin⁶ · Ahmed S. Hussien^{3,7} · Qi Huang¹

Received: 12 February 2018 / Accepted: 31 March 2018 / Published online: 13 April 2018 © This is a U.S. government work and its text is not subject to copyright protection in the United States; however, its text may be subject to foreign copyright protection 2018

Abstract

A novel *Ralstonia* phage was isolated from soil in Egypt. It was designated *Ralstonia* phage RsoP1EGY using our phage identifier naming approach to reflect the phage's bacterial host species, characteristics and origin. When tested, this phage specifically infected only race 3 biovar 2 phylotype IIB sequevar 1, and not non-race 3 biovar 2 strains of *Ralstonia solanacearum*. The phage has an icosahedral capsid of 60 ± 5 nm in diameter with a short tail of 15 ± 5 nm in length, typical of a podovirus. The genome of RsoP1EGY is 41,297 bp in size, containing 50 open reading frames, with no significant sequence identity to any other reported *R. solanacearum* or non-*Ralstonia* phages, except to the recently deposited but unreported and unclassified *Ralstonia* phage DU_RP_I. RsoP1EGY is the first sequenced and characterized *R. solanacearum* phage isolated in Egypt.

Potato is the second most important vegetable crop next to tomato in Egypt. Egypt also ranks among the world's

Handling Editor: Johannes Wittmann.

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s00705-018-3844-4) contains supplementary material, which is available to authorized users.

Qi Huang qi.huang@ars.usda.gov

- ¹ Floral and Nursery Plants Research Unit, United States National Arboretum, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, USA
- ² Department of Plant Pathology, Faculty of Agriculture, Minia University, El-minia, Egypt
- ³ Bacterial Diseases Research Department, Plant Pathology Research Institute, Agricultural Research Center (ARC), Giza, Egypt
- ⁴ Potato Brown Rot Project, Ministry of Agriculture, Dokki, Giza, Egypt
- ⁵ Department of Plant Protection, Faculty of Agriculture, University of Jember, Jember, Indonesia
- ⁶ Botany Department, Faculty of Science, Benha University, Benha, Qalyubiya Governorate, Egypt
- ⁷ Central Administration of Plant Quarantine, Dokki, Giza, Egypt

top potato exporters, mainly for markets in Europe. Potato brown rot caused by cool-temperature-adapted race 3 biovar 2 (r3b2) strains of *Ralstonia solanacearum* is a serious problem in Egypt that affects potato production and exportation. As a result, *R. solanacearum* is a quarantine pathogen with zero tolerance in Egypt. The r3b2 strains of *R. solanacearum* are also quarantined pathogens in Europe and select agents in the United States [1]. Bacteriophages that specifically infect *R. solanacearum* have recently been isolated from Japan, Thailand, Korea and the United States, and efforts have been made to determine their potential as biocontrol agents for *R. solanacearum* [2–5].

In this study, a bacteriophage was isolated from the soil of an *R. solanacearum*-infested potato field in 2017 in Al Qalyubia governorate of Egypt using *R. solanacearum* strain K3 as the host, based on the method of Ahmad et al. [6]. The phage was purified from a single-plaque isolate, and a phage susceptibility test was performed (Table 1) using a spot test followed by a plaque assay as described by Ahmad et al. [2]. *R. solanacearum* was cultured as described [2], and strains UW550 and UW552 were routinely used as hosts for propagation of the phage. A total of one liter of bacterial culture (ten 100-ml cultures) was grown in order to obtain a sufficient amount of the phage particles. When the bacterial culture reached the absorbance of 0.2 at 600 nm, phage suspension was added at a

A. A. Ahmad et al.

Table 1Susceptibility ofRalstonia solanacearum strainsto Ralstonia phage RsoP1EGY

Ralstonia solancearum (Phylotype-sequevar/biovar)	Origin	Source	RsoP1EGY ^a
Race 3 biovar 2 strains			
UW257 (IIB-1)	Costa Rica	C. Allen, USA	+
UW344 (IIB-1)	Brazil	C. Allen, USA	+
UW550 (IIB-1)	Netherlands	C. Allen, USA	+
UW551 (IIB-1)	Kenya	C. Allen, USA	+
UW552 (IIB-1)	Guatemala	C. Allen, USA	+
RUN035 (IIB-1)	Netherlands	P. Prior, France	+
IVIA1602.1 (IIB-1)	Spain	M. M. López, Spain	+
K3 (IIB-1)	Egypt	This study	+
K10 (IIB-1)	Egypt	This study	+
K13 (IIB-1)	Egypt	This study	+
Non-race 3 biovar 2 strains			
Rs5 (IIA-7/1)	USA	J. Jones, USA	-
Rs121 (I/1)	USA	J. Jones, USA	-
AW1)II-7/1)	USA	C. Allen, USA	-
K60 (IIA-7/1)	USA	C. Allen, USA	-
UW119 (I/3)	Costa Rica	C. Allen, USA	-
GM1000 (I-18/3)	French Guiana	P. Prior, France	-
RUN302 (IIB-4/1)	Brazil	P. Prior, France	-
Pss530 (I/3)	Taiwan	J. F. Wang, Taiwan	-
Pss51 (I-15, I/4)	Taiwan	J. F. Wang, Taiwan	<u> </u>
DT3 (I/3)	Indonesia	H. Addy, Indonesia	-

^aSusceptibility of *R. solanacearum* strains to phage RsoP1EGY was determined by spot test followed by plaque assay. It is shown as susceptible (+) when clear plaques were observed or resistant (-) when no plaques were observed

multiplicity of infection of 0.1. After further growth for approximately 20 h, bacterial cells were removed, and phage particles were purified for morphological characterization under a Hitachi 7700 transmission electron microscope and for DNA extraction as described by Ahmad et al. [2], except that the supernatant was filtered through a 0.45-µm membrane before ultracentrifugation.

The phage has an icosahedral capsid of 60 ± 5 nm in diameter with a short tail of 15 ± 5 nm in length (Fig. 1A), the typical morphology of members of the family *Podoviri*dae. We named the phage Ralstonia phage RsoP1EGY, since it is the first R. solancearum-infecting bacteriophage belonging to the family *Podoviridae*, and it was isolated in Egypt. Our naming of the phage was based on the informal guide by Adriaenessens and Brister [7] (i.e., a tripartite phage name - Ralstonia phage RsoP1EGY) and the scheme proposed by Kropinski et al. [8] (i.e. Rso for R. solanacearum and P for Podoviridae). Additionally, our name took the sequential number of the phage we isolated in the same virus family into consideration (i.e., 1 for the first isolated phage), as well as where the phage was originally isolated, by including a three-letter country code (i.e., EGY for Egypt). This naming approach aims to not only make the identifier name unique but also to make it systematic and meaningful to other researchers regarding the phage's bacterial host species, characteristics and origin.

Genomic DNA of phage RsoP1EGY was extracted from purified phage particles (described above) using either a Phage DNA Isolation Kit (Norgen Biotek Corp, Canada), or the phenol-chloroform method [9]. The phage DNA was sequenced and the genome sequence assembled commercially by SeqMatic LLC (Fremont, California). Potential open reading frames (ORFs) larger than 40 amino acids were identified using PHAST (http://phast.wishartlab.com) and GeneMark [10]. Homology searches for each identified ORF were performed using BLAST/PSI-BLAST [11] against NCBI's protein databases [12, 13]. An e-value threshold of e^{-4} or less was used for two proteins to be considered a match.

The genome of phage RsoP1EGY is a double-stranded DNA of 41,297 bp in length, with a G+C content of 59% (GenBank accession no. MG711516). When the complete genome sequence of phage RsoP1EGY was used as a query to search NCBI's nucleotide collection by BLASTn, the phage genome has no significant sequence identity to any of the previously published *R. solanacearum* phages, except that it shared 99% nucleotide sequence identity with 95% coverage with a recently deposited but unpublished




Fig. 1 Electron micrograph of *Ralstonia* phage RsoP1EGY virions (**A**) and genome organization of phages RsoP1EGY and DU_RP_I, with reference to the *Enterobacteria* phage T7 (**B**). The scale represents 100 nm. Arrows represent the size and direction of transcription of ORFs. Functional gene clusters I for early genes, II for DNA metabolism, and III for virion structure and assembly of phages

RsoP1EGY and **DU_RP_I** are assigned according to the typical podovirus genome represented by the T7 phage [12] and marked in red, green and blue, respectively. Numbers are annotated ORFs in each phage. RNAP, RNA polymerase; DNAP, DNA polymerase; MCP, major capsid protein; Ter, terminase

"*Ralstonia* phage DU_RP_I" (GenBank accession no. MF979559.1; Fig. 1B), although the Egyptian phage is 640-nucleotides longer than phage DU_RP_I. It is unknown, however, what *Ralstonia* species or *R. solanacearum* strains the DU_RP_I phage infects. The fact that phage RsoP1EGY belongs morphologically to the family *Podoviridae* but does not share significant sequence similarity with any of the previously characterized *R. solanacearum* phages of the family *Podoviridae* suggests that the Egyptian phage maybe a new member of the family *Podoviridae*.

Fifty putative ORFs were identified in phage RsoP1EGY. Their positions, best homologs and predicted functions are summarized in the Supplementary Table S1. Among the 50 ORFs, 20 had homologs with known functions, including phage structural proteins (ORFs 38-39, 41, 46-47, and 50), early genes involved in DNA processing (ORFs 6, 18, 20,-24, 27, 30, 34, and 42), and late gene products such as lysis proteins (ORFs 2, 3, and 25). Twelve are hypothetical proteins showing marginal similarities with hypothetical proteins in Burkholderia ubonensis (ORF 9) or the Ralstonia phage DU_RP_I (ORFs 4-5, 26, 28, 31-33, 35, 40, and 48-49). Eighteen ORFs, however, had no similarity to any proteins in the databases (Table S1). The genome arrangement of phage RsoP1EGY and the 39% amino acid sequence identity between the putative RNA polymerase of phage RsoP1EGY (ORF20) and the T7 RNA polymerase

(GenBank accession no. NP_041960.1) suggest that RsoP-1EGY may be a T7-like phage. A typical podovirus genome represented by Enterobacteria phage T7 generally consists of three functional gene clusters: cluster I for early functions, cluster II for DNA metabolism, and cluster III for structural proteins and virion assembly (Fig. 1B) [14]. We tentatively assigned the three clusters to phage RsoP1EGY and its closely related phage DU RP I and compared the two phages to the T7 phage (Fig. 1B). The putative cluster I of phage RsoP1EGY includes ORF8 to ORF20, containing a putative integrase (ORF18) and a RNA polymerase (ORF20) (Fig. 1B, Table S1). The putative cluster II includes ORF21 to ORF37, containing annotated phage single-stranded DNA-binding protein (ORF22), endonuclease VII (ORF24), lysozyme (ORF25), DNA primase (ORF27), DNA polymerase (ORF30), and exonuclease (ORF34) (Fig. 1B, Table S1). The putative cluster III includes ORF38 to ORF50 and ORF1 to ORF7 containing putative phage tail proteins (ORFs 38, 46 and 47), phage capsid assembly protein (ORF39), phage capsid protein (ORF41), lytic transglycosylase protein (ORF2), bacteriolytic protein (ORF3), and terminase protein large subunit (ORF6) (Fig. 1B, Table S1).

To determine the host specificity of phage RsoP1EGY, 20 *R. solanacearum* strains in our collection were tested for phage susceptibility using the method of Ahmad et al. [2]. Interestingly, none of the 10 non-r3b2 strains, but all 10

of the r3b2 strains (phylotype IIB sequevar 1) tested were susceptible to phage RsoP1EGY (Table 1). This is probably because phage RsoP1EGY was isolated from the Nile delta region of Egypt, where only r3b2 strains of *R. solanacearum* causing potato brown rot were detected [15]. Further research is needed to determine whether the Egyptian phage has potential as a biocontrol agent for r3b2 strains of *R. solanacearium* in Egypt or other countries where potato brown rot is a problem, and whether the phage can be used for specific detection of r3b2, since the phage is lytic to and specific for *R. solanacearum* r3b2.

It is interesting that phage RsoP1EGY is closely related to the *Ralstonia* phage DU_RP_I, but not to any other reported *R. solanacearum* or non-*Ralstonia* phages. It will be interesting to study the evolution of these two *Ralstonia* phages once the origin, isolation, morphology and host range of phage DU_RP_I becomes available.

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- 1. Agricultural Bioterrorism Protection Act of 2002 (2002) Possession, use, and transfer of biological agents and toxins; interim rule and request for comments. Fed Regist 240(67):76907–76938
- 2. Ahmad AA, Stulberg MJ, Mershon JP et al (2017) Molecular and biological characterization of ϕ Rs551, a filamentous bacteriophage isolated from a race 3 biovar 2 strain of *Ralstonia solanacearum*. PLoS One 12:e0185034. https://doi.org/10.1371/journ al.pone.0185034
- 3. Yamada T (2012) Bacteriophages of *Ralstonia solanacearum*: their diversity and utilization as biocontrol agents in agriculture.

In: Kurtboke I (ed) Bacteriophages. InTech, London, pp 113–138. https://doi.org/10.5772/33983

- Bhunchoth A, Phironrit N, Leksomboon C et al (2015) Isolation of *Ralstonia solanacearum*-infecting bacteriophages from tomato fields in Chiang Mai, Thailand, and their experimental use as biocontrol agents. J Appl Microbiol 118:1023–1033
- Murugaiyan S, Bae JY, Wu J et al (2011) Characterization of filamentous bacteriophage PE226 infecting *Ralstonia solanacearum* strains. Appl Microbiol 110:296–303
- Ahmad AA, Askora A, Kawasaki T et al (2014) The filamentous phage XacF1 causes loss of virulence in *Xanthomonas axonopodis* pv. *citri*, the causative agent of citrus canker disease. Front Microbiol 5:321. https://doi.org/10.3389/fmicb.2014.00321
- 7. Adriaenssens EM, Brister JR (2017) How to name and classify your phage: an informal guide. Viruses 9:70. https://doi. org/10.3390/v9040070
- 8. Kropinski AM, Prangishvili D, Lavigne R (2009) Position paper: the creation of a rational scheme for the nomenclature of viruses of bacteria and archaea. Environ Microbiol 11:2775–2777. https ://doi.org/10.1111/j.1462-2920.2009.01970.x
- Sambrook J, Russell DW (2001) Extraction of bacteriophage λ DNA from large-scale cultures using proteinase K and SDS. In: Sambrook J, Russell DW (eds) Molecular cloning: a laboratory manual, 3rd edn. Cold Spring Harbor Laboratory Press, New York, pp 2.56–2.58
- 10. Besemer J (2001) GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. Nucleic Acids Res 29:2607–2618
- 11. Altschul SF, Madden TL, Schaffer AA et al (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25:3389–3402
- 12. UniProt Consortium (2007) The universal protein resource (Uni-Prot). Nucleic Acids Res 35:D193–D197
- Wheeler DL, Barrett T, Benson DA et al (2007) Database resources of the national center for biotechnology information. Nucleic Acids Res 35:D5–D12
- Dunn JJ, Studier FW (1983) Complete nucleotide sequence of bacteriophage T7 DNA and the locations of T7 genetic elements. J Mol Biol 166:477–535
- 15. Tomlinson DL, Elphinstone JG, Soliman MY et al (2009) Recovery of *Ralstonia solanacearum* from canal water in traditional potato-growing areas of Egypt but not from designated Pest-Free Areas. Eur J Plant Pathol 125:589–601