

World Journal of Advance Heathcare Research



Published By
WJAHR

Editorial Board

Editor in Chief

Prof. Dr. Starodub, Nickolaj Fedorovich

nikstarodub1@yahoo.com

National University of Life and Environmental Sciences of Ukraine
Associate Editor

Prof. Dr. N.S. NEKI

drneki1234@gmail.com


(MD , DTN & H, FRCP (Edin), FRCP(Ire), FRCP(Glas), FCCP(USA), FRCP(Lon), FACC(USA), FACP (USA), FACE(USA), FESC, FICP, FGSI, FRSM(London), FNCCP , FICN, FACN(USA), FSCH(Ind)) Professor of Medicine, Govt. Medical College, Amritsar, Punjab, India

Editorial Board Member

S.No.	Name	Email	Address	Country
1	Dr. Ajay Bilandi	ajaybilandi20311@gmail.com	Associate Professor Seth G. L. Bihani S. D. College of Technical Education, Institute of Pharmaceutical Sciences & Drug Research, Sri Ganganagar (Rajasthan)	India
2	Dr. Khawaja Shakeel Ahmed	khajashakeel1@gmail.com	Assistant Professor (Microbiology) Head Department of BMSC School of Allied Health Professionals, Asmara College of Health Sciences, Asmara, State of Eritrea	Eritrea
3	Dr. Vineeta Negi	drvineetanegi1@gmail.com	Associate Professor, Chaudhary Devi Lal Group of Institutions, Haryana	India
4	Prof. Dr. N. Dora Babu	ndorababu811@gmail.com	Professor Mother Teresa Pharmacy College Sanketika Nagar, Sathupally, Khammam District. Telangana	India
5	Dr. Islam Mostafa Abdo Bayomi Elbaz	islam_mostafa_elbaz@yahoo.com	Lecturer of Pharmacognosy Faculty of Pharmacy, Zagazig University, 44519-Zagazig	Egypt
6	Dr. Kundlik Girhepunje	kundlik@gmail.com	School of Pharmacy, Peoples University, Bhopal	India
7	Dr. Shikha Thakur	patho.shikha@gmail.com	Assistant Professor - Department of Microbiology, SIPAS, Hemwati Nandan Bahuguna University, India	India
8	Obeagu, Emmanuel Ifeanyi	emmanuelobeagu@yahoo.com	Diagnostic Laboratory Unit, University Health Services Department, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria	Nigeria
9	Dr. Abd El-Aleem Saad Abdelaleem2011@yahoo.com Soliman Desoky		Professor of Agricultural Zoology Assistant in Plant Protection Department), Faculty of Agriculture, New Campus (ElKawamel), Sohag University, New Sohag City, Sohag, Egypt	Egypt
10	Prof. Katalin Prokai-Tatrai	Katalin.Prokai@unthsc.edu	Center for Neuroscience Discovery, Department of Pharmaceutical Sciences University of North Texas Health Science Center 3500 Camp Bowie Boulevard Ft. Worth, Texas 76107,	USA

Digital Repository Universitas Jember

S.No.	Name	Email	Address	Country
11	Dr. Franck YOVO	yovofranck62@gmail.com	USA. Doctor in Physical Chemistry/ Environmental Chemistry, Beninese.	Beninese
12	Dr. Mohammed Nader Shalaby	dr.m.nader@hotmail.com	Lecture, Department of Biological Sciences and Sports Health, Faculty of Physical Education Suez Canal University, Egypt	Egypt
13	Professor Dr. Jacinta A. jaopara@yahoo.com Opara	jaopara@yahoo.com	Professor and Director, Centre for Health and Environmental Studies, University of Maiduguri, PMB 1069, Maiduguri- Nigeria	Nigeria
14	Dr. Sreekanth sreekanthbose@gmail.com B.	sreekanthbose@gmail.com	BDS, MDS, Department of Public Health Dentistry, India	India
15	Dr. U. usreenivas1966@rediffmail.com	Srinivasa	Professor and Head of the Department of Pharmacognosy and Phytochemistry, India	India
16	Dr. Mahesh mchsahu@gmail.com	Chandra Sahu	Assistant Professor, Directorate of Medical Research, IMS & Sum Hospital (Faculty of Medical Sciences), Siksha 'O' Anusandhan University, Kalinga Nagar, Bhubaneswar 751003, Odisha, India	India
17	Dr. Nisa Uyar nisa.uyar@hotmail.com	Alpaslan	Mersin University Medical Faculty, Department of Medical Biology and Genetics, Turkey	Turkey



DETERMINANTS OF MATERNAL HEALTH, MATERNAL MORTALITY AND MOBILITY IN ADO-EKITI, EKITI STATE OF NIGERIA

Okpoko P. I.* and James R.

ORBITO-OCULAR MALIGNANCIES IN MAKURDI, NIGERIA

***CO. Ojabo, BA. Ojo, I. Ugwu, E. Umabong, J. Ngbea, R., Vhritherhire, I. Akpor, C. Udu**

PEDIATRIC ORBITAL AND OCULAR MALIGNANCIES IN MAKURDI, NIGERIA.

***C. O. Ojabo, B. A. Ojo, I. Ugwu, E. Umabong, J. Ngbea, R. Vhritherhire, I. Akpor and C. Udu**

THE LIPID PROFILE IN CHRONIC OBSTRUCTIVE PULMONARY DISEASE AND CORRELATION WITH DISEASE SEVERITY

Dr. Ali Raheem Ali, Dr. Jaleel Okaish Hadi, Professor Dr. Adnan M. Al, Jubouri and Professor Dr. Mohammed Waheeb Al-Obaidy*

UTILIZATION AND ALLOCATION FOR VILLAGE FUND IN THE PROMOTION OF FAMILY PLANNING VIA UPTAKE LONG ACTING AND PERMANENT METHOD (LAPM) IN WEST NUSA TENGGARA PROVINCE, INDONESIA

Surotul Ilmiyah*, Rita Damayanti, Ede Surya Darmawan, Iwan Ariawan, Hoirun Nisa, Hendri Hartati, Ousman Bajinka

EFFECT OF CONVENIENCE AND BENEFIT ON THE ATTITUDE OF USING THE HOSPITAL MANAGEMENT INFORMATION SYSTEM IN BALUNG GENERAL HOSPITAL JEMBER REGENCY

Hafis Nur Wicaksono*, Isa Ma'rufi and Dwi Wahyuni

ANAESTHETIC TECHNIQUES FOR OPHTHALMIC SURGERIES IN BENUE STATE UNIVERSITY TEACHING HOSPITAL (BSUTH):- A REVIEW OF 100 CONSECUTIVE CASES.

***Efu ME, Ojabo CO, Ojo BA, Chaha K, Eke BA Ozuagu MA, Anefu GO**

HAEMOSTATIC FACTORS ASSESSMENT AMONG MALE AND FEMALE GERIATRICS IN NNEWI, ANAMBRA STATE NIGERIA.

Onwurah O. W.*, Amilo G. I., Ajuba I., Fasakin K. A., Muhibi M. A. and Ezeugwunne I. P.

IMPACT OF EDUCATIONAL INTERVENTION ON KNOWLEDGE OF UNIVERSAL PRECAUTIONS AMONG HEALTH CARE STAFF OF A TERTIARY CARE HOSPITAL OF SOUTH GUJARAT

Kamlesh Shah*, Baria Hinal, Alka Nerurkar and Diwarkar Sharma

CONFRONTING THE CHALLENGES OF REDUCING HIV AND OTHER SEXUALLY TRANSMITTED INFECTIONS AMONG IN-SCHOOL ADOLESCENTS IN THE RURAL AREAS OF IMO STATE OF NIGERIA.

Ezeama, M.C. and Enwereji, E. E.*

EPIDEMIOLOGICAL PARTICULARITY OF ANTHRAX IN RUMINANTS IN KANKAN AND KOUBIA PREFECTURES (REPUBLIC OF GUINEA)

***Mamadou Dian D., Youssouf S., Ekatherina Vladimirovna N., Sanaba B. and Ansoumane S.**

CONCURRENT INFECTION OF DENGUE CHIKUNGUNYA AND SCRUB TYPHUS IN JHARKHAND, INDIA

Dr. Amar Verma*, Dr. Antony Kisku and Dr. Vishwajeet Pratap

PREVALENCE OF STRESS AMONG PARENTS HAVING HEARING IMPAIRED CHILD IN NORTH INDIA

Rina Kumari*, Sunita Tiwari and Sunil Kumar

ANALYSIS BASIC EMERGENCY OBSTETRIC AND NEONATAL ON THE QA PROGRAM IN PONED PUBLIC HEALTH CENTER X BONDOWOSO REGENCY

Asti Nuris Mida Rahmawati*, Al Munawir and Dewi Rokhmah

CONTENT OF SULFUR DIOXIDE (SO₂) IN THE AIR ON THE OCCURRENCE OF UPPER RESPIRATORY INFECTION (URI) IN FARMERS IN IJEN DISTRICT WATERSHED, BONDOWOSO

Aditya Sapta Wardana*, Isa Ma'rufi and Ristya Widi Endah Yani

THE COMMONEST RESPIRATORY INFECTIONS THAT INDICATE HOSPITAL ADMISSION AMONG CHILDREN IN MOSUL

***Dr. Hajir H. Al-Ridhwany, Khulood J. Mahmood and Farah S. Yonan**

FACTORS AFFECTING UNMET NEED FOR CONTRACEPTION AMONG REPRODUCTIVE AGED WOMEN IN BANGLADESH: DOES EMPLOYMENT STATUS MATTER?

Ahmed Zohirul Islam*, Mt. Laily Khatun, Md. Nazrul Islam Mondal, Ahmed Eftekharul Islam, Md. Rafiqul Islam and Md. Golam Mostofa

PHARMACOGNOSTICAL AND CHEMICAL COMPARISON OF THE SEED AND ARIL OF MYRISTICA FRAGRANS HOUTT.

***Gayathri Devi V., Anitha John, Natarajan M., Neethu Kannan B. and Kanagarajan A.**

EVALUATION OF C-REACTIVE PROTEIN LEVEL IN PATIENTS WITH RENAL DISEASE IN NNAMDI AZIKIWE UNIVERSITY TEACHING HOSPITAL, NNEWI, ANAMBRA STATE, NIGERIA

Onwurah O. W.*, Ikebido O. C., EZE. H. T., Nkemjika U. N., Ifeanyichukwu M. O., Amilo G. I., Ajuba I., NWACHUKWU E.P

FREQUENCY AND DURATION OF BREASTFEEDING AMONG CHILDREN HAVING ACUTE RESPIRATORY INFECTIONS IN MOSUL

Dr. Khulood J. Mahmood* and Zaid M. Yassen

IMPROPER TECHNIQUE OF USING TURBUHALER IN A SAMPLE GROUP OF IRAQI ASTHMATIC PATIENTS, AFFECTING FACTORS AND ITS EFFECT ON ASTHMA CONTROL

Dr. Mahmood Fouad Mahmood M.B.Ch.B., Dr. Muftafa Ali Al. Sakini M.B.Ch.B., Dr. Ibtihal Shukri Abd Alhaleem M.B.Ch.B, FICMS-FM. and Professor Dr. Mohammed Waheeb. Al-Obaidy*

COMPARISON OF DEXAMETHASONE AND TRIAMCINOLONE AS EPIDURAL STEROIDS IN LOW BACK PAIN

Harparam Singh Ghuman*, Hardayal Singh Ghuman, Sumera Rahim

EFFECTIVITY OF LEECH THERAPY IN VARICOSE VEINS

Md. Anwer Alam, Zarnigar and Md. Tanwir Alam*

PREVALENCE OF ANEMIA AMONG THE ADOLESCENT FEMALES

Dr. Talat Nahid Ramzan Ali, Prof. S. T. A. Bilgrami, D. Md Razi Ahmad, Dr. Md Najibur Rahman and Dr. Md Tanwir Alam*

PEDIATRIC AKI - A COMPARISON OF PRIFLE, AKIN, AND KDIGO DEFINITIONS

Dr. Nirranjana Sakthivelu and Dr. Divya Singh*

DYSLIPIDAEMIA AMONG PATIENTS WITH END STAGE RENAL DISEASE IN MAKURDI, NORTH CENTRAL, NIGERIA

Dr. Monday O. Ogiator*, Joseph E. Ojobi and Ochoche O. Ijachi

MODERN VIEW ON THE MECHANISM OF "THOUGHT" FORMATION AND ITS IMPLEMENTATION "PROGRAM" AT THE SUPRAMOLECULAR LEVEL

Dr. Bitsoev Vladimir Dодtievich*

CASE REPORT

ACQUIRED TESTICULAR AND EPIDIDYMAL LYMPHANGIOMA: A RARE CASE REPORT AND REVIEW

CASE REPORT OF TWO RARE TUMORS IN THE SPHENOID SINUS

Juvaina P., Priyanka Venu*, Devarajan E, Rajan P., Naufal P. and Saanida M. P.

ESTHETIC REHABILITATION OF DISCOLORED MAXILLARY ANTERIORS WITH MINIMALLY INVASIVE PORCELAIN LAMINATE VENEERS – A CASE REPORT

*Dr. Param Dev Singh (MDS), Dr. Geetha Bhavani Paluri (MDS), Dr. Geeta Arya (MDS) and Dr. Payal Gumber

A RARE CASE OF DIFFUSE INFILTRATING RENAL CELL CARCINOMA WITH EXTENSIVE VENOUS THROMBOSIS

Juvaina P.*, Neema Jaleel, Devarajan E., Rajan P., Naufal P. and Saanida M. P.

POST-DIARRHEA HEMOLYTIC UREMIC SYNDROME MIMICKING SEVERE ACUTE COLITIS IN ADULTS: ABOUT A RARE CASE AND REVIEW OF THE LITERATURE

Krati K.*, Lemfadli Y., Jiddi S., Benjelloun Z., Ait Errami A., Oubaha S. and Samlani Z.

DIGESTIVE HAEMORRHAGE: PLACE OF SCINTIGRAPHY COUPLED WITH TECHNICIUM 99

Jiddi S.*, Jellal S., Lmfadli Y., Bouchrit S., Ait Errami A., Samlani Z., Oubaha S. and Krati K.

MALE BREAST CANCER: A CASE REPORT AND REVIEW OF LITERATURE

Dr. Akpor I. O.*, Vhritherhire R. A., Ngbea J. A., Ojo B. A., Eke B. A., Nyaga T., Gyenger D.

GIANT OVARIAN FIBROMA: A CASE REPORT AND REVIEW OF LITERATURE

*Akpor I. O., Vhritherhire R. A., Ojo B. A., Nyaga T., Terkula D. G. and Agwa M.

AYURVEDIC MANAGEMENT OF AN INFECTED WOUND: A CASE STUDY

Manorma Singh*, Sanjeev Sharma, Rahul Sharma

REVIEW ARTICLE

EMERGENCY MEDICAL TREATMENT AND LABOR ACT

*Dr. Shinu Kuriakose DHSc PA-C

POST-LOBECTOMY DIFFICULTY IN WEANING OFF THE VENTILATOR-IS IT EDAC?

Dr. Hemant Gopal Bhirud*, Dr. Amit Bodkhe, Dr. Chitra Pitale, Dr. Sanjay Upadhye

CARDIOVASCULAR DISEASE - RISE IN INDIA: NURSES ROLE TO CURB THE MENACE

Vinil Upendrababu* and Dr. Prof. S. N. Nanjunde Gowda

MANAGEMENT OF COMMON POISONING

Dr. Bhartendra Sharma*

FIBER REINFORCED MATERIALS: A NOVEL APPROACH TOWARDS IMPROVED ESTHETIC DENTISTRY

Dr. Anwetakshmi Ray*, Dr. Kundabala M., Dr. Ankita Kanotra, Dr. Ayushi Bajoria, Dr. Janina Loren DSouza and Dr. Afroz

FACTORS INFLUENCING CONTRACEPTIVE USE AMONG WOMEN OF REPRODUCTIVE AGE IN THE BIA WEST DISTRICT OF GHANA: A COMMUNITY BASED CROSS-SECTIONAL STUDY

Awuni Prosper Mandela Amaltinga*, Mbinta Fenibe James and Abugri Osman Bright

PERTUSIS: FACTS AND MANAGEMENT

K. Sen*, Dr. B. Ray and Prof. S. K. Mahapatra

A COMPREHENSIVE REVIEW ON MATRIX TYPE TRANSDERMAL PATCHES

Asheesh Chauhan* and Dr. Kapil Kumar

MODERN CONCEPTS OF THE "MECHANISM OF ACTION" OF THE PHYSICAL FACTOR ON THE WHOLE ORGANISM FROM THE POSITION OF QUANTUM PHYSICS

Dr. Bitsoev Vladimir Dottiievich*

CLINICAL EFFICACY OF COMPOUND UNANI FORMULATION IN NIQRIS (GOUT)


Md. Kalam Ansari, Md. Razi Ahmad, Md. Najibur Rahman and Md. Tanwir Alam*

DAWALI (VARICOSE VEINS): DESCRIPTION IN UNANI SYSTEM OF MEDICINE

Md. Anwer Alam, Zarnigar and Md. Tanwir Alam*

LEECHING: THE ANCIENT ART OF HEALING

Md. Anwer Alam, Zarnigar and Md. Tanwir Alam*



CONTENT OF SULFUR DIOXIDE (SO₂) IN THE AIR ON THE OCCURRENCE OF UPPER RESPIRATORY INFECTION (URI) IN FARMERS IN IJEN DISTRICT WATERSHED, BONDOWOSO

Aditya Sapta Wardana*¹, Isa Ma'rufi² and Ristya Widi Endah Yani³

¹Occupational Health and Safety, Magister of Public Health, University of Jember.

²Lecturer at Faculty of Public Health, University of Jember.

³Lecturer at Faculty of Dentistry, University of Jember.

Received date: 06 February 2020

Revised date: 27 February 2020

Accepted date: 17 March 2020

*Corresponding author: Aditya Sapta Wardana

Occupational Health and Safety, Magister of Public Health, University of Jember.

ABSTRACT

The objective of this research was to analyze the effect of Sulfur Dioxide (SO₂) content on Upper Respiratory Infection through individual factors (age, duration of exposure, length of work, smoking habits, nutritional status, independent efforts to reduce exposure (personal protective equipment) and environment factors (temperature, humidity, ventilation). This is an observational analytic research with a cross sectional approach with a sample of 101 respondents. The data used are primary data obtained from SO₂ measurements, Upper Respiratory Infection examination, and questionnaires for 101 farmer respondents. The analysis instrument uses SPSS 22.0. The results of data analysis show that: 1) SO₂ source of exposure has a significant effect on the incidence of URI. 2) Individual factors (age, length of work, duration of exposure, smoking habits, independent efforts to reduce exposure) and environment factors (temperature, humidity, ventilation) have a significant effect on the occurrence of URI.

KEYWORDS: Sulfur dioxide, upper respiratory infection, farmer.

INTRODUCTION

The activity of Ijen Crater can produce toxic gases, sulfur sublimation, volcanic dust, and acidity in the crater water and the content of fluoride (F). This crater water flows upstream of Kali Pahit until it flows to Blawan plantation area of Ijen District such as three hamlets, which are Curah Macan, Margahayu and Watu Capil, then it flows across several regions in the eastern Situbondo District (Aminuddin 2014).^[1]

The content of SO₂ toxic gas in river water can pollute the environment in Ijen District watershed area, and it can cause harm to health. WHO states that the ambient air quality has a chemical effect on the health of the respiratory tract. Poor air quality can also cause the possibility of being infected with several respiratory diseases and one of them is URI (Upper Respiratory Infection). Viruses or bacteria are the causes of human being exposed to acute respiratory infections. This disease lasts for 14 days and can also more than 14 days. The etiology of URI consists of 300 types of bacteria, viruses, and rickets. Common symptoms of this disease are cough, sore throat, difficulty of breathing, runny

nose, fever and earache (Ministry of Health of the Republic of Indonesia 2014).^[2]

The data obtained from the Community Health Center of Ijen shows that URI disease always becomes the highest disease in the monthly report in Ijen District. Wednesday (3/21/2018) at 19.13 Western Indonesian Time. Kali Pahit Riverflow exposes a very pungent odor of toxic gas, which is then inhaled by residents living in the watershed of Watu Capil River, Margahayu, Kali Bahit, Kalianyar Village, Ijen District, Bondowoso Regency, East Java. Two hundred residents in the area were evacuated. While the number of victims resulting from exposure to poison gas reached 30 people. Twenty four of them were treated at the Community Health Center of Ijen, then four people were treated at the Community Health Center of Tlogosari, and two people were rushed to the Regional Hospital of Koesnadi (Indonesian Regional Board for Disaster Management of Bondowoso 2018).^[3]

The research on SO₂ content and URI occurrence conducted in Surabaya, based on the level of pollution caused by motorized vehicles with an average SO₂

content of 11.6 $\mu\text{g}/\text{m}^3$ in Rungkut District, causes URI occurrence of 12.73 per 1,000 population and 25.12 per 1,000 residents in Jambangan District. Therefore, it can be concluded that there is a relation between the occurrence of URI and SO_2 content (Firdaus A, 2017).^[4] The factors that can affect the risk of someone being exposed to URI can be divided into four broad lines, namely pollution, individual characteristics, working behavior, and environment factors. From some of the problems aforementioned above, the researchers are interested in conducting a study and measurement of air SO_2 content in the watershed area of Ijen District, Bondowoso. In addition, the researchers are also willing to know the distribution of farmers who experience symptoms of URI.

MATERIALS AND METHODS

Quantitative approach is the method used in this research. The research used is analytic observational

RESULT AND DISCUSSION

1.1 Univariate Analysis

Table 1: Source of SO_2 Exposure.

Variable	Category	SO_2 Content	Information
Source of SO_2	Kali Pahit	33,6	Above Threshold Value
	Watu Capil	6,9	Above Threshold Value
	Farming	4,7	Above Threshold Value
	Residents' House	9,27	Above Threshold Value

Table 1 The highest SO_2 measurement (Table 1) in Kali Pahit watershed is 33.65 (mg/m^3) above the Threshold Value (NAV). The lowest measurement result (SO_2) is

study with cross sectional approach. The population in this study was all farmers around Watucapil watershed of 101 people. The sample used in this study was all farmers who were around Watucapil watershed (total sampling).

This research has passed the ethics test in the research ethics commission at the Faculty of Dentistry, University of Jember with number 540/UN25.8/KEPK/DL/2019. Data collection is done by interview, documentation, observation and measurement. Measurement is made to measure content of SO_2 in collaboration with Technical Implementing Unit of Occupational Health and Safety of Surabaya by using minipump and implinger. Then, the URI examination is carried out by a doctor from the local health center. The analysis used is logistic regression using SPSS 22.0.

in the watershed are (plantation) that is 4.7 (mg/m^3) below the Threshold Value (NAV).

Individual Factors

Table 2: The distribution of Individual Factors.

Variable	Category	N	%
Age	36 – 45 y.o	25	24,7
	46 – 55 y.o	42	41,6
	>56 y.o	34	33,7
	Total	101	100
Duration of Exposure	<4 hrs	54	53,5
	>4 hrs	47	46,5
	Total	101	100
Length of Work	6-10 yrs	43	42,6
	>10 yrs	58	57,4
	Total	101	100
Smoking Habit	Not smoking	47	46,5
	Light Smoker	54	53,5
	Total	101	100
Nutritional Status	Thin	15	15
	Normal	78	77
	Fat	8	8
	Total	101	100
Independent Efforts to Reduce Exposure	Frequent	26	25,7
	Sometimes	65	64,3
	Never	10	10
	Total	101	100

The age group of 46-55 years (Table 2) has a greater percentage among other categories of groups with 41.6% or as many as 42 respondents; farmers exposed to < 4 hours per day ie as many as 54 people or about 53.5%; then as many 58 or around 57.4% of respondents in Ijen

District have more than 10 years of work, 54 smokers or 53.4%, 78 people or 77% of farmers have normal nutritional status, 65 people or around 64.3% of farmers sometimes make independent efforts to reduce exposure to pollutants.

Environment Factors

Table 3: Temperature, Humidity, Ventilation.

Variable	Category	N	%
Temperature	Qualified	47	46,6
	Not qualified	54	53,4
	Total	101	100
Humidity	Qualified	47	46,6
	Not qualified	54	53,4
	Total	101	100
Ventilation	Qualified	90	89,1
	Not qualified	11	10,9
	Total	101	100

Environment factors (Table 3) as many as 54 houses or around 53.4% have temperatures that do not meet the requirements; as many as 54 or 53.4% of houses do not meet the humidity requirements and as many as 90

houses or 89.1% have ventilation of farmer houses that meet the requirements. Almost all houses in this study have the same model in terms of size and shape; there are only a few houses which are different

Bivariate Analysis

Table 4: Result of Logistic Regression.

Variable	Chi Square	Sig
SO ₂	61,559	0,000
Age	7,141	0,028
Length of Work	20,532	0,000
Smoking Habit	27,466	0,000
Duration of Exposure	47,096	0,000
Independent Efforts to Reduce Exposure	31,466	0,000
Nutritional Status	0,777	0,678
Temperature	65,044	0,000
Humidity	74,889	0,000
Ventilation	9,152	0,002

*it is regarded as significant if the p value < α (0.05)

The results of the analysis (Table 4) reveal that almost all variables are at the value of p value < α (0.05) so that it can be interpreted that they have a significant effect, except for nutritional status with a significance of 0.678 > α (0.05) so that it does not significantly affect the occurrence of URI in farmers in the watershed of Ijen District, Bondowoso.

URI examination conducted by doctors on farmers found that as many as 57.4% or 58 people were affected by light URI, while as many as 42.6% or 43 people were not affected by URI. The results of statistical test (Table 4) obtained a chi square value of SO₂ content of 61.559 which was greater than the value of chi square table 3.84 and p-value of 0.000 which was less than α (error level 0.05). These results indicate that SO₂ affects the occurrence of URI in farmers in Ijen District.

The individual factors (age, length of working, smoking habits, duration of exposure, independent efforts to

reduce exposure) and environmental factors (temperature, humidity, ventilation) have a chi square value greater than 3.84 and a p-value less than α (0, 05) are regarded to affect the occurrence of URI in farmers in Ijen District. The nutritional status variable was regarded not to affect the occurrence of URI disease in farmers in Ijen District because it had a p value of 0.678 greater than α (error level of 0.05) so that it did not meet the assumption of affecting the URI disease occurrence.

The average age of the farmer, specifically 51 years old, can be stated that it has come to the early elderly period. Decreased respiratory muscles and tissue elasticity will occur in line with age, then when breathing oxygen, the strength of the respiratory muscles will decrease, and it can make a person vulnerable to have URI exposure. At the age of 30 years, the lung function value will usually decrease slowly.^[5] The farmers exposed to SO₂ pollutant for more than 4 hours every day experience more light URI than the farmers who work less than 4 hours every

day. The duration of exposure is closely related to the length of time a farmer works in a day. Someone works usually 4 - 8 hours a day, if the working hours are longer, then it will have a negative impact on workers due to the exposure to these pollutants.^[6]

The average length of work of farmers affected by URI is > 10 years. The longer someone works, the more possible that person be contaminated with more pollutants.^[7] The accumulation of pollutants in the lungs has an effect on the narrowing of the respiratory tract, then the accumulation of accumulated dust causes the elasticity of the respiratory tissue to decrease so that the lung capacity also decreases.^[8] Smoking habit has a bad impact on the function and the structure of tissues in the respiratory tract. It has a close relation with the increased risk of respiratory disorders. Cigarette smoke causes the ability of vibrating hair in capturing foreign objects decreased which result in dust easily entering the lungs.^[9]

Independent efforts to reduce exposure (personal protective equipment) used by farmers are cuff, shoes, head coverings and cloth. However, not all farmers wear complete personal protective equipment, because they are less comfortable and it can interfere with their work productivity. Personal protective equipment is worn when the smell of SO₂ gas is extremely strong by using a wet cloth. There is no guarantee that someone using personal protective equipment will avoid the possibility of humans affected by respiratory disorders.^[10] The analysis results on the nutritional status of URI occurrence showed that there was no significant effect. There was no significant effect between nutritional status and respiratory disorders in textile industry workers X in Semarang Regency.^[11]

The research results found that the temperature condition of house that did not meet the requirements was at temperature less than 18oC. The condition of residential density in the house also affected the temperature in the room. Low temperature can cause an increase in relative humidity, so that it can increase the corrosive effects of pollutants in places where the air is polluted when in cold temperatures; the air condition becomes dense so that the pollution concentration in the air is higher.^[12] The average housing model is the same as the area that can be regarded as small and is habited of 3 or 4 people in one house in average. The more people in a room that has high humidity, the higher concentration of microorganism will be. Everyone has the possibility of containing microorganisms that has the risk of transmitting the disease to fellow residents of the room.

The results of the house ventilation found that in average it met the requirements, but this was not followed by people's behavior to open the window in the morning and also in the afternoon. It is because in the morning, SO₂ gas had started to cause a pungent odor so that the average respondent did not open the window. It is feared

that the pungent odor will become severe until it enters the house.^[13] explained that the concentration of germs is more in the un-exchanged air. It is closely related to the limited ventilation that functions to guarantee the quality and the adequacy of air circulation in and out of the room so that it is safe for breathing.

This is the first research conducted on exposure to natural SO₂ pollutants on farmers in Ijen Crater watershed, because the previous studies focus more on SO₂ exposure from industry. Then, this research combines two methods, namely SO₂ measurement and URI examination. The limitation of this study is that the URI examination is only done in a short time.

CONCLUSION

The sources of SO₂ exposure affect the occurrence of URI in farmers in Ijen District watershed. Individual factors (age, duration of exposure, length of work, smoking habits and independent efforts to reduce exposure) and environment factors (temperature, humidity, and ventilation) affect the occurrence of ARI in Ijen District watershed, Bondowoso. Besides, nutritional status does not affect the occurrence of URI. As a suggestion for this research, it is needed to conduct a research related to chronic respiratory disorder.

ACKNOWLEDGEMENT

Our sincerest appreciation is dedicated to PTP XII Blawan Bondowoso Regency who have been pleased to give permission to the implementation of this research.

REFERENCES

1. Aminuddin. Perekayasa di Pusat Sumber Daya Air Tanah dan Geologi Lingkungan, Badan Geologi, 2014.
2. Kementerian Kesehatan Republik Indonesia. Profil Kesehatan Indonesia. Jakarta Kementerian Kesehatan Republik Indonesia, 2014.
3. Badan Penanggulangan Bencana Daerah (BPBD). Bondowoso, 2018.
4. Firdaus, A P dan Sulistyorini L. Hubungan Antara SO₂ Dengan Kejadian ISPA Di Kota Surabaya 2013-2015. Jurnal.Fakultas Kesehatan Masyarakat Universitas Airlangga, 2017; 9(1).
5. Rahmatullah, P. Pneumonitis Dan Penyakit Paru Lingkungan. Edisi 5. Jakarta: Pusat Penerbitan Ilmu Penyakit Dalam, 2009.
6. Mengkidi, D N S. Gangguan Fungsi Paru dan Faktor-Faktor yang Mempengaruhinya pada Karyawan PT. Semen Tonasa Pangkep Sulawesi Selatan. Jurnal Kesehatan Lingkungan Masyarakat, 2009; 5(2).
7. Astorina N. Hubungan Masa Kerja terhadap Gangguan Fungsi Paru pada Petugas Penyapu Jalan di Protokol 3, 4 dan 6 Kota Semarang. Jurnal Kesehatan Lingkungan. Fakultas Kesehatan Masyarakat Universitas Diponegoro. 2015; 3(3).

8. Yusnabeti, Ririn, Ruth. PM10 dan Infeksi Saluran pernapasan Akut pada Pekerja Industri Mebel. Skripsi. Jakarta: Universitas Indonesia, 2011.
9. Suma'mur, P. K. Higiene Perusahaan Dan Kesehatan Kerja (Hiperkes). Jakarta: Sagung Seto, 2013.
10. Afiani E E. Faktor – Faktor yang Berhubungan dengan Gangguan Fungsi Paru pada Pekerja di Unit Boiler Industri Tekstil X Kabupaten Semarang. Jurnal. Fakultas Kesehatan Masyarakat Universitas Diponegoro, 2016; 4(3).
11. Siregar. Hubungan Tingkat Konsentrasi Sulfur Dioksida (SO₂), Total Suspended Particle (TSP) dan Lingkungan Fisik Dendan Jumlah Kejadian Infeksi Saluran Pernapasan (ISPA) Pada Penduduk DI Kotamadya Jakarta Timur. Skripsi. Universitas Indonesia, 2011.
12. Prajapati, B., Talsania, N., Sonaliya, K.N.” A Study on Prevalence of Acute Respiratory Infections (ARI) in Under Five Children in Urban and Rural Communities of Ahmedabad District, Gujarat”. National Journal of Community Medicine, 2011; 2(2).

